



Microwave-assisted Synthesis, Characterization, Antimicrobial and Antioxidant Activities of 1-Benzyl-3-[(4-Methylphenyl)imino]-Indolin-2-One and Its Co(II) Complex

Adebomi A. Ikotun^{1*} and Tolulope O. Omolekan²

¹Department of Chemistry and Industrial Chemistry, Bowen University, Iwo, Osun State, Nigeria.

²Department of Biochemistry, Bowen University, Iwo, Osun State, Nigeria.

Authors' contributions

This work was carried out in collaboration between the authors. Author AAI designed the work, carried out the chemistry and antimicrobial studies, while author TOO carried out the antioxidant studies and the reducing power assay. Both authors wrote the first draft, read and approved the final manuscript.

Article Information

DOI: 10.9734/CSJI/2019/v28i130130

Editor(s):

- (1) Dr. Ho Soon Min, Department of Chemistry, Faculty of Science, INTI International University, Malaysia.
(2) Thomas P. West, Ph.D. Professor and Head, Department of Chemistry, Texas A&M University-Commerce, Commerce, TX 75429, USA.

Reviewers:

- (1) Joel Omar Martínez, Universidad Autónoma de San Luis Potosí, México.
(2) Hasan Küçükbay, İnönü University, Turkey.
(3) Pipat Chooto, Prince of Songkla University, Thailand.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/51225>

Original Research Article

Received 07 July 2019
Accepted 09 September 2019
Published 17 September 2019

ABSTRACT

Aims: To develop a simple, efficient microwave-assisted synthetic method to prepare 1-Benzyl-3-[(4-methylphenyl)imino]-indolin-2-one and its Co(II) complex, characterize and ascertain their biological significance.

Place and Duration of Study: School of Chemistry, Cardiff University, Wales, United Kingdom (November, 2007); Department of Chemistry & Industrial Chemistry, Department of Biological Sciences and Department of Biochemistry, Bowen University, Iwo, Nigeria (September 2015 and June 2018).

Methodology: N-benzylisatin and 4-methylaniline were microwave irradiated in acetic acid to give 1-Benzyl-3-[(4-methylphenyl)imino]-indolin-2-one (C₂₂H₁₈N₂O, L; 58 %). L was microwave irradiated

*Corresponding author: E-mail: ikotunadebomi@gmail.com, lhbcards@yahoo.com;

with $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ in ethanol to yield its Co(II) complex. **L** was characterized using $^1\text{H-NMR}$, $^{13}\text{C-NMR}$, IR and Electronic spectra analyses, exact mass and melting point. IR and Electronic Spectra analyses with melting point confirmed the Co(II) complex was formed. The *in-vitro* antimicrobial activities of **L** were evaluated against three gram-positive bacteria (*Staphylococcus aureus*, *Bacillus subtilis* and Haemolytic *Staphylococcus aureus*), three gram-negative bacteria (*Pseudomonas aeruginosa*, *Escherichia coli* and *Klebsiella* sp.) and three fungi (*Aspergillus niger*, *Trichoderma viride* and *Penicillium citrinum*). The IC_{50} of **L** and its Co(II) complex were carried out against DPPH, H_2O_2 and NO radicals, as well as their reducing power abilities.

Results: Antimicrobial studies revealed **L** was active against *Pseudomonas aeruginosa* with a high zone of inhibition (thrice that of tetracycline) and *Penicillium citrinum*. The IC_{50} of **L** against DPPH, H_2O_2 and NO radicals were 0.561 ± 0.02 , 0.3 ± 0.01 and 0.53 ± 0.01 $\mu\text{g/ml}$ respectively, while they were 0.200 ± 0.01 , 0.9 ± 0.02 and 0.26 ± 0.03 $\mu\text{g/ml}$ for Co(II) complex. Their reducing power abilities at IC_{50} were 0.53 $\mu\text{g/ml}$ (for **L**) and 0.6 ± 0.03 $\mu\text{g/ml}$ (the complex).

Conclusion: **L** was synthesized within 15 min and its Co(II) complex within 5 min. **L** showed good free radical scavenging activities and reducing power when compared with ascorbic acid, while its Co(II) complex even performed better.

Keywords: Microwave-assisted; synthesis; antimicrobial and antioxidant activities.

1-Benzyl-3-[(4-methylphenyl)imino]-indolin-2-one;

1. INTRODUCTION

The use of microwave irradiation synthetic method has recently gained much attention due to the fact that it speedily facilitates the synthesis of new chemical entities with pharmaceutical applications by improving the efficiency of such chemical reactions [1,2]. Isatin is a versatile precursor in fine organic/inorganic syntheses of many amine-, ether-, nitrile- and oxazole-derivatives [3,4,5], as well as several metal complexes [6,7,8,9], with antimicrobial [10], antiviral, anti-inflammatory, analgesic [7,11] and anticonvulsant activities [12]. Quite an innumerable number of isatin derivatives have proven to be good therapeutic agents for several coronary diseases including ischemic heart disease, cardiac arrhythmia, hypertension, depression and even cancer [6]. The preparation and x-ray crystallographic structure of *N*-benzylindole-2, 3-dione (*N*-benzylisatin) has been reported [13]. *N*-alkylated isatins have also been reported to possess interesting chemistry and pharmacological activities such as antibacterial, antiviral and anticancer [14,15]. Hence the need to synthesize biologically active Schiff bases of *N*-benzyl isatin and their complexes using a simple and time efficient method and also establishing their biological significance are the propelling forces for this research. We have reported the single crystal structure of 1-Benzyl-3-[(4-methylphenyl)imino]-indolin-2-one grown after conventional refluxing of isatin and *P*-toluidine for 6 hours [16]. We hereby report the convenient microwave-assisted synthesis of 1-Benzyl-3-[(4-methylphenyl)imino]-

indolin-2-one (**L**) within 15 minutes, 30 seconds, its full spectroscopic characterization, antimicrobial studies, free radical scavenging activities and the reducing power assay. This report also includes the successful microwave irradiation synthesis (within 4 minutes, 55 seconds), free radical scavenging activities and the reducing power assay of its Co(II) metal complex.

2. EXPERIMENTAL DETAILS

2.1 Chemical

Isatin, *P*-Toluidine, $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 1,1-Diphenyl-2-picryl hydrazyl (DPPH), ascorbic acid, potassium dihydrogen phosphate, dipotassium hydrogen phosphate, sodium chloride, sodium nitroprusside and sulphanilamide were obtained from Aldrich. All solvents used (ethanol, methanol, chloroform, acetone, dichloromethane, pyridine, diethyl ether, *n*-hexane, and *N,N*-dimethylformamide; DMF) were purchased as analytical grades from Sigma-Aldrich and SAARChem.

2.2 Instrumentation

Microwave experiments were performed inside a domestic oven (24 L, 800W light-up Dial Microwave). $^1\text{H NMR}$ and $^{13}\text{C NMR}$ spectra were recorded at room temperature on Bruker 400 MHz Spectrometer. The infrared spectra were recorded on Agilent Cary 630 FTIR spectrometer. The UV-Visible spectra were recorded on a Shimadzu UV-1800 spectrometer.

Mass spectrum was determined using a Fisons VG Quattro Spectrometer. The purity of the compounds were checked by Thin-Layer Chromatography (TLC) on Silica Gel 60 F254 alumina plates (E Merck) using appropriate solvent mixtures of diethyl ether: petroleum ether or chloroform as the eluent and visualized in UV chamber (365 nm). Melting points were determined using a Gallenkemp variable heater apparatus.

2.3 Synthetic Work

2.3.1 Preparation of 1-benzyl-3-[(4-methylphenyl)imino]-indolin-2-one (L)

This was done in two steps.

2.3.2 Synthesis of *N*-benzylisatin (C₁₅H₁₁NO₂)

N-benzylisatin was prepared according to literature [13,16]. Isatin (1.00 g; 6.79 mmol) was dissolved in DMF (20 mL). NaH (0.16 g; 6.79 mmol) was added while stirring for 1 h. Benzyl bromide (0.81 mL; 6.79 mmol) was added and stirring was continued for another 24 h at room temperature. T.L.C. was done using chloroform as the eluent and the R_f value of the product was 0.95. The precipitated light yellow product, *N*-benzylisatin was filtered out under vacuum and recrystallized in ethanol. The yield after recrystallization was 0.52 g (32%).

2.3.3 Microwave synthesis of 1-benzyl-3-[(4methylphenyl)imino]indolin-2-one (C₂₂H₁₈N₂O; L)

N-Benzylisatin (1 g; 4.22 mmol) and *P*-toluidine (0.45 g; 4.22 mmol) were both dissolved in 20 mL acetic acid with heat. The mixture was poured inside a beaker and put inside a 24 L, 800W light-up Dial Microwave. This was irradiated for 7 minutes, 30 seconds (microwave medium; 149 - 177°C) and thereafter 8 minutes (microwave high; 218 -260°C) at 30 seconds intervals. A brown oil was got. This was recrystallized in ethanol and left to cool. A light orange solid 0.8 g (58%) was obtained, m.p. 152.4-154.6°C (154°C from Ikotun et al. 2012b); λ_{\max} (CH₃OH)/nm 417, 300, 250 (E/dm³mol⁻¹cm⁻¹) 3139, 4559, and 24914); ν_{\max} /cm⁻¹ 1716 (C=O), 1597 and 1683 (C=N + C=C), 1467 (CH_{bend}), 1100 and 1079 (C-N + C-C); δ H (400 MHz, DMSO-d₆)/ppm 5.00 (2H, s, CH₂), 2.35 (3H, s, CH₃), 6.55-7.70 (13H, m, ArH); δ C (DMSO-d₆)/ppm 162.90 (C=O), 154.23 (C=N), 148.18 (Cq), 147.30 (Cq), 136.38 (Cq), 134.86 (Cq),

134.60 (CH), 130.5 (CH), 129.25 (CH), 129.20 (CH), 129.15 (CH), 128.03 (CH), 127.84 (CH), 125.54 (CH), 122.87 (CH), 120.44 (CH), 117.95 (CH), 115.90 (CH), 111.10 (CH), 110.00 (CH), 21.06 (CH₃), 43.36 (CH₂); m/z (ESI) 327 (M+1⁺ peak, 100 %).

2.3.4 Synthesis of the Co(II) Complex

L (1 g; 0.00422 mol) was dissolved in 20 mL ethanol. CoCl₂.6H₂O (1 g; 0.00422 mol) was added to the solution. This was microwave irradiated (medium) for 4 minutes, 55 seconds at an interval of 30 seconds to yield a brown amorphous solid, 0.64 g (20%), m.p. > 320°C; λ_{\max} (CH₃OH)/nm 666, 619, 423, 310, 288 (15015, 16151, 23640, 32258 and 34722 cm⁻¹); ν_{\max} /cm⁻¹ 1698 (C=O), 1590 and 1614 (C=N + C=C), 1461 (CH_{bend}) and 1098 (C-N + C-C).

2.4 Antimicrobial Activity

The synthesized compounds were screened for *in-vitro* antibacterial and antifungal activities using Mueller-Hinton agar (MHA) and Potato dextrose agar (PDA) media. The antibacterial activity was evaluated against three gram-positive bacteria (*Staphylococcus aureus*, *Bacillus subtilis* and Haemolytic *Staphylococcus aureus*) and three gram-negative bacteria (*Pseudomonas aeruginosa*, *Escherichia coli* and *Klebsiella* sp.). Also, the antifungal activities of the compounds were evaluated against three fungi (*Aspergillus niger*, *Trichoderma viride* and *Penicillium citrinum*). Preliminary identification of the bacteria was carried out following the methods previously described by [17]. Tetracycline (30 µg; antibiotic test kit) was used as a standard drug for the bacteria, while dimethylformamide (DMF) was used as a control.

2.4.1 Antibacterial test

An 18 h culture of each test bacteria was suspended in a sterile universal bottle containing nutrient broth. Normal saline was added gradually in order to compare its turbidity to that of 0.5 McFarland standard corresponding to approximately 10⁸ cells/mL. This was diluted to produce 10⁶ cells/ml used for the experiment [18]. For the antibacterial susceptibility test, one milliliter (1 mL) of test organism (10⁶ cells/mL) was inoculated into Petri plates (90 mm diameter). 19 mL molten Mueller Hinton agar (MHA) sterilized at 121°C for 15 min was also added. The plates were shaken gently for even mixing. The agar was left on the bench to

solidify. Disc diffusion method was used to evaluate the antimicrobial activities of compounds. Filter paper discs were cut (diameter 8 mm) and sterilized. **L** was dissolved in dimethylformamide (DMF) at a concentration of 100 µg/mL. The discs were impregnated with the solution (100 µg/mL), picked with sterile forceps and placed on the MHA plates containing the test organisms [19]. The plates were left on the bench for 1 h to diffuse before incubating at 37°C for 24 h.

2.4.2 Antifungal test

The fungal isolates were allowed to grow on Potato dextrose agar (PDA) (LabM) at 25°C for 5 – 7 days to sporulate. After that, the fungal spores were harvested by pouring a mixture of sterile glycerol and distilled water unto the surface of the plate. The spores were scraped using a sterile glass rod. The harvested fungal spores were standardized to 10⁶ spores per mL. One milliliter of the standardized spore suspension was evenly spread on solidified PDA (LabM) plates using a glass spreader. The plates were placed on the work bench for 1 h for the spore suspension to diffuse into the agar. The sterile discs were impregnated with the test compounds and placed aseptically using sterile forceps on the surface of the agar plates. The plates were then allowed to stand on the laboratory bench for 1 h to allow proper diffusion of the compounds into the media. Plates were incubated at 25°C for 96 h and observed for zones of inhibition. Activity was evaluated by measuring the diameters of zones of growth inhibition in triplicates and the mean of three results determined.

2.4.3 Minimum inhibitory concentration (MIC)

This was carried out by adding 10, 5.0, 2.5, 1.25, 0.625, and 0.3125 µg/mL of **L** into test tubes containing sterile nutrient broth. *Pseudomonas aeruginosa* was thereafter introduced into the broths containing different concentrations of **L**. The tubes were then incubated for 24 h at 37°C. The MIC was taken as the lowest concentration of **L** that did not permit any visible growth [20].

2.5 Antioxidant Properties

The antioxidant properties determined were the free radical scavenging activities and the reducing power abilities of the synthesized compounds. These values were determined as a

mean ± standard deviation of three different readings.

2.5.1 Free radical scavenging activity and reducing power assays

All synthesized compounds were screened for *in vitro* free radical scavenging activities using DPPH, H₂O₂, nitric oxide (NO) radical scavenging activity assay, and reducing power assay.

2.5.2 DPPH radical scavenging activity

DPPH radical scavenging activity of the compounds was determined using the method according to the literature [21]. To 0.1 mL of different concentrations (0.1 to 1.0 µg/mL) of the test compounds, 2.5 mL of methanol and 0.5 mL of 0.2 mM DPPH solutions were added and mixed thoroughly and the absorbance was read at 517 nm against blank. Ascorbic acid was used as a reference standard. The IC₅₀ (Inhibitory concentration of the test compound required to scavenge 50% of DPPH free radicals) was thereafter calculated.

2.5.3 H₂O₂ radical scavenging activity

This was determined according to the literature [22,23]. The solution of hydrogen peroxide (20 mM) was prepared in phosphate buffered saline (pH 7.4). Various concentrations (0.1 to 1.0 µg/mL) of 1 mL of test compounds and standard were added to 2 mL of H₂O₂. Absorbance of H₂O₂ at 230 nm was determined 10 min later against a blank solution containing the phosphate buffered saline without H₂O₂. The IC₅₀ of H₂O₂ was thereafter calculated.

2.5.4 NO radical scavenging activity


Nitric Oxide generated from sodium nitroprusside in aqueous solution at physiological pH interacts with oxygen to produce nitrite ions which were measured by Griess reaction [24,25]. The reaction mixture (3 mL) containing sodium nitroprusside (10 mM) in phosphate buffered saline and test compounds in different concentrations was incubated at 25°C for 150 min. At intervals, samples (0.5 mL) of incubation solution were removed and 0.5 mL of Griess reagent (1% sulphanilamide, 2% H₃PO₄ and 0.1% naphthylethylene diamine dihydrochloride) was added. The absorbance of the chromophore formed was measured at 546 nm.

2.5.5 Reducing power


The reducing power of the test compounds was carried out according to literature [21] (Bliss 1958). 2.5 mL of 0.2 M phosphate buffer (pH 6.6) and potassium ferric cyanide (1% w/v) were added to 1.0 mL of the sample dissolved in DMF. The resulting mixture was incubated at 50°C for 20 mins, followed by the addition of 2.5 mL trichloroacetic acid (10% w/v). The mixture was centrifuged at 3000 rpm for 10 min to collect the upper layer of the solution (2.5 mL) mixed with distilled water (2.5 mL) and ferric chloride (0.1% w/v), the absorbance was then measured at 700 nm against blank sample.

3. RESULTS AND DISCUSSION

The reaction for the microwave synthesis of **L** from *N*-benzylisatin is presented as Scheme 1.



Scheme 1. Microwave synthesis of 1-benzyl-3-[(4-methylphenyl)imino]-indolin-2-one ($C_{22}H_{18}N_2O$; **L) from the prepared *N*-benzylisatin**



Scheme 2. Microwave synthesis of the cobalt(II) complex

Scheme 2 presents the microwave synthesis of the Cobalt(II) complex similar to literature [10]. The metal has picked up two molecules of the ligand making the reaction ratio to be 1:2 (Metal : Ligand).

3.1 Infrared Spectra of **L** and Its Co(II) Complex

The characteristic vibrational frequencies in the infrared spectra have been identified by comparing the spectrum of the Schiff base with the Co(II) complex as shown in Table 1.

The assignments of these absorption bands have also been made by comparing the spectra of the compounds with reported literature on transition metal complexes of isatin Schiff bases [6,10]. There are two potential donor sites in **L** including isatin. These are the isatin nitrogen and the isatin

Table 1. Relevant infrared spectral data of the L and its co(II) complex

Compound	$\nu(\text{C=O})$ (cm^{-1})	$\nu(\text{C=N+C=C})$ (cm^{-1})	$\nu(\text{CH})_{\text{bend}}$ (cm^{-1})	$\nu(\text{C-N+C-C})$ (cm^{-1})
L ($\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$)	1716 s	1683 s 1597 m	1467 s	1100 m 1079 m
Co(II) Complex	1698 s	1614 s 1590 m	1461 s	- 1098 m

Note: *ν* stretching; *m*, medium and *s*, strong

oxygen. The FTIR spectrum of **L** ($\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$) showed a strong intensity band at 1716 cm^{-1} attributed to $\nu(\text{C=O})$ stretching vibration. This band has undergone a shift to a lower frequency of 1698 cm^{-1} in the spectrum of its Co(II) complex signifying the involvement of the keto oxygen in coordination to Co(II). The spectrum of **L** showed a strong and a medium band at 1683 and 1597 cm^{-1} attributed to $\nu(\text{C=N} + \text{C=C})$ stretching vibration. These bands have moved to lower frequencies of 1614 and 1590 cm^{-1} . This also signifies the involvement of the imine nitrogen in coordination to Co(II). The strong band appearing at 1467 cm^{-1} in the spectrum of **L** was attributed to the $\nu(\text{CH}_{\text{bend}})$ vibration and it has shifted slightly to a lower frequency of 1461 cm^{-1} in the spectrum of its complex. The medium bands appearing at 1100 and 1079 cm^{-1} in the spectrum of **L** were attributed to $\nu(\text{C-N} + \text{C-C})$ stretching vibration. One of these bands has disappeared, while the other appeared at 1098 cm^{-1} in the spectrum of the Co(II) complex. All confirm the formation of the Co(II) complex of this ligand.

3.2 ^1H - and ^{13}C -NMR Spectra of L ($\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$)

Fig. 1 presents the numbering system of atoms in **L**.




Fig. 1. The numbering system of atoms in L ($\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$)

In the ^1H -NMR spectrum of **L**, a singlet at δ 5.00 ppm appeared, which is due to the methylene group. The singlet that appeared at 2.35 ppm is due to the methyl group. The aromatic protons appeared as groups of multiplets in the range 6.55-7.70 ppm. In the ^{13}C -NMR spectrum (DMSO- d_6) of this compound, the expected 22 signals were observed as follows (δ ppm): 162.90 (C=O), 154.23 (C=N), 148.18 (Cq), 147.30 (Cq), 136.38 (Cq), 134.86 (Cq), 134.60 (CH), 130.5 (CH), 129.25 (CH), 129.20 (CH), 129.15 (CH), 128.03 (CH), 127.84 (CH), 125.54 (CH), 122.87 (CH), 120.44 (CH), 117.95 (CH), 115.90 (CH), 111.10 (CH), 110.00 (CH), 21.06 (CH_3). The peak that appeared at 43.36 ppm is the characteristic of the CH_2 group.

3.3 Mass Spectrum of L ($\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$)

Exact mass analysis of **L** showed the elemental composition to be $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$, which corresponds to the expected molecular formula and revealed the molecular ion peak at m/z (ESI) 327 (100%) corresponding to $[\text{M}+\text{H}]^+$. This ion fragments by the loss of $\text{C}_7\text{H}_{14}\text{O}$ (114 mass units) to give the ion at m/z 213 (18%) corresponding to $[\text{M}+\text{H}+\text{C}_6\text{H}_6+\text{CH}_4+\text{H}_2\text{O}+\text{H}_2]^+$ or $[\text{M}+\text{H}+\text{C}_7\text{H}_8+\text{H}_2\text{O}+2\text{H}_2]^+$. The ion at m/z 213 fragments by the loss of 9 mass units corresponding to 4H_2 and a proton (H) to give the ion at m/z 204 (5%). The spectral fragmentation is presented in Table 2, while the fragmentation pattern is presented as Scheme 3.

3.4 UV-Visible Spectra

The ultraviolet spectra analyses of the prepared compounds are presented in Table 3.

The absorption bands have been assigned by comparing the spectra of the compounds with reported literature on transition metal complexes of isatin Schiff bases [6,10]. The spectrum of **L** showed absorption bands at 33,333 and $40,000\text{ cm}^{-1}$ (300 and 250 nm) which have been assigned to $\pi - \pi^*$ transition. The band appearing at $23,981\text{ cm}^{-1}$ (417 nm) has been assigned to $n - \pi^*$ transition. The Co(II) complex

was characterized with bands appearing at 15,015 and 16,151 cm^{-1} (666 and 619 nm), which have been assigned as d-d transitions [6,10]. The Co(II) complex spectrum also showed a band at 23,640 cm^{-1} (423 nm) assigned to $n-\pi^*$ transition, as well as bands appearing at 32,258 and 34,722 cm^{-1} (310 and 288 nm) assigned to $\pi-\pi^*$ transitions.

3.5 Antimicrobial Activities

Fig. 2 shows the comparison of the results of the antibacterial studies of L and tetracycline as a bar chart. The compound was only active against *Pseudomonas aeruginosa*, which is a very drug resistant bacterium that causes wound infections including gangrin [26,27].





Fig. 2. Comparative results of the antibacterial activities of L and tetracycline

Note Keys: Bac = *Bacillus subtilis* (+); SA = *Staphylococcus aureus* (+); HAS = *Haemolytic Staphylococcus aureus* (+); L = $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$ (L); Kleb (-) = *Klebsiella* sp.; PAE (-) = *Pseudomonas aeruginosa*; EC (-) = *Escherichia coli*; Tetra = Tetracycline

Table 2. Mass spectral data of L ($\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$)

m/z	Fragment
327	$[\text{M}+\text{H}]^+$
213	$[\text{M}+\text{H}+\text{C}_6\text{H}_6+\text{CH}_4+\text{H}_2\text{O}+\text{H}_2]^+$ or $[\text{M}+\text{H}+\text{C}_7\text{H}_8+\text{H}_2\text{O}+2\text{H}_2]^+$
204	$[\text{M}+\text{H}+\text{C}_6\text{H}_6+\text{CH}_4+\text{H}_2\text{O}+5\text{H}_2+\text{H}]^+$ or $[\text{M}+\text{H}+\text{C}_7\text{H}_8+\text{H}_2\text{O}+6\text{H}_2+\text{H}]^+$



Scheme 3. Fragmentation pattern for L ($\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}$)

Table 3. Electronic spectra of L and its co(II) complex

Compound	Band position (nm)	Band position (cm ⁻¹)	Band assignment
L	417	23,981	n -π*
	300	33,333	π -π*
	250	40,000	π -π*
Co(II) Complex	666	15,015	d-d
	619	16,151	d-d
	423	23,640	n -π*
	310	32,258	π -π*
	228	34,722	π -π*

L had a high zone of inhibition against this bacterium, which was thrice that of tetracycline. Thus it can be developed as a drug against *Pseudomonas aeruginosa*. Fig. 3 also presents the results of the antifungal studies of L as a bar chart.

It revealed that L was active against one of the tested fungi (*Penicillium citrinum*) and could be a potential drug against it.

3.6 Antioxidant Activities

The results of the antioxidant activities are presented below.

3.6.1 DPPH radical scavenging activity


Fig. 4 presents the comparison of the results of DPPH radical scavenging activities of L, its Co(II) complex and ascorbic acid as a bar chart.

DPPH mimics many free radicals produced in the biological system which have a stable but highly delocalized spare electron. The compounds under study are not readily broken down to less

reactive species. They either form dimers or attack macromolecules in the biological system which leads to abrogated cellular function, carcinogenesis, aging or cell death. However, they become less reactive when an antioxidant donates hydrogen atom to their molecules or when they form chelates. The Cobalt(II) complex had a lower IC₅₀ value compared to L and ascorbic acid. This shows the potency of this metal complex in scavenging a charged and highly reactive radical like DPPH in a biological system. It is most likely that the positively charged cobalt ion in this metal complex molecule readily chelates the nitrogen donor (DPPH molecule) to prevent its attacks on macromolecules in a biological system. Therefore, the Co(II) metal complex possesses better DPPH radical scavenging activity than its ligand (L) and ascorbic acid.

3.6.2 H₂O₂ radical scavenging activity

Fig. 5 presents the comparison of the results of H₂O₂ radical scavenging activity for L, its Co(II) complex and ascorbic acid as a bar chart.

**Fig. 3. Results of the antifungal studies of L**




Fig. 4. IC₅₀ values of test compounds for DPPH radical scavenging activity




Fig. 5. IC₅₀ values of test compounds for H₂O₂ radical scavenging activity

Hydrogen peroxide is a very reactive and unstable radical which the human body produces during infection. It rapidly decomposes into a more reactive but relatively stable hydroxyl radical (OH[•]). Both of them combat the invading microorganisms and eventually kill it. They initiate lipid peroxidation, DNA mutation and oxidative stress in the microorganism which eventually lead to their death. Although the human body has its mechanism of detoxifying these radicals, their sustained production after the death of the microorganisms could lead to similar effects in the human body. **L** showed higher potency in scavenging hydrogen peroxide than its Co(II) complex. This means that **L** is highly sensitive to hydrogen peroxide at a very low concentration. The higher hydrogen peroxide scavenging activity of **L** can be explained by the

presence of the carbonyl and imino groups in its molecule, which show high sensitivity to H₂O₂. However, the metal complex has both the oxygen of the carbonyl group and the imine nitrogen involved in coordination to the Co(II) ion. This is a possible reason for its reduced ability in scavenging hydrogen peroxide. And this could invariably account for its bactericidal effect by sustaining the hydrogen peroxide oxidation of macromolecules in the bacterial cell which could lead to the death, inhibition of growth or reduced population of the bacterial cell.

3.6.3 NO radical scavenging activity

Fig. 6 presents the comparison of the results of NO radical scavenging activities for **L**, its Co(II) complex and ascorbic acid as a bar chart.

NO is readily oxidized to a peroxy nitrite radical (ONOO[•]), which is a more stable radical in the presence of oxygen. However, protonation of peroxy nitrite radical gives a highly reactive peroxy nitrous acid (ONOOH) which has strong oxidizing properties toward various cellular constituents and can cause cell death, lipid peroxidation, carcinogenesis and aging [28]. An antioxidant compound which could scavenge rather than donating a hydrogen atom would be necessary to detoxify NO[•]. This prevents activating it to more cytotoxic radicals. The Co(II) complex at a low concentration (0.26 µg/mL) has shown that it possesses the capacity of scavenging NO[•]. It is probably capable of coordinating NO[•] as a ligand to form a stable

chelate in a ring structure rather than donating hydrogen atom to NO[•]. This complex is therefore capable of reducing NO[•] to a non toxic end product. The comparison of IC₅₀ values of Co(II) complex with L and ascorbic acid shows that Co(II) complex has a better mechanism of scavenging NO[•] radical. This is because the Co(II) ion present in this complex is capable of reducing NO[•] to a non toxic metabolite.

The reducing power of an antioxidant is its ability to abstract or receive electrons from a free radical to form a more stable compound. Fig. 7 therefore shows the comparison of the results of the reducing power activities of the tested compounds.




Fig. 6. IC₅₀ values of test compounds for NO radical scavenging activity




Fig. 7. IC₅₀ values of test compounds for reducing power activity of the test compounds

These results revealed that **L** and its Co(II) complex have very effective reducing power than ascorbic acid. Thus they could scavenge negatively charged free radicals at very low concentrations compared with ascorbic acid the reference antioxidant compound. Also, the Co(II) ion in the metal complex is capable of receiving the lone pairs of electrons on free radicals into its empty d-orbitals, thereby forming a new stable and harmless complex. This serves as an explanation for its effective reducing power.

4. CONCLUSION

Microwave irradiation technique has been successfully developed for the synthesis of 1-Benzyl-3-[(4-methylphenyl)imino]-indolin-2-one (**L**) and its Co(II) complex. The *in-vitro* antibacterial activities of **L** evaluated against three gram-positive bacteria (*Staphylococcus aureus*, *Bacillus subtilis* and Haemolytic *Staphylococcus aureus*) and three gram-negative bacteria (*Pseudomonas aeruginosa*, *Escherichia coli* and *Klebsiella* sp.) revealed it was active against *Pseudomonas aeruginosa* with a very high zone of inhibition, about thrice that of tetracycline (clinical drug). The *in-vitro* antifungal activities of **L** evaluated against three fungi (*Aspergillus niger*, *Trichoderma viride* and *Penicillium citrinum*) revealed it was active against *Penicillium citrinum*. 1-Benzyl-3-[(4-methylphenyl)imino]-indolin-2-one has also proven to be a good antioxidant compound through its potent reducing ability, nitric oxide and DPPH radical scavenging activities. Test results showed it could scavenge relatively stable free radicals before they decompose into unstable and highly reactive radicals. However, its bactericidal effect might be through its hydrogen peroxide sparing effects. Hydrogen peroxide is a relatively unstable radical which easily decomposes to hydroxyl radical and finally oxygen and water. If hydrogen peroxide is spared in a living system, it is capable of initiating oxidative stress, DNA damage, abrogation of cellular functions and eventually cell death. As expected, the Co(II) complex has proven to be a better antioxidant compound than its ligand **L** and ascorbic acid through its potent reducing ability, nitric oxide and DPPH radical scavenging activities. It has a bi-dimensional approach to its activity: it confers protection to humans while being toxic to bacteria. However, it might be genotoxic which encourages involving research for further investigations.

ACKNOWLEDGEMENTS

The authors appreciate the kind assistance of Dr. Abimbola A. Owoseni towards the success of the antimicrobial studies. We are immensely grateful to the Royal Society of Chemistry (RSC) for supporting this research with funds.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Wagner RW. Microwave-assisted synthesis in the pharmaceutical industry a current perspective and future prospects. *Drug Disc. World.* 2006;7:59–66.
2. Krstenansky JL, Cotterill I. Recent advances in microwave-assisted organic syntheses. *Curr. Opin. Drug Discovery Dev.* 2000;3:454–461.
3. Abele E, Abele R, Dzenitis O, Lukevics E. Indole and isatin oximes: Synthesis, reactions and biological activity. *Chem Heterocycl Comp.* 2003;39:3-35.
4. Zhou F, Liu YL, Zhou J. Catalytic asymmetric synthesis of oxindoles bearing tetrasubstituted stereocenter at the C-3 position. *Adv Synth Catal.* 2010;352:1381-1407.
5. Nathani BR, Pandya KS, Jeni MM, Prasad A. Synthesis, characterisation and antimicrobial study of some new Isatins derivatives. *J Chem Pharm Res.* 2011;3: 687-692.
6. Chohan ZH, Shaikh AU, Naseer MM. Metal-based isatin-bearing Sulfonamides: Their synthesis, characterization and biological properties. *Appl Organomet Chem.* 2006;20:729-739.
7. Cerchiaro G, Ferreora AMDC. Oxindoles and copper complexes with oxindole-derivatives as potential pharmacological agents. *J Braz Chem Soc.* 2006;17:1473-1485.
8. Khan BT, Najmuddin K, Shamsuddin S, Zakeeruddin SM. Mixed ligand complexes of cis-dichloroethionine palladium (II) with purines, pyrimidines and nucleosides. *Inorg Chim Acta.* 1990;170:129-131.
9. Singh RV, Fahmi N, Biyala MK. Coordination behavior and biopotency of N and S/O donor ligands with their

- Paladium(II) and Platinum(II) complexes. *J Iranian Chem Soc.* 2005;2:40-46.
10. Ikotun AA, Egharevba GO, Obafemi CA, Owoseni AO. Ring deactivating effect on antimicrobial activities of metal complexes of the Schiff base of *p*-nitroaniline and isatin. *J Chem Pharm.* 2012;4:416-422.
 11. Jarrahpour A, Khalili D, De Clercq E, Salmi C, Brunel JM. Synthesis, antibacterial, antifungal and antiviral activity evaluation of some new bis-schiff bases of Isatin and their derivatives. *Molecules.* 2007;12:1720-1730.
 12. Verma M, Pandeya SN, Singh KN, Stables JP. Anticonvulsant activity of Schiff bases of isatin derivatives. *Acta Pharm.* 2004;54:49-56.
 13. Akkurt M, Turktekin S, Jarrahpour A, Khalili D, Buyukgungor O. *N*-Benzylindole-2,3-dione (*N*-benzylisatin). *Acta Cryst.* 2006;E62:1575-1577.
 14. Bauer DJ, Sadler PW. The structure-activity relationships of the antiviral chemotherapeutic activity of isatin β -thiosemicarbazone. *Brit J Pharmacol.* 1960;15:101-110.
 15. Rainoldi G, Lesma G, Picozzi C, Lo Presti L, Silvani A. One step access to oxindole-based β -Lactams throug Ugi four-center three-component reaction. *RSC Adv.* 2018;8:34903-34910.
 16. Ikotun AA, Adelani PO, Egharevba GO. 1-Benzyl-3-[(4methylphenyl)imino]-indolin-2-one. *Acta Cryst.* 2012;E68:2098.
 17. Cheesbrough M. Biochemical tests to identify bacteria. In: *Laboratory practice in tropical countries.* Cheesbrough M, (Ed.). Cambridge Edition. 2002;63-87.
 18. NCCLS (National Committee for Clinical Laboratory Standards). *Methods for dilution in antimicrobial susceptibility tests: Approved Standard M2-A5.* Villanova PA, NCCLS; 1993.
 19. Emeruwa KC. Antimicrobial substances from *Cacaya papaya* fruit extracts. *J Nat Prod.* 1982;45:123-127.
 20. NCCLS (National Committee for Clinical Laboratory Standards). *Methods for dilution of antimicrobial susceptibility tests of bacteria that grow aerobically. Approved Standard M100-S12.* Wayne PA, NCCLS; 2002.
 21. Blios MS. Antioxidant determination by the use of stable free radical. *Nature.* 1958;26:1199.
 22. Sanchez M. Methods used to evaluate the free radical scavenging activity in foods and biological systems. *Food Science Technology Institute.* 2001;8:121-137.
 23. Famey EJC, Luyengi L, Lee SK, Zhu BN, Zhou HHS. Prog, Antioxidant flavonoid glycosides from *Daphniphyllum calycium*. *J Nat Prod.* 1998;61:706-708.
 24. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JK, Tannenbaum SR. Analysis of nitrate, nitrite and 15N nitrate in biological fluids. *Anal Biochem.* 1982;126:131-138.
 25. Marcocci LMJJ, Droy-Lefaix MT, Packer L. The nitric oxide scavenging property of *Ginkgo biloba* extract EGb 761. *Biochem Bioph Res Co.* 1994;201:748-755.
 26. Hancock REW, Speert DP. Antibiotic resistance in *Pseudomonas aeruginosa*: Mechanisms and impact on treatment. *Drug Resist Updat.* 2000;3:247-255.
 27. Lutz JK, Lee J. Prevalence and antimicrobial-resistance of *Pseudomonas aeruginosa* in swimming pools and hot tubs. *Int J Environ Res Public Health.* 2011;8:554-564.
 28. Alam MN, Bristi NJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J.* 2013;3(21):143-152.

© 2019 Ikotun and Omolekan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
 The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/51225>