

Asian Journal of Fisheries and Aquatic Research

Volume 26, Issue 3, Page 1-11, 2024; Article no.AJFAR.113570 ISSN: 2582-3760

The Utilization of IoT Using Grow My Fish App in African Catfish (*Clarias sp*) Cultivation Employing Biofloc Technology in Apung Village, Bulungan Regency, North Kalimantan Province

Tesya Despira ^{a*}, Gatot Hari Priowirjanto ^b, Rita Rostika ^b, and Mochhamad Ikhsan Cahya Utama ^a

^a Tropical Marine Fisheries Study Program, Off-Main Campus Study Program (PSDKU) Pangandaran, Padjadjaran University, Indonesia.

^b Fisheries Study Program, Fisheries and Marine Sciences Faculty, Padjadjaran University, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2024/v26i3741

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/113570

Original Research Article

Received: 04/01/2024 Accepted: 08/03/2024 Published: 11/03/2024

ABSTRACT

This research aims at introducing the Grow My Fish application (or app) as an IoT for collecting and processing data on water quality in biofloc tank for cultivators. IoT in itself has been widely used by several sectors, and one of them is the fisheries sector. Grow My Fish is an app specifically designed to support research in the fisheries field. This research was conducted in Apung Village, Tanjung Selor District, Bulungan Regency from October through December 2023. Biofloc is an innovation in fisheries field. It is known as an eco-friendly innovation thanks to its

Asian J. Fish. Aqu. Res., vol. 26, no. 3, pp. 1-11, 2024

^{*}Corresponding author: Email: tesya20002@mail.unpad.ac.id;

ability to process the wastes produced by fish into natural feed in the assistance of the probiotic that was mixed into the water. However, the persistent main issue until recently in this biofloc innovation is water quality. Many cultivators so far recorded their water quality measurements on a book. This made it a little hard for these cultivators to process the data. For this reason, an IoT that facilitated cultivators in processing the data on water quality, mortality, fish weight, and amount of feed was born.

Keywords: African catfish; biofloc; IoT; grow my fish; water quality.

1. INTRODUCTION

The fisheries sector has given birth to an ecofriendly innovation, namely fish cultivation employing biofloc technology. Making a good use beneficial microorganisms. this of biofloc innovation is well-known among cultivators for its effectiveness and efficiency in increasing the harvests. Biofloc consists of 2 words, namely "Bio" which means life and "Floc" which means a lump. From these two words, it can be defined as a group of various organisms such as bacteria, fungi, protozoa, and algae in one lump [1]. Biofloc also utilizes oxygen and microorganisms that can be turned into fish feed in the tank. The oxygen in the biofloc cultivation is supplied through the aerator and it needs to be constantly turned on for 24 hours, to prevent any sedimentation. When sedimentation occurs, farmers will need to drain the tank manually [2,3].

Excessive feeding leads the toxic ammoniac to deposit, resulting in unsafe environment for the fish. The cultivation employing biofloc technology produces microorganisms from probiotic to decompose wastes and generate feed raw materials for the cultivated fish.

One freshwater fish many cultivators are fans of is African catfish. In addition, the African catfish is still continuously developed by the government to increase the fisheries sector production [4,5]. The problem is that the successful cultivation of African catfish with biofloc technology is affected water quality parameters such by as temperature, pH, and DO (Suprivanto et al., 2019) in Ilham Firman Ashari et al. [6]. Poor water quality will negatively affect the fish growth. This is because when the water quality is poor, the floc will not grow or form and it eventually affects the fish. This will lower the fish's appetite because of the water's strong odor as a result of ungrown floc [7-9]. The worst consequence is that it can lead to the fish's mass death. The water quality is poor for fish cultivation with biofloc technology when the temperature is 2530°C, pH is 6.5–8, DO is >3 and the water is greenish (SNI 6484.3:2014). Additionally, fish cultivators still use books to record their water quality measurement results so far. This makes it hard for them to process the data they have obtained.

Currently, Indonesia is one of those countries that has entered the 4.0 era, in which Artificial Intelligence (AI) and internet of Things (IoT) have begun to be widely used. In this era. technologies have been created to help cultivators in their attempt to achieve food security [10,11]. One of these technologies is Grow My Fish, an app designed to support research in fisheries field to facilitate data processing. Using this Grow My fish app, cultivators can simply input the data on water quality, fish weight, amount of feed, fish mortality, and farmer's observation data and market real time.

This research aims at introducing the Grow My Fish app as a means to collect and process data while cultivating fish. In addition, it aims at discovering the effectiveness of data collection and processing when using the manual method and Grow My Fish app.

2. METHODS

The research was conducted in Apung Village, Bulungan Regency, North Kalimantan Province, from October through December 2023. It began with making the biofloc tank and this was assisted by students and employees from Borneo Tarakan University. The biofloc tank was a circle of 3-meter diameter in shape, made of tarpaulin with an iron frame. In this project, the research participated in the land processing, tank assembly, and even pavement construction. The method used involved collecting the data on water quality and inputting them into the Grow My Fish app. This made it easier for data processing and the data would be permanently saved in the app. Before using the Grow My Fish app to collect the water quality data, the app needed to be installed first in the mobile phones or notebooks of each cultivator. This Grow My Fish app had been available on Play Store. The steps to install it are as follows:

First, open Play Store then click the search bar and write Grow My Fish on it. Click install Afterwards, login to the app in the login menu. The only problem is that this application can only be used if the cultivator has registered with PT SCM as the owner of this app. The registration was intended to let the company know what institution was the cultivator from and to obtain an account.




Fig. 1. Grow my fish app



Fig. 2. Login menu appearance

After logging in as cultivator, a dashboard will pop up. There, the cultivator could complete such data as the tank, cultivation activity, farmer's survey, market survey and also mini games as an entertainment to let the cultivator rest while having free time.

Next, go to the cultivation menu to input your cultivation data. In addition, the cultivator can also input the data from their monitoring of water quality, amount of feed, fish weight, and mortality. In this cultivation menu, cultivators can input the data on their tanks such as the cultivation method (biofloc, RAS, Torrential current), fish species, number of fish stocked,

stocking/restocking date and estimated fish harvest. In addition to going straight to the app, the data will automatically be inputted to Excel sheet and can be immediately downloaded to make things easier for the cultivator.

The data from the monitoring can be inputted every day and repeatedly. In addition, if the water quality in the tank is poor and the cultivator takes an action, the cultivator can also inputted the data regarding the actions taken into the Grow My Fish app. Thus, when what this cultivator did solved the problem, this data can be used as a reference and it can also help other cultivators in dealing with the same problems.








Fig. 3. Dashboard menu



Tanggal Awal 8 Oktober 2023 18 Desember 202					
*		1		Hardnes	
9(12)2023 02:45	6.5	47.2. %C	24 mgT.	Q ang L	
n 12/2023 16:50	7	27.5 °C	25.4 mg/L	O ang L	
6(12:2023	6.4	29.7 *0	24.3 mgL	0 mgT.	
612:3023 00:00	6.6	25.8 °C	$26.3 \mathrm{mg} \mathrm{L}$	O may'E.	
5(12)2023 05:00	6.2	264.ºC	26.3 mgL	0 mg/L	
4/12/2023 17:68	6.6	27.4 °C	27.6 mgL	0 mg/L	
012/2023 (0:00	6.1	26.9 °C	27 mg L	0 mg/L	
3(12)2021 17.00	6.5	28.5 °C	29.3 mg/L	0 mg/1.	
2;12:2023 17:00	6.6	33.50	23 mg/L	0 mg/L	
212/2023	65	28.5 °C	32.0 mg L	0 mg/L	

Fig. 4. Input of data from monitoring

In addition to the data on water quality, the cultivator can also input the data regarding the amount of feed used everyday, the fish average weight, and the fish mortality. This app is used not only by cultivators who raise the fish, but also by those who keep the fish for seeding and nurserv purposes. And, of course. the available menu will also differ. However, both share the same main issue, i.e., water quality monitoring. Other than inputting the cultivation data, the cultivator can also input data on the tank they used, including the tank type and land area.

If the cultivators want time series data, then they need to go to the dashboard menu on the data processed by the users. The monitoring data chart can only be seen by the users.

3. RESULTS AND DISCUSSION

3.1 Collected Data

The water quality data were important for cultivators, since they could be used for the cultivation in the next cycle. Furthermore, the quality needed to be checked on a regular basis. In this research, the research checked the water quality every morning and late afternoon on daily basis. In addition, the researcher also sampled the fish once every 10 days. This was to figure out the fish weight to recalculate the amount of feed that needed to be given later.

3.1.1 Data on pH

The good pH for cultivating African catfish using biofloc technology was 6.5–8 (SNI 6484.3:2014). The neutral pH was 6–7. Hence, a pH <6 could be interpreted as acidic and >7 was considered basic. An action needed to be immediately taken when the water pH decreased or increased. Less optimal pH might make the fish stressed, more vulnerable to disease, and could reduce its growth and productivity (Diansari et al, 2013) in Safsafubun et al. [12].

From the chart above, it could be concluded that during the cultivation, the water pH was still in compliance with SNI. However, some days witnessed that the water pH was >8, thus the researcher took a measure of putting two Catappa leaves into the tank. The chart was the data that had been processed from the ones that the researcher inputted in the Grow My Fish app.

3.1.2 Temperature data

Other than pH, temperature is also highly important in fish cultivation. Temperature is the controlling factor that can affect the digestion and metabolism in the fish body. Changes in temperature typically occurred during season transition. This has the potential of making it difficult for the fish to digest its food. This was also the case in Apung Village. The season in this research site kept on changing. The change occurring in the research site was that it would rain for one week and dried the next. This made the fish growth varied as a result of its difficulty in digesting the food.




Fig. 5. A chart from processed pH data

Temperature	Feed consumption response	
Close to 0	Minimum crisis condition	
8–10	No response to feeding	
15	Decreased feeding	
22	50% optimum	
28–30	Decreased feeding	
33	50% optimum	
35	Decreased feeding	
36–38	No response to feeding	
38–42	Minimum crisis condition	

Table 1. Effect of temperature on African catfish's feed consumption response [13]




Fig. 6. Temperature chart

From Table 1, it could be interpreted that African catfish in biofloc tank would live better at warm temperature ranging from 25°C to 30°C. It was imperative to maintain the temperature to prevent it from significantly increasing more than 4°C since this would lead to the fish death and stress.

From Fig. 6, it can be seen that the high temperature was recorded at 37°C, yet it occurred only in one day and led to the death of 25 fish. The chart was obtained from the data that had been processed from the Grow My Fish app.

3.1.3 DO (Dissolved oxygen) data

DO is one factor that affects the water quality in biofloc technology. The amount of oxygen dissolved in water is important to keep the water conducive for aquatic organisms. The DO concentration in water can be affected by some factors such as water temperature, atmospheric pressure, biological activities and pollution. Low dissolved oxygen can render the decomposition, reproduction and growth processes in the tank less optimal and it can eventually lead to death for the fish. The good oxygen level for a biofloc tank is >3 mg/l. To stabilize the dissolved oxygen, an aerator is used. Aerator itself is a tool to help increase the oxygen value in the water, to allow more oxygen to get into the water.

Fig. 7 showed that the dissolved oxygen still frequently increased drastically. This made the death of fish inevitable in this research. The chart above was the processed data from the Grow My Fish app.

In addition to separate data, this app also the combined water quality chart. This allowed us to see the comparison. Fig. 8 shows the combined chart from the water quality parameter data. From the chart above, it can be seen that some

parts declined steeply. This was the data on the day where the water quality was not controlled.

3.2 Other Parameter Data (Fish Weight, Amount of Feed, Mortality)

The collected data were not limited only to water quality. In the Grow My Fish app, the researcher also inputted such data as fish weight, amount of feed and mortality. In addition to water quality, the cultivator should also own data on fish weight for every sampling, amount of feed used and the fish mortality in one cycle. These data could be inputted by the cultivator to the Grow My Fish app. This allowed the cultivator to readily discover the FCR and losses they might suffer from. Yet, in these parameters, the data had not been processed as a chart.

3.2.1 Data on amount of feed

The feed data were calculated whenever a sampling was carried out. This sampling itself was done once every 10 days. Other than to discover the amount of feed, the sampling was also aimed at finding out the fish growth. To feed the African catfish, the researcher used FR 3%. From the data in Fig. 9, the amount of feed for every 10 days would increase by 100 gr. And this could be interpreted that the fish also grew.

3.2.2 Fish weight data

The fish weight data were also obtained after the sampling. The sampling was done once every 10

days. From Fig. 10, it can be seen that the fish weight increased every 10 days. These were the data on fish weight per 1 fish. As many as 5 fish were sampled from the tank.

3.2.3 Fish mortality data

Fish mortality is the rate or level of fish death in general. In this research, the fish mortality was relatively low. This was because African catfish was cannibalistic. From the data in Fig. 11, 77 fish died in total. The cause for cannibalism in African catfish was generally because of their varied sizes (Yang et al., 2015 in Putri et al. 2022).

From the research that Enggar Patrio et al. [14] conducted on water quality in tarpaulin tank for African catfish's growth, the data on water quality parameters showed that the lowest temperature was 26.6 °C and the highest was 28.8 °C.

In addition, the pH data showed that it ranged from 7.5 to 7.9. It means the tank was still feasible for cultivating the fish

The data collected for DO show that the dissolved oxygen level in the tank of 7.06 m2 wide, and 9.42 m2 diameter, and fish stocking density of ± 4000 fish ranged between 0.80 and 1.19 ppm. This means that it did not meet the good standard quality requirement for the fish growth and survival.




Fig. 7. DO data chart





Fig. 8. Combined water quality data chart

← Monitoring Pakan				
Tanggal Awal 7 Oktober 2023 28		Tanggal Akhir Desember 2023		
**	8			
9/12/2023 09:53	960 Gr	匬		
29/11/2023 08:20	870 Gr	创		
19/11/2023 08:19	739 Gr	创		
9/11/2023 08:19	348 Gr	创		
29/10/2023 08:18	261 Gr	创		
18/10/2023 08:18	174 Gr	创		
08:19 29/10/2023 08:18 18/10/2023	261 Gr	Ū		

Fig. 9. Data on amount of feed

← Monitoring Berat Ikan				
Tanggal Awal 7 Oktober 2023		Tanggal Akhir 28 Desember 2023		
—	ō			
29/11/2023 08:22	20 G	r 🔟		
19/11/2023 08:22	17 G	r 🔟		
9/11/2023 08:22	8 Gr	创		
29/10/2023 08:21	6 Gr	Ū		
18/10/2023 08:03	4 Gr	D		

Fig. 10. Fish weight data

← Monitoring Mortality				
Tanggal Awal 7 Oktober 202	23 28	Tanggal Akhir B Desember 2023		
	ø			
6/12/2023 17:10	3	创		
6/12/2023 08:10	3	创		
4/12/2023 08:10	3	创		
3/12/2023 17:10	5	创		
2/12/2023 08:10	2	创		
1/12/2023 17:09	25	创		
1/12/2023 08:09	5	创		
30/11/2023 08:08	7	创		
7/12/2023 08:45	2	创		

Despira et al.; Asian J. Fish. Aqu. Res., vol. 26, no. 3, pp. 1-11, 2024; Article no.AJFAR.113570

Fig. 11. Mortality ikan data




Chart 1. Results of tarpaulin tank temperature measurement




Chart 2. Results of tarpaulin tank pH measurement

Despira et al.; Asian J. Fish. Aqu. Res., vol. 26, no. 3, pp. 1-11, 2024; Article no.AJFAR.113570



Chart 3. Results of tarpaulin tank DO measurement

4. CONCLUSION

From the research that been carried out, it could be concluded that the use of Grow My Fish app was highly effective in helping cultivators process and collect the data they needed. Additionally, from the data they had inputted from the African catfish cultivation employing biofloc technology, it was found that the water quality parameters were still compliant with SNI 6484.3:2014. The pH ranged from 6.5 to 8. The lowest temperature was 25.2°C and the highest was 37°C, in which this occurred only once. Meanwhile, the DO value was >3 mg/l. Nevertheless, the fish death in the biofloc tank cultivation was still inevitable since the DO increased drastically. In addition, the high temperature led to the death of 25 fish. Thus, it could be interpreted that the DO value had met the SNI, despite its frequent drastic increase.

When the African catfish cultivation using biofloc technology was compared to that using tarpaulin tank, there were still some weaknesses. For the pH and temperature values, they both met the provisions of SNI. Hence, it can be said that they were still feasible for use to cultivate the fish (optimal for fish cultivation). Meanwhile, for their DO values, both could still be said less optimal. The DO value for tarpaulin tank ranged from 0.80 to 1.19 ppm. This means it was still <3 ppm and could possibly hinder the fish growth.

ACKNOWLEDGMENT

The researcher would like to thank Padjadjaran University and Seamolec for their guidance and assistance in the completion of this research in MBKM (Merdeka Belajar Kampus Merdeka) project. The gratitude would also be extended to the Partner PT. Pesona Khatulistiwa Nusantara for their facilitation to this MBKM (Merdeka Belajar Kampus Merdeka) project, as well as PT. Sebelas Cipta Mandiri, Mr. Ir. Siswoyo, Msi and Mr. Syahroni S.Pi.Gr who have developed the Grow My Fish app that has helped make this research easier to complete. Lastly, the writer would also offer her appreciation to her parents and families who have prayed for the successful implementation of this project.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Suprapto, Samsir SL. Bioflok-165 secrets of successful catfish cultivation technology, Depok (ID): AGRO 165; 2013.
- Mukherjee P, Sarka G, Saha A, Sanyal T. Extensive study and data collection on the pituitary gland: A promising prospect revealed by surveying the fish market during the monsoon season. International Journal of Experimental Research and Review. 2022;29:73-79. Available:https://doi.org/10.52756/ijerr.202 2.v29.008
- Mondal P, Adhikary P, Sadhu S, Choudhary D, Thakur D, Shadab M, Mukherjee D, Parvez S, Pradhan S, Kuntia M, Manna U, Das A. Assessment of the impact of the different point sources of pollutants on the river water quality and the evaluation of bioaccumulation of heavy metals into the fish ecosystem thereof. Int. J. Exp. Res. Rev. 2022;27:32-38.

Available:https://doi.org/10.52756/ijerr.202 2.v27.003

- Irmawanto R, Panggayudi DS, Fanani AK. Design and development of a photovoltaic power management system for fish ponds based on arduino and Internet of Things (IoT). Computer Electrical Power Telecommunications Control Electronics. 2021;4(2).
- Maghfiroh H, Hermanu C, Adriyanto F. Automatic feeder protype with IoT monitoring for biofloc fisheries. Journal of Electrical Engineering and Technology; 2019.
- Ashari IF, Untoro CU, Praseptiawan M, Afriansyah A, Nurazmi Eka. IoT-Based tilapia cultivation monitoring and control system with biofloc (Case study: Sadewa Mandiri Fish Cultivation Group, Pringsewu). Scientific Journal of Community Service. 2022;22(2):471-472. Availablehttps://doi.org//10.24036/sb.0268.
- 7. Oktavia RP, Febrianti DA, Septianingsih AD, Ramadhan MIR, Diana L. Application the biofloc method in catfish of cultivation in the greenhouse, pucang sewu village, gubeng district. Surabaya. Journal of Community Service. 2022;1(2).
- 8. Ombong F, Salindeho IRN. Application of Biofloc Technology (BFT) in tilapia fish (*Orechromis niloticus*) culture. Aquaculture. 2016;4(2):16-25.

- 9. Windriani U. Biofloc system catfish cultivation. Directorate of Production and Cultivation Business. 2017;1-38.
- Firyadha MNA, Waluyo W, Hidayati N. Design and build automatic catfish feeding biofloc. Journal of Telecommunication Networks. 2023;13(2).
- Faridah, Diana S, Yuniati. Catfish cultivation using the biofloc method in conventional catfish farmers. Journal of Community Service. 2019;1(2). Available:https://doi.org/10.31960/caradde. vli2.74
- 12. Safsafubun FR, Undap SL, Salindeho IRN, Pangemanan NPL, Watung JC, Pangkey H. Fluctuations in water quality parameters floc development in and tilapia (Oreochromis niloticus) cultivation with the biofloc svstem at BPBAT Talelu. Aquaculture e-Journal. 2023;11(2):213-226.
- Rafiq A. Design of an IoT-based Biofloc fish pond water temperature and floc volume monitoring tool to support food security 4.0. Syarif Hidayahtullah State Islamic University; 2022.
- Patriono E, Amalia R, Sitia M. Water quality of cultivation ponds and tarpaulins for the growth of sangkuriang catfish (*Clarias gariepinus*) in catfish cultivator groups in pali regency, South Sumatra. Biosanins Sriwijaya. 2022;2(3). DOI:10.24233/ sribios.2.3.2021.378

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/113570