

Spontaneous Recanalization of an Occluded Left Internal Mammary Graft after Documented Atresia

Sergey Yalonetsky ^{a*}, Yair Feld ^a, Amir Solomonica ^a, Anees Musalem ^a,
Gil Gross ^a and Arthur Kerner ^a

^a Interventional Cardiology Unit, Rambam Healthcare Campus, Technion Faculty of Medicine, Haifa, Israel.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85035>

Received 25 January 2022

Accepted 31 March 2022

Published 05 April 2022

Case Study

ABSTRACT

An internal mammary graft atresia after CABG surgery is a well-known complication which could be related to competitive flow in the grafted artery. Spontaneous recanalization of a previously occluded LIMA graft is an extremely rare phenomenon. We describe a patient in whom the LIMA was occluded 2.5 years after CABG surgery, but was found to be completely patent 10 years later, probably due to disease progression in the native grafted vessel.

Keywords: Internal mammary graft; coronary bypass graft occlusion; coronary bypass graft recanalization.

1. INTRODUCTION

Internal mammary arteries (IMA) are the grafts of choice for coronary artery bypass (CABG) because they are generally free of atherosclerosis and they have high patency rates compared with venous grafts [1]. While IMA atresia is a well-known complication post CABG surgery [2,3], spontaneous patency restoration of an occluded IMA graft is an extremely rare

phenomenon. We report a patient in whom the left internal mammary graft (LIMA) was occluded 2.5 years after CABG surgery, but was found to be completely patent 10 years later.

1.1 Learning Objective

Spontaneous recanalization of an atretic mammary graft may be observed as long as 10 years after its documented occlusion. While graft

*Corresponding author: Email: s_yalonetsky@rmc.gov.il;

degeneration is usually related to competitive flow, its spontaneous return may be explained by disease progression in the grafted vessel.

2. CASE REPORT

A 70-year-old woman with a known atherosclerotic cardiovascular disease, hypertension and diabetes melitus was admitted for elective cardiac catheterization as a part of investigation for suspected diastolic heart failure.

Ischemic heart disease was diagnosed 12 years earlier when she was first admitted for unstable angina pectoris. Coronary at angiography (CA) at that time revealed 70% narrowing of the ostial left main coronary artery (LMCA) and a 50-75% narrowing in the mid left anterior descending coronary artery (LAD) segment. In the light of these findings the patient underwent CABG surgery which included a LIMA graft to the LAD and a free right internal mammary artery (RIMA) graft to the obtuse marginal artery (OM). Sixteen months following the surgery the patient started to experience recurrent angina and therefore underwent another cardiac catheterization. It revealed 75-90% narrowing of the distial LMCA, a patent LIMA graft to LAD and an occluded RIMA graft to OM. Percutaneous coronary intervention (PCI) to the LMCA with a bare metal stent (BMS) (Driver 4.0/9) was completed

successfully aiming to augment the blood flow into the LCX artery.

Fourteen months later the patient underwent repeat coronary angiography for exertional dyspnea which was considered to be anginal equivalent. Total LIMA graft obliteration ("string phenomenon") was demonstrated (Fig. 1), and no additional flow limiting lesions were detected.

Repeat coronary angiography in 2014 revealed similar findings.

Current elective admission in April 2021 for diagnostic cardiac catheterization was related to ongoing effort dyspnea and left ventricular diastolic dysfunction revealed by echocardiography. Her ongoing medication included antiplatelets (clopidogrel), statins and beta blocker, angiotensin receptor antagonist and insulin. The hemodynamic measurements demonstrated moderate pulmonary hypertension (mean arterial pressure of 38 mm Hg) with normal pulmonary vascular resistance and elevated left ventricular end diastolic pressure, consistent with diastolic heart failure, probably attributed to age related myocardial compliance reduction as well as chronic hypertension and ischemic heart disease. Coronary angiography demonstrated a 75-90% mid LAD narrowing while the LIMA graft which was previously shown to be occluded was now patent (Fig. 2).




Fig. 1. Coronary angiography from 2011 showing atretic LIMA graft ("string phenomenon")




Fig. 2. Coronary angiography from 2021 showing patent LIMA

3. DISCUSSION

We describe a case of spontaneous recanalization of a LIMA graft 10 years after it was shown to be occluded. Postoperative degeneration of a LIMA graft after CABG surgery is a well-known complication which could be related to competitive flow in the grafted artery [2,3] as well as to an on-going atherosclerotic disease in the native vessel, which has extended to the anastomotic area, or other mechanisms [4]. However only a few reports describe the recanalization of a previously occluded LIMA graft [5-8].

In the presented case the LMCA stenting was performed to improve LCX blood flow since the free RIMA graft to OM had been shown to be occluded; this probably also enhanced the blood flow into the LAD causing competitive haemodynamics with subsequent LIMA degeneration. The recanalization of the LIMA graft which was later observed could probably be explained by the progressive mid LAD narrowing over the time.

Feld et al described an early postoperative LIMA to LAD occlusion (3 months after surgery) related to progressive narrowing of the LAD segment located distal to the anastomosis which was successfully stented. Repeat angiography

performed 4 months later revealed patent a LIMA graft along with proximal LAD narrowing progression and widely patent distal angioplasty site [5].

In the report of Nawaz et al the LIMA graft was shown to be occluded 2 years after CABG surgery, but was found completely patent 5 years later, as a result of worsening native LAD disease [6].

A similar case had been described by Khalid et al showing LIMA graft "return" 9 years after its documented occlusion [7].

Meredith et al described a patient who underwent bypass surgery with a LIMA to LAD graft for an anomalous LMCA originating from the right coronary cusp with retro-pulmonary interarterial course. The LIMA graft was found to be atretic 3 months after surgery. Repeat coronary imaging 7 years later showed recanalization of the previously atretic LIMA probably explained by atherosclerotic plaque progression in the distal LMCA [8].

Similarly to the previously published reports, in the case presented here, LIMA recanalization was related to the progression of native LAD disease; however, the ten-years' time interval between the documented occlusion and

recanalization is longer compared to the previously published cases.

4. CONCLUSION

The case presented here demonstrates that LIMA graft atresia caused by competitive flow, can be reversible, while the restoration of its patency is probably related to disease progression in the native grafted vessel.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

DECLARATION

The authors declare that:

- 1) The paper is not under consideration elsewhere;
- 2) None of the paper's contents have been previously published;
- 3) All authors have read and approved the manuscript;
- 4) The authors do not have any relationship with industry or other conflict of interest.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Windecker S, Kolh P, Alfonso F, Collet JP, Cremer J, Falk V, A. 2014 ESC/EACTS Guidelines on myocardial revascularization: The Task Force on

- Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS) Developed with the special contribution of the European Association of Percutaneous Cardiovascular Interventions (EAPCI). *Eur Heart J*. 2014 Oct 1;35(37):2541-619.
2. Villareal RP, Mathur VS. The string phenomenon: an important cause of internal mammary artery graft failure. *Tex Heart Inst J*. 2000;27(4):346-9.
3. Hashimoto H, Isshiki T, Ikari Y, Hara K, Saeki F, Tamura T et al. Effects of competitive blood flow on arterial graft patency and diameter. Medium-term postoperative follow-up. *J Thorac Cardiovasc Surg*. 1996 Feb;111(2):399-407.
4. Berger A, MacCarthy PA, Vanermen H, De Bruyne B. Occlusion of internal mammary grafts: A review of the potential causative factors. *Acta Chir Belg*. 2004 Nov-Dec;104(6):630-4.
5. Feld H, Navarro V, Kleeman H, Shani J. Early postoperative occlusion of a left internal mammary artery bypass graft with subsequent restoration of patency. *Catheterization and Cardiovascular Diagnosis* 1992 Dec;27(4): 280–3.
6. Nawaz R, Brener SJ. Flow-dependent changes resulting in restoration of patency of the left internal mammary artery graft after documented atresia. *The Journal of Invasive Cardiology*. 2011 May;23(5):207–10.
7. Khalid N, Shlofmitz E, Chen Y, Chen Y, Case B, Dheendsa A et al. Return of the Left Internal Mammary Artery. *Cardiovasc Revasc Med*. 2021;23:119-20.
8. Meredith T, Jansz P, Sesel K, Kathir K. Recanalization of an atretic left internal mammary graft after bypass surgery for an anomalous left main coronary artery origin: a case report. *Eur Heart J Case Rep*. 2018;3(1):yty152.

© 2022 Yalonetsky et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85035>