



## Antioxidant Potential and Anti-Sickling Activity of Different Organs of *Curcuma longa*: Correlation of the Antioxidant Capacity on Anti-Sickling Activity

Clément M. Mbadiko<sup>1\*</sup>, Koto-te-Nyiwa Ngbolua<sup>1</sup>, Pius T. Mpiana<sup>2</sup>,  
Nadège K. Ngombe<sup>3</sup>, Paulin M. Kapepula<sup>3</sup>, Landry L. Kemfine<sup>1,4</sup>,  
Gédéon N. Bongo<sup>1</sup> and Théophile F. Mbemba<sup>1</sup>

<sup>1</sup>Department of Biology, Faculty of Sciences, University of Kinshasa, P.O.Box 190, Kinshasa XI, Democratic Republic of the Congo.

<sup>2</sup>Department of Chemistry, Faculty of Sciences, University of Kinshasa, P.O.Box 190, Kinshasa XI, Democratic Republic of the Congo.

<sup>3</sup>Centre d'Etudes des Substances Naturelles d'Origine Végétale (CESNOV), Faculty of Pharmaceutical Sciences, University of Kinshasa, Kinshasa XI, Democratic Republic of Congo.

<sup>4</sup>Chemical, Biological, Radiological and Nuclear Centre of Excellence, Ministry of Scientific Research, Kinshasa, Democratic Republic of the Congo.

### Authors' contributions

This work was carried out in collaboration among all authors. Authors PTM and TFM designed the study. Author PMK performed the statistical analysis. Author NKN wrote the protocol. Authors KNN and CMM wrote the first draft of the manuscript. Authors CMM and GNB managed the analyses of the study. Authors LLK and GNB managed the literature searches. All authors read and approved the final manuscript.

### Article Information

#### Editor(s):

- (1) Dr. MVNL Chaitanya, Associate Professor, Department of Pharmacognosy and Phytopharmacy, Guru Nanak School of Pharmacy, Hyderabad, India.
- (2) Dr. Prasong Srihanam, Professor, Department of Chemistry, Faculty of Science, Mahasarakham University, Thailand.
- (3) Dr. Dildar Ahmed, Professor, Department of Chemistry, Forman Christian College University Lahore, Pakistan.
- (4) Dr. Stefano Manfredini, Professor of Pharmaceutical Chemistry and Toxicology, Department of Life Sciences and Biotechnology, University of Ferrara, Italy.

#### Reviewers:

- (1) Muhammad Shahzad Aslam, Xiamen University, China.
- (2) Tchoffo Herve, University of Dschang, Cameroon.
- (3) Syed Umer Jan, University of Balochistan, Pakistan.
- (4) Timur Chis, Ovidius University, Romania.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46718>

Original Research Article

Received 27 January 2019  
Accepted 15 April 2019  
Published 07 May 2019

### ABSTRACT

**Aims:** The aim of this research was to evaluate comparatively the anti-sickling activity of aqueous and methanol extracts of different organs of *C. longa*, to determine the polyphenolic compounds and to evaluate the antioxidant potential of different parts of *C. longa*. Also, to establish a possible

correlation between the antioxidant potential of the extracts on the anti-sickling activity.

**Study Design:** The study used an experimental design. The antioxidant and anti-sickling activities of the aqueous and methanol extracts of different organs of *C. longa* were assessed along with their correlation.

**Place and Duration of Study:** The study was carried out at the Laboratory of Natural Products, Faculty of Science and at Centre for the Study of Natural Products from Plant Origin (CESNOV), Faculty of Pharmaceutical Sciences, both at University of Kinshasa in 2018.

**Methodology:** *C. longa* was planted in an experimental monoculture setting from where were collected different organs of this plant. After processing these parts, they were macerated in methanol and water respectively. After filtration of the macerates, the filtrates were dried in the oven to obtain aqueous and methanol extracts which were used for biological analyses and the determination of polyphenolic compounds. This determination was performed using a spectrophotometer while the biological activities was performed using the ABTS, DPPH and Emmel tests respectively. Data were analyzed using GrapPad Prism 6.0 software for statistical analyzes and the determination of IC<sub>50</sub>.

**Results:** The findings showed that the polyphenolic compounds are not distributed in the same way in different organs of *C. longa*. The extracts of different parts of *C. longa* do not all have a great antioxidant potential. However, the rhizome and root extracts showed a better anti-scavenging activity using ABTS method compared to other organs, while rhizome and leaf extracts showed better anti-scavenging activity using the DPPH test. The analyses carried out revealed: a positive correlation between (i) the total polyphenol contents and the anti-sickling activity of the studied extracts ( $r = 0.08328$ ) and (ii) between the antioxidant activity using ABTS and DPPH tests along with the anti-sickling activity of different extracts ( $r = 0.1221, 0.4900$  and  $0.3114$  respectively).

**Conclusion:** The tested extracts displayed good antioxidant and anti-sickling activities. Thus, different organs of *C. longa* are a source of natural antioxidants, which can be used to manage sickle cell disease. The effect of the antioxidant potential of the extracts tested on their anti-sickling activity was demonstrated. Nevertheless, in-depth studies are required in order to confirm this correlation.

**Keywords:** *Curcuma longa*; antioxidant activity; anti-sickling activity; polyphenol content; correlation.

## 1. INTRODUCTION

Sickle cell disease is a genetic disease with autosomal recessive inheritance linked to an abnormality of hemoglobin precisely hemoglobin S. At the molecular level, it is a hemoglobinopathy resulting from a point mutation at the sixth codon of the  $\beta$ -globin gene carried by chromosome 11. In fact, it is a substitution of Adenine by Thymine at the 6<sup>th</sup> codon in the  $\beta$ -globin gene. At the protein level, this mutation leads to the replacement of glutamic acid (polar amino acid) at the 6<sup>th</sup> position by a valine (apolar amino acid) ( $\beta 6$  Glu  $\rightarrow$  Val), thus the synthesis of an abnormal hemoglobin, hemoglobin S (Hb S), in English "sickle" [1-4]. Sickle cell disease is responsible for the loss of life worldwide, and the most vulnerable layer is children under 5 years old [5]. From its regions of origin (Central and West Africa), it has moved and continues to move with migratory flows (Mediterranean region, Asia, USA, the Caribbean). However, the African populations are particularly affected. In some

West African countries, 30-40% of the population is carrier this gene [3]. Several treatment modalities have been considered namely the bone marrow transplantation, gene therapy, blood transfusion and the hydroxyl-urea intake in order to relieve patients. The truth is that these treatments are not only ineffective but very costly for poor people in Africa, but can also be the cause of several adverse effects (toxicity, blood incompatibility, graft versus host reaction, HIV/AIDS contamination) [5-7]. Due to all the known side effects, medicinal plants are used in Africa to relieve several ailments. In the Democratic Republic of the Congo (DRC), the phytotherapy is currently used as an alternative, which can provide a relief to sickle cell patients. The review of the literature indicated that the screening of several plants have been revealed a potential anti-sickling. Among these plants were also mentioned *Curcuma longa* [2,5-9,10-18].

Belonging to Zingiberaceae family, *C. longa* L. is a perennial herbaceous plant measuring up to 1m high. It is mostly cultivated in the tropics but a

large part of the production is performed in India and Southeast Asia [9,19]. Several studies reported that the intense cultivation of *C. longa* is due to the use of its rhizomes. For centuries, *C. longa* rhizomes were efficiently used in various sectors including the food, cosmetology, agri-food, pharmaceutical, textile and traditional medicine industries [9,19-47]. Furthermore, its leaves are used for the extraction of essential oils and the seasoning of dishes or as a colouring agent in the industry and in the traditional kitchens respectively. In traditional medicine, these leaves are used as tonic, stimulant and blood purifier in the valley districts of Manipur, India [32,37,43-45]. Mbadiko et al. [9] and other researchers [29,35,40,43,48-51], reported that extracts of rhizomes of *C. longa* have broad pharmacological potential such as anti-cancer, anti-inflammatory, healing, cholesterol-lowering, hypoglycemic, anti-Alzheimer's, antiplasmodial, anti-inflammatory, antioxidant, antibacterial, antifungal, anti-venom, antipyretic, analgesic, inhibits the action of the integrase of HIV-1 replication, protects against diabetic retinopathy and numerous other pathologies. However, other studies have shown that the pharmacological actions reported on rhizome extracts of *C. longa* are linked to the presence of polyphenolic compounds, particularly curcuminoids notably curcumin, but also to its essential oils [8-9,19,21,32-33,52-66]. Ajay et al. [33] and Camatari et al. [26], showed that *C. longa* flowers possess the antioxidant potential. In addition, Lassa [67] reported that *C. longa* rhizomes is used for the treatment of SS anemia in traditional medicine in DRC. This use was scientifically justified by Mbadiko et al. [9] whom showed the anti-sickling potential of total methanol extracts from different organs (rhizomes, roots, leaves, sepals and petals) of *C. longa*. Henceforth, the interest to carry out a comparative study on the anti-sickling activity of methanol and aqueous extracts of *C. longa* different organs.

By the way, several reports have showed the involvement of oxidative stress in sickle cell anemia. Hierso [1] reported that sickled red blood cells produce more free radicals than normal red blood cells, this leads to the increase of the production of oxidative biomarkers, which would actively participate in the physiopathology mechanisms of the disease. Then, it plays a role in the hemolysis and the onset of vaso-occlusive crises as well the attacks of various organs. Due to oxidative damage associated with sickle cell disease, the development of therapeutic

antioxidant strategies is proven essential and relevant. Henceforth, the aims of this study were to (i) evaluate the antioxidant potential of different parts of *C. longa* and (ii) look for a possible correlation between the content of polyphenolic compounds, the antioxidant potential and the anti-sickling activity of the extracts. Studies raising from this perspective have not yet been reported in the literature.

## 2. MATERIALS AND METHODS

### 2.1 Biological Material

Different organs of *C. longa* (rhizome, root, leaf, petal and sepal) were used as biological material. The plant was first cultivated as a monoculture in an experimental setting and different parts were harvested. These organs brought to the laboratory where they were washed, dried in the oven (Melag Nur Für Wechselstrom brand) at 30°C and crushed using a Moulinex electric mill. Different powders obtained were used for the preparation of different extracts that were used for the phytochemical screening and the evaluation of biological activities.

This study was carried out at the laboratory of Natural Products, Department of Chemistry and at the laboratory of Analyses and Research on Food and Nutrition (LARAN), both at Faculty of Science and at Centre for the Study of Natural Products from Plant Origin (CESNOV), Faculty of Pharmaceutical Sciences, both at University of Kinshasa in 2018.

### 2.2 Determination of Total Phenolic Content

The dosage of the polyphenolic compounds was carried out according to the method described by Kapepula et al. and Adepado et al. [68-69]. In fact, 10 mg/mL of each extract were diluted in methanol 80% in order to obtain a solution of 1 mg/mL for each extract. Then, a mixture consisted of 0.5 mL of the extract; 5.0 mL distilled water and 0.5 mL of Folin-Ciocalteu reagent was prepared for each extract. After three minutes, 1.0 mL of a saturated solution of Na<sub>2</sub>CO<sub>3</sub> 20% was added. The mixtures prepared were stirred and incubated at laboratory temperature under shade for one hour. After the incubation, different absorbances were read using a spectrophotometer at 725 nm. Each analysis was performed in triplicate. The quantity of total polyphenols was expressed in mg of

gallic acid equivalents (GAE)/g using the following equation from the calibration line:

$$y = 1.7097 \ln(x) + 5.2062 \quad (1)$$

and  $R^2=0,965$ , where  $x$  is the absorbance and  $y$  is the equivalent of gallic acid (mg/g).

### 2.3 Dosage of Total Flavonoids

The estimation of the total flavonoid content of our extracts was performed using a spectrophotometer. Aluminium trichloride forms a yellow complex with flavonoids that absorbs at 415 nm [69]. First, a mixed reaction containing 1mL of the methanol solution (80%) of each extract (concentration 10mg/mL) and 1ml of  $AlCl_3$  (2%) (dissolved in methanol) was prepared. The mixtures were then stirred and incubated for one hour at the laboratory temperature under shade and the absorbances were read at 415 nm. For the preparation of the control, the process was the same as described above except that instead of using the extract, 1 mL of methanol was added. Each analysis was performed in triplicate. The flavonoid content of the extracts was expressed in mg quercetin equivalent (QE)/g using the following equation from the calibration line:

$$y = 0.5001 \ln(x) + 3.442 \quad (2)$$

$R^2 = 0,944$ , where  $x$  is the absorbance and  $y$  is the equivalent of quercetin (mg/g).

### 2.4 Determination of Total Anthocyanins

The determination of the anthocyanin content of extracts was carried out following the protocol as per Adedapo et al. [69]. We took 0.5 mL of the 1 mg/mL solution of extract prepared with methanol (80%), which was mixed with 3 mL of vanillin methanol solution (4%) and 1.5 mL hydrochloric acid. The mixture was incubated for one hour, then the absorbance was measured at 500 nm. The same procedure was repeated for the control (D-catechin) in the concentration range (from 4 mg/mL to 0.125 mg/mL) and allowed the calibration line to be established. The anthocyanin content of the extracts was expressed in mg equivalent of D-catechin per g of dry vegetable matter (EC/g) using the following equation from the calibration line:

$$y = 0.0728 x + 0.0171 \quad (3)$$

and  $R^2 = 0.994$ , where  $x$  is the absorbance and  $y$  is the equivalent of catechin (mg/g).

## 2.5 Assessment of the Anti-Sickling Activity

### 2.5.1 Blood samples

The blood samples used in our experiments were collected at the Centre de Médecine Mixte et d'anémies SS (CMMSS) from the sickle cell patients. Once in the laboratory, they were kept in the refrigerator and the Emmel test was performed 24 hours later to ensure that these samples were actually taken from sickle cell patients.

### 2.5.2 Preparation of dry methanol extracts of various organs of *C. longa*

Using a precision scale (Kern 440-35N), it was weighed respectively 25 g of roots and rhizomes, 20 g of leaves, 3.69 g of sepals and 5 g of petals which were macerated in 100 mL of methanol during 24 hours. The macerates were then filtered using a Whatman paper and dried in the oven at 37°C (Melag Nurfur Wechselstrom) for five days.

### 2.5.3 Preparation of dry aqueous extracts of various organs of *C. longa*

With a precision scale (Kern 440-35N), it was weighed respectively 20 g of rhizomes and roots, 15 g of leaves and 10 g of sepals and petals of *C. longa* powders that were diluted in 150 mL of distilled water. After 24 hours of maceration, the macerates were filtered using a Whatman no1 filter paper, poured into petri dishes and dried in an oven at 40°C for a week. The dry extracts obtained were used for the Emmel's test.

### 2.5.4 Emmel's test

The principle of this test is the following: Hemoglobin S is less soluble when the oxygen content decreases (hypoxia,  $pO_2 < 45 \text{ mmHg}$ ). In sickle cell disease, the sickling of erythrocytes is observed due to the intracellular crystallization of hemoglobin S in the form of tactoids. Therefore, unlike the normal red blood cell, which has a biconcave or circular shape, the red blood cells of sickle cell patients have a sickle or banana shape under hypoxic conditions. The effect of crude plant extracts, which prevents cell sickling under hypoxic conditions (created by the addition of sodium metabisulfite 2%, m/v), is used as an indication of the pharmacological activity of these plants.

The anti-sickling activity of the aqueous and methanol extracts of different parts of *C. longa* was evaluated using the Emmel's test, as described previously [8-18,70]. Different concentrations of extracts were prepared notably: 1 mg/mL, 500 µg/mL, 250 µg/mL, 125 µg/mL, 62.5 µg/mL, 31.25 µg/mL and 15.62 µg/mL, using saline solution (NaCl 0.9%, m/v) as a solvent. From this solution, we were able to make a microscopic preparation whereby a drop of SS blood was placed on the slide to which was a drop of Na<sub>2</sub>S<sub>2</sub>O<sub>5</sub> 2% in order to create a hypoxic condition. Then, a drop of the extract was added to that preparation, which was mixed thoroughly and carefully covered to avoid the formation of air bubbles. Another microscopic preparation not containing the extracts was made under the same conditions and served as a negative control. The edges of the slide was supercooled with paraffin to reinforce the hypoxic solution. The results was read after 24 hours using a microscope (Bresser) at 40 of magnitude. All the analyses were carried out in triplicate.

## 2.6 Antioxidant Activity

The evaluation of the antioxidant activity was performed using the ABTS and DPPH tests as described by Kapepula et al. [68].

### 2.6.1 Preparation of samples

In fact, 10 mg of dry extract from each sample was dissolved in 1mL of methanol for polar extracts and dichloromethane-methanol mixture (1:1) for apolar extracts (solution A: 10 mg/mL) respectively. We then diluted to the following concentration: 8mg/mL, 6 mg/mL, 4mg/mL and 2 mg/mL.

### 2.6.2 Preparation of ABTS radical

In 500 µL of distilled water was dissolved a quantity of the ABTS reagent corresponding to 20 millimoles: solution A. In another 500µL of distilled water was dissolved a quantity of potassium persulphate (K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>) corresponding to 10 millimoles: solution B. Both solutions were mixed at the same volume and the mixture was kept under shade between 12 and 16 hours. This mixture constituted the ABTS•<sup>+</sup> radical stock solution. Thereafter, the stock solution of the radical was diluted several times with methanol to obtain an analytical solution having an absorbance ranging from 0.800 to 1.000.

### 2.6.3 ABTS test

When reacted with potassium persulfate (K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>), ABTS (2,2'-azino-bis-3ethylBenz-

Thiazolin-6-Sulfonic acid) forms the cationic radical ABTS<sup>+</sup> from blue to green. The addition of the antioxidant reduces this radical and causes the mixture to discolour. The radical discoloration measured by spectrophotometer at 734 nm is proportional to the antioxidant concentration [68]. For each sample and concentration (8 mg/mL, 6 mg/mL, 4 mg/mL and 2 mg/mL), 20 µL of the solution was collected and 1980 µL of the ABTS<sup>+</sup> radical analysis solution was added later. The reaction mixtures were incubated for 30 minutes and the solutions were successively read with a spectrophotometer at 734 nm. The same procedure was repeated for the preparation of the control consisting of methanol (20 µL) and ABTS (1980 µL). The analysis was performed in triplicate.

The percentage of inhibition of the ABTS<sup>+</sup> radical by the sample is determined using the following formula:

$$\%inhibition = [1 - \left(\frac{A_x}{A_c}\right)] \times 100 \quad (4)$$

Where

A<sub>x</sub>: the absorbance of the ABTS<sup>+</sup> radical in the presence of the extract

A<sub>c</sub>: the absorbance of the ABTS<sup>+</sup> radical (control solution)

### 2.6.4 Preparation of the DPPH radical

A quantity of 3.2 mg of DPPH reagent was dissolved in 100 mL of methanol 80% and this solution was kept under shade for at least one hour. The absorbance of this solution was adjusted to 0.7 ± 0.05 using methanol 80%.

### 2.6.5 DPPH test

The DPPH test is based on the degradation of the DPPH- (2,2 DiPhenyl-1- PicrylHydrazyl) radical. The DPPH- radical is a violet coloured radical, the addition of antioxidant reduces this radical and causes the discoloration of the mixture; this discoloration of the radical measured by spectrophotometer at 517 nm is proportional to the concentration of antioxidants added [68]. For each sample and dilution level, 20 µL of the solution was taken and 1980µL of the DPPH- radical analysis solution was added. The reaction mixtures were incubated under shade for 30 minutes and the solutions were read successively with a spectrophotometer at 517 nm. We proceeded in the same way for the

preparation of the control, except that instead of the extract, 20  $\mu$ L of methanol was added. As with the ABTS test, the analysis was performed in triplicate.

The percentage of inhibition of the ABTS+ radical by the sample is determined using the following formula:

$$\%inhibition = [1 - \left(\frac{A_x}{A_c}\right)] \times 100 \quad (5)$$

Where

A<sub>x</sub> : the absorbance of the DPPH- radical in the presence of the extract

A<sub>c</sub> : the absorbance of the DPPH- radical (control solution)

## 2.7 Data Analysis

The GraphPad Prism 6.0 software was used for data analysis for statistical analyzes and the determination of IC<sub>50</sub>. For data presented as mean  $\pm$  standard deviation, we performed the ANOVA one Way test for comparison of sample averages, followed by Tukey's pairwise multiple comparison test. We considered the threshold of significance at  $\alpha = 0.05$ . In addition, Pearson's test was used in order to assess the correlation between some variables.

## 3. RESULTS AND DISCUSSION

### 3.1 Total Phenolic Content

The content of polyphenolic phytoconstituents of different extracts of *C. longa* is presented in Table 1.

The ANOVA showed that extracts from different organs of *C. longa* showed very highly significant differences ( $p$  value <00001) in total polyphenol content. However, the Tukey pair multiple comparison test (HSD = 0.4938) does not reveal a significant difference between roots and leaves

and between sepals and petals. This indicates that the total polyphenol content is low in the sepals and petals. It has the following descending order: Rhizomes > roots, leaves > petals and sepals. This shows that rhizomes are an important source of total polyphenol in *C. longa*. Maizura et al. [23] reported that *C. longa* rhizomes are rich in polyphenolic compounds. Although the rhizome samples used in their study had a high content of total polyphenols (67.9  $\pm$  1.0 mg GAE/100g extracts) compared to those used in this study. This may be explained by the fact that these two samples were not collected in the same geographical environment. Gayatri et al. [71], reported that the rhizomes of *C. longa* have a high content of total polyphenols compared to other species of *Curcuma* genus.


However, the floral extracts showed low levels of total polyphenols Camatari et al. [26], Ajay et al. [33] and Yan et al. [35] revealed the presence of large amounts of total polyphenols in floral extracts (2013.09  $\pm$  5.13 mg GAE/100 g, 210.45  $\pm$  1.32 mg GAE/100 g and 119 mg  $\pm$  2 GAE/g respectively). Chang et al. [72] also reported that *C. longa* flowers were rich in polyphenolic compounds (including curcuminoids). In fact, the content of secondary metabolites precisely polyphenolic compounds for the current study may vary with respect to the geographical origin, the period of collection, the physicochemical and biochemical characteristics of the soil, to the method and solvents used for the extraction, to the storage and post-harvest treatment conditions, the genotype of the plant and the part of the plant used [21-22,50,73-76].

In this study, the post-harvest treatment of floral pieces may have influenced polyphenolic content. Indeed, while drying floral pieces under shade at the laboratory temperature; these organs tend to soften and lose color; which implies a deterioration of pigmentary principles including polyphenolic compounds. Yan et al. [35] and Chang et al. [72] have also reported the

**Table 1. Content of polyphenolic phytoconstituents in different organs of *C. longa***

Samples	Tota polyphenol content (mg/EAG) (n=3)	Flavonoid content (mg/EQ) (n=3)	Anthocyanin content (mg/EC) (n=3)
Rhizomes	12.82 $\pm$ 0.63 <sup>a</sup>	0.020 $\pm$ 0.001 <sup>a</sup>	0.29 $\pm$ 0.05 <sup>a</sup>
Roots	5.91 $\pm$ 0.87 <sup>b</sup>	0.010 $\pm$ 0.001 <sup>b</sup>	0.17 $\pm$ 0.04 <sup>a,b</sup>
Leaves	4.31 $\pm$ 0.68 <sup>c</sup>	0.020 $\pm$ 0.002 <sup>c</sup>	0.30 $\pm$ 0.19 <sup>a</sup>
Petals	1.53 $\pm$ 0.28 <sup>d</sup>	0.007 $\pm$ 0.000 <sup>d</sup>	0.05 $\pm$ 0.01 <sup>b,c</sup>
Sepals	0.59 $\pm$ 0.30 <sup>d</sup>	0.006 $\pm$ 0.001 <sup>d</sup>	0.01 $\pm$ 0.00 <sup>b,c</sup>

Mean values followed by the same letters in each column do not differ significantly at  $p \leq 0.05$  by the Tukey test, n= number of repetition of the experiment



**Fig. 1. Anti-sickling activity of methanol and aqueous extracts of different organs of *Curcuma longa* L**

**Table 2. Evaluation of the anti-scavenging activity of methanol extracts from different organs of *C. longa* expressed in IC<sub>50</sub> by the ABTS and DPPH techniques**

Samples	ABTS (µg/mL) (n=3)	DPPH (µg/mL) (n=3)
Roots	4.66 ± 0.34 <sup>a</sup>	-
Rhizomes	4.43 ± 0.18 <sup>a</sup>	15.63 ± 7.81 <sup>a</sup>
Sepals	46.24 ± 2.51 <sup>b</sup>	1129.8 ± 559.64 <sup>b</sup>
Petals	24.27 ± 1.43 <sup>c</sup>	553.35 ± 250.15 <sup>c</sup>
Leaves	14.93 ± 4.30 <sup>d</sup>	54.83 ± 19.35 <sup>d</sup>
Gallic acid (Control)	0.71 ± 0.08	1.07 ± 0.10

Mean values followed by the same letters in each column do not differ significantly at  $p \leq 0.05$  by the Tukey test, n= number of repetition of the experiment

**Table 3. Comparison of total polyphenol content and antioxidant activity of different organs of *C. longa***

Plant extracts	Total polyphenol content (mg/EAG)(n=3)	Antioxidant activity	
		ABTS (µg/mL)	DPPH (µg/mL)
Rhizomes	12.82 ± 0.63	4.43 ± 0.18	15.63 ± 7.81
Roots	5.91 ± 0.87	4.66 ± 0.34	-
Leaves	4.31 ± 0.68	14.93 ± 4.30	54.83 ± 19.35
Petals	1.53 ± 0.28	24.27 ± 1.43	553.35 ± 250.15
Sepals	0.59 ± 0.30	46.24 ± 2.51	1129.80 ± 559.64

degradation of plant polyphenolic compounds under cellular conditions i.e. fresh samples. Moreover, Yan et al. [35], reported that dried and

powdered *C. longa* leaves have a high total polyphenol content as fresh samples. The author justified this observation by the fact that the

degradation of phytoconstituents in cellular conditions probably due to the activation of hydrolases and/or cellular oxidases. Secondary metabolites are synthesized mainly in response to environmental constraints (biotic or abiotic stresses) and intervene in plant-environment interactions. Some secondary metabolites are considered as messenger molecules and are involved in the initiation of certain biochemical and/or physiological processes of the plant. These processes explain the presence or absence of phytoconstituents during a period of the plant cycle or following the environment conditions [77]. In addition, the concentration of certain polyphenolic compounds, especially curcuminoids in the rhizomes of *C. longa*, becomes important five months after planting and declines between the 9<sup>th</sup> and 10<sup>th</sup> month [21]. This author also stated that the primary rhizomes are rich in phenolic compounds as well as secondary rhizomes [21].

However, we observed low levels of flavonoids and total anthocyanins in different extracts studied. The ANOVA also showed a very highly significant difference in the distribution of flavonoids in different parts of *C. longa* (p value <0, 0001). The Pairwise multiple comparison using the Tukey test (HDS = 0.0005357) shows that there is no significant difference in flavonoid levels in petals and sepals. The order of flavonoid concentration in the different parts of *C. longa* is as follows: Leaves> rhizomes> roots> sepals and petals. The flavonoid content of these extracts is expressed in mg equivalent of quercetin per g of dry extracts. Meanwhile, the ANOVA revealed a highly significant difference in anthocyanin levels of different extracts (p value = 0.0073). Although the Tukey multiple comparison test (HSD = 0.07237) revealed a significant difference only between rhizomes and sepals, sepals and leaves as well as petals and leaves. This shows that the petals and sepals are poor in anthocyanins. In addition, it was observed that the polyphenolic phytoconstituents studied are not distributed in the same way in different organs of *C. longa*. It should be noted that studies on the determination of flavonoid and anthocyanin content of different organs of *C. longa* are still poor.

### 3.2 Anti-Sickling Activity

The anti-sickling activity of aqueous and methanol extracts of different organs of *C. longa* are evident from the findings of this study (Fig. 1). The anthocyanin-rich leaf extracts showed a

great anti-sickling activity (15.62 µg/mL). According to Ngbolua et al. [7], Mpiana et al. [10-18] and Tula [70], the anti-sickling activity of plants is generally related to the presence of anthocyanins and organic acids. In fact, anthocyanins interact with the hemoglobin S and compete with the polymerization reaction by inhibiting the erythrocyte sickling process [2,70]. This could justify the fact that leaf extracts showed at the same time high anthocyanin content, and a strong anti-sickling activity. However, the aqueous extracts of rhizomes, leaves and sepals showed a high anti-sickling activity compared to methanol extracts. Mbadiko et al. [9] also reported the anti-sickling activity of methanol extracts of different organs of *C. longa*.

Fig. 1 displays the normalization of sickle cells treated with methanol leaf extracts of 31.25 µg/mL (B); rhizomes of 62.50 µg/mL (C), petals of 62.50 µg/mL (D), roots of 62.50 µg/mL (E), sepals of 62.50 µg/mL (F), and on the other hand, sickle cell treated with aqueous rhizome extracts of 31.25 µg/mL (G), roots of 62.50 µg/mL (H), petals of 62.50 µg/mL (I), sepals of 31.25 µg / mL (J) and leaves 15.62 µg/mL (K) compared to control, blood SS erythrocytes not treated (A).

The ANOVA showed a very highly significant difference for the anti-sickling activity of methanol extracts of different organs of *C. longa* (p value <0.0001). The Tukey multiple comparison test did not show a significant difference between the activities of rhizome and root, rhizome and sepal as well as rhizome and petal. This reflects the following order in relation to the anti-sickling of extracts from different organs: Leaves > Rhizome, root, petal and sepal. Furthermore, the ANOVA also showed a very highly significant difference for the anti-sickling activity of aqueous extracts of different organs of *C. longa* (p value <0.0001). However, the Tukey Multiple Comparison Test did not reveal a difference between the rhizome and sepal, and then between root and petal. The anti-sickling activity of aqueous extracts follows the following order: Leaves> rhizome, sepal >Petal root.

### 3.3 Antioxidant Activity

The antioxidant activity of different extracts from different organs of *C. longa*, the relationship between the polyphenolic content and the antioxidant activity, and the relationship between total polyphenol content and/or anti-scavenging activity and the anti-sickling activity is presented in the Table 2.



**Table 4. Relationship between polyphenol contents, antioxidant and anti-sickling activities**

Extracts from plants	Total polyphenol content (mg/EAG) (n=3)	Antioxidant activity		Anti-sickling activity /M.E (MNC en µg/mL)	Anti-sickling activity /A.E (MNC en µg/mL)
		ABTS (µg/mL)	DPPH (µg/mL)		
<b>Rhizomes</b>	12.82 ± 0.63	4.43 ± 0.18	15.63 ± 7.81	62.50	31.25
<b>Roots</b>	5.91 ± 0.87	4.66 ± 0.34	-	62.50	62.50
<b>Leaves</b>	4.31 ± 0.68	14.93 ± 4.30	54.83 ± 19.35	31.25	15.62
<b>Petals</b>	1.53 ± 0.28	24.27 ± 1.43	553.35 ± 250.15	62.50	62.50
<b>Sepals</b>	0.59 ± 0.30	46.24 ± 2.51	1129.80 ± 559.64	62.50	31.25

Legend: MNC: Minimum Normalization Concentration, M.E.: Methanol extracts, E.A.: Aqueous Extract, n= number of repetition of the experiment

The ANOVA revealed a very highly significant difference regarding the antioxidant activity (ABTS test) of *C. longa* extracts (p value = 0.0001). The Tukey pair multiple comparison test (HSD = 0.5365) does not reveal a significant difference between rhizome and root extracts. The intensity of anti-scavenging activity using the ABTS test and follows the following descending order: Rhizomes, roots > leaves > petals > sepals. However, the ABTS method helps to evaluate the antioxidant activity of the substance mixtures but also to distinguish their additive and synergistic effects. Because the radical ABTS reacts at the same time with the lipophilic and hydrophilic compounds (in particular the polyphenols) while the DPPH reacts only with the polar or hydrophilic compounds [68,78]. Thus, the fact that the rhizomes and roots of *C. longa* showed better activity with the ABTS test proves that these organs have a high content of lipophilic (non-polyphenolic compounds) and hydrophilic (polyphenolic compounds) compounds responsible for the antioxidant activity. It could also demonstrate that the anti-scavenging activity of rhizome and root extracts is related to the additive or synergistic effect of polyphenolic and non-polyphenolic compounds. The ANOVA on the antioxidant activity of extracts of different organs of *C. longa* using DPPH test showed a very highly significant difference in the antioxidant activity of these organs (p value= 0, 0001). However, the Tukey multiple-pair comparison test indicated that there is no significant difference between rhizome and leaf extracts. The anti-scavenging activity of extracts from different parts of *C. longa* follows the following descending order: Rhizomes> leaves > petals > sepals. It seems that the antioxidant activity of *C. longa* leaves is largely related to polyphenolic compounds because the DPPH radical reacts particularly with hydrophilic compounds i.e. polyphenols [68,78].

In addition, for the same sample, the analysis of the antioxidant activity using the ABTS test showed in most cases lower IC<sub>50</sub> values than those of the DPPH test. This asserts the hypothesis on the additive or synergistic effect of polyphenolic and non-polyphenolic compounds in the antioxidant activity of extracts studied and revealed that non-polyphenolic compounds contribute significantly to their antioxidant property. Maizura et al. [23] also reported the synergistic potential of the polyphenolic and non-polyphenolic fraction on the antioxidant potential of plant extracts. Several studies have shown that rhizome extracts of *C. longa* exhibited an antioxidant activity; and this antioxidant potential is mainly due to the presence of polyphenolic compounds, like curcuminoids. [21-22,24-25,27, 36,41-42,46-47,53,62,64,74,79-87]. According to these authors, curcuminoids notably Curcumin is ten times more antioxidant than Vitamin E.

By the way, Stanojevicv et al. [37], reported that the essential oils extracted from *C. longa* rhizomes also have a very strong antioxidant activity. There is little information in the literature on the antioxidant activity of root, leaves and floral extracts. The antioxidant activity of *C. longa* leaf, roots and flower extracts was reported as well by Camatari et al. [26], Ajay et al. [33] and Ritwiz et al. [48].

The comparison of total polyphenol content and the antioxidant activity is presented in Table 3.

The correlation between the total Polyphenol content and the antioxidant activity of ABTS and DPPH test of different extracts was evaluated using the Pearson test. The results obtained highlighted the existence of a negative correlation between the concentration of total polyphenols and the antioxidant activity of different extracts using the ABTS test (r = - 0.7807) but yet this correlation is not significant.

The analyses also showed a negative correlation between the concentration of total polyphenols content and the antioxidant activity using the DPPH test ( $r = -0.7445$ ). The non-significant negative correlation between total polyphenol content and the *in vitro* anti-scavenging activity of the ABTS and DPPH test indicated that the antiradical activity of the studied extracts was not related solely to the presence of the polyphenol compounds. Nevertheless, many studies have reported a positive correlation between the total polyphenol concentration of plant extracts and their antioxidant potential [21-22,26,33,35,71, 76]. Shan et al. [88] and Wu et al. [89], have also reported that phenolic compounds contribute significantly to the antioxidant activity of plant extracts. Velioglu et al. [90] and Kaur et al. [91], have also shown that there is a linear correlation between the total polyphenol content of plants and their antioxidant potential.

The relationship existing between the polyphenol content, antioxidant and anti-sickling activities is presented in Table 4.

Correlations between the total polyphenol content, antioxidant activity and the anti-sickling activity of the extracts were demonstrated using the Pearson test. The analyses carried out revealed: a positive correlation between (i) the total polyphenol contents and the anti-sickling activity of the studied extracts, in particular methanol extracts ( $r = 0.08328$ ) and (ii) between the antioxidant activity using ABTS and DPPH tests along with the anti-sickling activity of different extracts, in particular methanol extracts ( $r = 0.1281$  and  $0.4900$  respectively). However, these correlations were not significant. The positive correlation between total polyphenol contents and anti-sickling activity demonstrated that the polyphenolic compounds would contribute effectively at the same time to the inhibition of hemoglobin S polymerization and to erythrocyte sickling. However, the fact that this correlation (total polyphenol content and antioxidant activity of methanol extracts) was not significant suggests that polyphenolic compounds act in synergy with other active ingredients. Tshilanda [2] and Ngbolua et al. [7], reported that anthocyanins have the ability to inhibit the polymerization of hemoglobin S by engaging in a reaction with this protein.

In addition, the analysis of the correlation using Pearson test did not show a correlation between the total polyphenol content or antioxidant activity using ABTS test and the antioxidant activity of aqueous extracts ( $r = -0.1740$ ,  $r = -0.1481$

respectively). Although the correlation between the antioxidant activity of the extracts using DPPH test and the antioxidant activity of aqueous extracts was positive ( $r = 0.3114$ ). This would predict the involvement of the polyphenolic compounds of *C. longa* and their antioxidant potential on their anti-sickling potential; because the DPPH radical reacts exclusively with the polyphenolic compounds. However, in-depth and especially *in vivo* investigations are necessary to elucidate the implications of plant antioxidants on their anti-sickling potential.

Furthermore, the fact that the results of the present study showed a positive correlation between the antioxidant power of the extracts and their anti-sickling potential would reveal the involvement of the extract antioxidants in the normalization of erythrocytes *in vitro*. Considering these findings, Curcuma is a good candidate for the management of sickle cell disease because it has both anti-sickling and antioxidant activities and because of the involvement of oxidative stress in the pathophysiology of sickle cell disease. According to Ngbolua et al. [7], anthocyanins are effective scavengers of free radicals and therefore potent inhibitors of lipid peroxidation and can prevent the lysis of erythrocytes through their antioxidant properties. A phytomedicine based on these compounds could ensure the protection of the membrane in sickle cell patients by supplementing the enzymatic defense systems of the erythrocyte, which are defective in sickle cell patients [1].

Several studies reported that some polyphenolic compounds, notably curcumin extracted from *C. longa* rhizomes have the ability to trap free radicals (superoxide anions, hydroxyl radicals,  $H_2O_2$ , nitrite radical, etc.), inhibit or regulate the inducible enzyme for the synthesis of nitrite oxide (NO), iNO-Synthetase (iNOS), prevent lipid peroxidation, allow the chelation of metals, maintain the activity of antioxidant enzymes such as superoxide dismutase, catalase, and glutathione peroxidase [20,27,32,92,93,94-96]. Hierso [1] reported that administration of N-acetylcysteine, a non-essential amino acid stimulating glutathione production, would significantly decrease the formation of sickle red blood cells and return them to their biconcave form.

#### 4. CONCLUSION AND PERSPECTIVES

In the current study was highlighted the distribution of polyphenolic compounds in

different organs (rhizome, root, leaf, petal and sepal) of *C. longa*, evaluated the anti-sickling activity of aqueous and methanol extracts of *C. longa* and their antioxidant potential. The findings showed that the polyphenolic compounds are not evenly distributed in different organs (Rhizome, roots, leaves, floral parts) of *C. longa*. Total polyphenol content was higher in the rhizomes followed by roots, leaves, petals and sepals. The flavonoid content is higher in the leaves followed by rhizomes, roots, sepals and petals, while the anthocyanin content is highly in the leaves then rhizomes and other parts are poor in anthocyanins. Aqueous extracts of rhizomes, leaves and sepals were more active than methanol extracts of the same organs. Moreover, it was also found that different organs of this plant do not all have a good antioxidant potential, rhizome and root extracts showed better antiradical activity using ABTS method compared to other organs, while rhizome and leaf extracts showed better antiradical activity using the DPPH test.

A positive correlation between the total polyphenol content and the *in vitro* anti-sickling activity of the methanol extracts, and on the other hand between the *in vitro* antioxidant activity and anti-sickling activity was observed. In view of the involvement of oxidative stress in the pathophysiology of sickle cell disease, it is necessary to carry out *in vivo* studies not only to prove the anti-sickling activity of the polyphenolic compounds of *C. longa* but also to highlight the involvement of their antioxidant potential in the pathophysiology of this disease.

Studies on the anti-sickling activity of aqueous extracts from different parts of *C. longa* or those analyzing the correlation between total polyphenolic content and/or the correlation between the antioxidant and anti-sickling activities have not been reported yet in the literature. Further studies are required in order to (i) better understand the roles of plant antioxidants (especially polyphenols) in improving the clinical signs of sickle cell disease and their *in vivo* mechanisms of action and (ii) identify the active ingredients responsible for the anti-sickling activity.

## ACKNOWLEDGEMENT

Authors thank International Foundation for Science (IFS) for grant. One of the authors (Pius T Mpiana) thanks the TWAS and the Swedish

International Development Agency (SIDA) for the grant.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Hierso R. Implication du stress oxydant dans la pathologie de la drépanocytose: Crises vaso-occlusives, taux d'anticorps anti-bande 3 et oxydation du globule rouge. These, faculte de sciences exactes et naturelles. Universite Des Antilles et De Guyane. 2015;185.
2. Tshilanda DD. Etude phytochimique et activité antidrépanocytaire de trois espèces du genre *Ocimum*: *Ocimum basilicum* L., *O. canum* Sims, *O. gratissum* L. (Lamiaceae). These De Doctorat, Universite De Kinshasa, Faculte Des Sciences, Departement De Chimie. 2016; 274.
3. Janky E, Etienne-Julan M, Kadhel PH, Leborgne-Samuel Y, Melki E. Drépanocytose et grossesse. Extrait des mises à jour en Gynécologie et obstétrique, Tome XXX; 2006.
4. Ravelojaona M. Analyse histologique des répercussions musculaires, structurales, énergétiques et microvasculaires chez des hommes et des femmes drépanocytaires. These De Doctorat, Faculte De Medecine Jacques Lisfranc, Universite Jean Monnet Saint Etienne. 2015;206.
5. Mulungulu DN, Mpiana PT, Mbayo M, Tshisand PP, Badibanga LM. Etude de l'activité antidrépanocytaire et de la thermo-dégradation des extraits bruts aqueux, méthanoliques et éthanoliques d'*Ipomoea batatas*, une plante comestible à vertu thérapeutiques. International Journal of Innovation and Applied Studies. 2015;11(3):684-690. Available:<http://www.ijias.issr-journals.org>
6. Mpiana PT, Mudogo V, Tshibangu DST, Shetonde OM, Ngbolua KN, Mbala BM. Antisickling activity of some Congolese plants. In: Drug discovery from African flora. The 12th Symposium of the Natural Product Research Network for Eastern and Central Africa, July 22-26; University of Makerere, Kampala, Uganda. 2007d; 45(PS-6).

7. Ngbolua KN, Mudogo V, Mpiana PT, Malekani MJ, Rafatro H, Urvergratsimamanga S, Takoy L, Rakotoarimana H, Tshibangu DST. Evaluation de l'activité antidrepanocytaire et antipaludique de quelques taxons végétaux de la République Démocratique du Congo et de Madagascar. *J. Esthnopharmacologia*. 2013;5:19.
8. Mbadiko CM, Ngbolua KNN, Mpiana PT, Tshilanda DD, Makengo GK, Pambu AL, Kemfine LL, Bamvingana CK, Bongo GN, Mbemba FT. Assessment of the antisickling activity of total methanol extracts from the rhizomes and roots of *C. longa* and the effect of photodegradation on the antisickling activity. *J. of Advancement in Medical and Life Sciences*. 2017;5(1):1-5.
9. Mbadiko MM, Ngbolua KN, Mpiana PT, Mutambel H, Kikakedimau RN, Makengo GK, Pambu AL, Kemfine LL, Bongo GN, Mbemba FT. Phytochemical screening and assessment of anti-sickling activity of total methanol extracts of different organs of *Curcuma longa* L. (Zingiberaceae). *The Pharmaceutical and Chemical Journal*. 2017;4(1):32-40.
10. Mpiana PT, Mudogo V, Ngbolua KN, Tshibangu DST, Shetonde OM, Mbala MB. *In vitro* antisickling activity of anthocyanins from *Ocimum basilicum* L. (Lamiaceae). *International Journal of Pharmacology*. 2007;3(4):371-374.
11. Mpiana PT, Mudogo V, Tshibangu DST, Ekitwa EK, Kanangila AB, Lumbu BS, Ngbolua KN, Atibu EN, Kakule MK. Antisickling activity of anthocyanins from *Bombax pentadrum*, *Ficus capensis* and *Ziziphus mucronata*: Photodegradation effect. *Ethnopharmacology*. 2008;120:413-418.
12. Mpiana PT, Kitwa EK, Kanangila AB, Tshibangu DS, Ngbolua KN, Atibu EK, Kalonda EM. Influence of photodegradation on the anti-sickling activity of anthocyanins extracted from a few plants in Lubumbashi (DR Congo). *Ann. Fac. Sci*. 2009;45-53.
13. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Tshilanda DD, Atibu EK. Antisickling activity of anthocyanins of *Jatropha curcas* L. *Recent Progress in Medicinal Plants*. 2009;25:101-108.
14. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Atibu EN, Kitwa EK, Kanangila AB. Antisickling activity of anthocyanins extracts of *Vigna unguiculata* (L.) Walp. *Recent Progress in Medicinal Plants*. 2009;25:91-98.
15. Mpiana PT, Tshibangu DST, Shetonde OM, Ngbolua KN. *In vitro* antidrepanocytary activity (anti-sickle cell anemia) of some Congolese plants. *Phytomedicine*. 2007;14:192-195.
16. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Shetonde OM, Mangwala KP, Mavakala BK. *In vitro* antisickling activity of anthocyanins extract of a Congolese plant: *Alchornea cordifolia* M. Arg. *J. Med. Sci*. 2007;7(7):1182-1186.
17. Mpiana PT, Makelele LK, Oleko RW, Bokota MT, Tshibangu DST, Ngbolua KN, Mbala MB, Atibu EK, Nsimba SM. Antisickling activity of medicinal plants used in the management of Sicklec cell Disease in Tshopo district, D.R. Congo. *Australian Journal of Medical Herbalism*. 2010;22(4):132-137.
18. Mpiana PT, Makelele LK, Oleko RW, Bokota MT, Tshibangu DST, Ngbolua KN, Mbala MB, Atibu EK, Nsimba SM. Antisickling activity of medicinal plants used in the management of Sicklec cell Disease in Tshopo district, D.R. Congo. *Australian Journal of Medical Herbalism*. 2010;22(4):132-137.
19. Hombourger C. Le Curcuma de l'épice au médicament. These De Doctorat, Université Henri Poincaré-Nancy 1, Faculté De Pharmacie. 2010 ;222. Available:<http://www.culture.gouv.fr/culture/infos-pratique/droits/protection.htm>
20. Tilak JC, Banerjee M, Hari M, Devasagayam TPA. Antioxidant availability of turmeric in relation to its medicinal and culinary uses. *Phytother. Res*. 2004;18:798–804.
21. Shiyou L, Wei Y, Guangrui D, Ping W, Peiying Y, Aggarwal BB. Chemical composition and product quality control of turmeric (*Curcuma longa* L.). *Pharmaceutical Crops*. 2011;(2)29.
22. Tanzeela N, Muneeb I, Ahmad R, Madiha S, Fatima I, Marwa W. Turmeric: A promising spice for phytochemical and antimicrobial activities. *American-Eurasian J. Agric. & Environ. Sci*. 2015;15(7):1278-1288.
23. Maizura M, Aminah A, Wan AWM. Total phenolic content and antioxidant activity of kesum (*Polygonum minus*), ginger (*Zingiber officinale*) and turmeric (*Curcuma*

- longa*) extract. International Food Research Journal. 2011;18:529-534.
24. Prafulla S, Arjun M, Vidya S. *Curcuma longa* Linn. A phytochemical and phytopharmacological review. Research Journal of Pharmacognosy and Phytochemistry. 2013;5(2):59-68.
  25. Roshan PY, Gaur T. Versatility of turmeric: A review the golden spice of life. Journal of Pharmacognosy and Phytochemistry. 2017;6(1):41-46.
  26. Camatari FOS, Lopes KH, Valentim BI, Xavier JA, da Costa JG, Santana AEG, Goulart MOF. Antioxidant potential of flours from cereals, tubers, beans and seeds chemical profile of *curcuma longa* flour. Journal of Nutrition & Food Sciences. 2016;2(2):1-8.
  27. Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: A component of turmeric (*Curcuma longa*). J. Altern. Complement Med. 2003;9(1):161-168.
  28. Harshal P, Mugdha K, Nilesh M, Pravin JK. Phytochemical evaluation and curcumin content determination of turmeric rhizomes collected from bhandara district of Maharashtra (India). Medicinal chemistry. 2014;4(8):588-591.
  29. Mohammad NH, Aysha F, Nargis A, Shahnaz R, Shahadat HMD, Mohammed R. Preliminary phytochemical screening, toxicity, antihyperglycemic and analgesic activity studies with *Curcuma longa* leaves. World Journal of Pharmacy and Pharmaceutical Sciences. 2014;3(9):81-91.
  30. Rohini S, Mehta A, Mehta P, Shukla K. Anthelmintic activity of rhizome extracts of *Curcuma longa* and *Zingiber officinale* (Zingiberaceae). Int J Pharm Pharm Sci. 2011;3(2):236-237.
  31. Jansen PCM, Grubben GJH, Cardon D. Ressources végétales de l'Afrique tropicale 3. Colorants et tanins. Wageningen. Pays-Bas: PROTA. 2005 ; 238.
  32. Vaquier ARL. Intérêt d'un nouveau nutriment a visée anti-inflammatoire dans la gestion de troubles locomoteurs de cheval. Aspects bibliographiques et études cliniques. Ecole Vétérinaire d'Alfort, Faculté de Médecine de Cretal. 2010;199.
  33. Ajay K, Monika S, Prem PS, Sandeep KS, Pratima R, Kapil PD. Antioxidant efficacy and curcumin content of turmeric (*Curcuma-longa* L.) Flower. International Journal of Current Pharmaceutical Research. 2016;8(3):112-114.
  34. Kholi K, Ali J, Ansari MJ, Raheman Z. Curcumin: A natural anti-inflammatory agent. Indian J. Pharmacol. 2005;37(3): 141–147
  35. Yan SW, Asmah R. Comparison of total phenolic contents and antioxidant activities of turmeric leaf, pandan leaf and torch ginger flower. International Food Research Journal. 2010;17:417-423.
  36. Rafie H, Soheila H, Mohsen H, Mahnaz S, Roxanna H. Turmeric (*Curcuma longa*): from a variety of traditional medicinal applications to its novel roles as active antioxidant, anti-inflammatory, anti-cancer, and anti-diabetes. International Journal of Pharmacology, Phytochemistry and Ethnomedicine. 2015;1:37-45.
  37. Stanojević JS, Stanojević LP, Cvetković DJ, Danilović BR. Chemical composition, antioxidant and antimicrobial activity of the turmeric essential oil (*Curcuma longa* L.). Advanced Technologies. 2016;4(2):19-25.
  38. Ajaiyeoba EO, Sama W, Essien EE, Olayemi IO, Ekundayo OT, Walker M, Setzer WN. Larvicidal activity of turmerone-rich essential oils of *Curcuma longa* leaf and rhizome from Nigeria on *Anopheles gambiae*. Pharmaceutical Biology. 2008;46(4):279–282.
  39. Fathima ND. Curcumin -Role in Cancer. Pharm. Sci. & Res. 2014;6(11):351-355.
  40. Essien EE, Schmidt JN, Tameka MW, Setzer WN, Olusegun E. Chemotaxonomic characterization and in-vitro antimicrobial and cytotoxic activities of the leaf essential oil of *Curcuma longa* grown in Southern Nigeria. Medicines. 2015;2(4):340-349. Available:<https://www.ncbi.nlm.nih.gov>
  41. Rajesh H, Rao SN, Rani MN, Prathima KS, Rejeesh EP, Chandrashekar R. Phytochemical analysis of methanol extract of *Curcuma longa* Linn rhizome. International Journal of Universal Pharmacy and Bio Sciences. 2013;2(2):39-45.
  42. Savina P. Le Curcuma, un agent naturel de lutte contre le vieillissement cutané. Synthèse Bibliographique en Biologie et Biotechnologie–Master Biologie et Gestion, Université de Rennes 1 UFR Sciences de la Vie et de l'Environnement. 2014;34.
  43. Arutselvi R, Balasaravanan T, Ponmurugan P, Saranji MN, Suresh P. Phytochemical screening and comparative study of anti-microbial activity of leaves and rhizomes of turmeric varieties. Asian

- Journal of Plant Science and Research. 2012;2(2):212-219.
44. Seema PK. Comparative study of pharmacognostical and preliminary phytochemical investigation of curcuma longa leaves and rhizomes. *Pharma Tutor*. 2016;4(10):31-36.
  45. Chairman K, Jayamala M, Vijila CR, Ranjit SAJA. Phytochemical screening and antimicrobial activity of *Curcuma longa*. *Natural Dye. Med*. 2015;3(2).
  46. Wagner de ST, Yasmin A, Padoan GLG, Cola JZ, Murray BI. Turmeric powder and its derivatives from *Curcuma longa* rhizomes: Insecticidal effects on cabbage looper and the role of synergists. *Scientific Reports*. 2016;6:1-11.
  47. Panpatil V, Shalini T, Nirmala K, Chetan N, Kalpagam P. *In vitro* evaluation on antioxidant and antimicrobial activity of spice extracts of ginger, turmeric and garlic. *Journal of Pharmacognosy and Phytochemistry*. 2013;2(3):143-148.
  48. Ritwiz BS, Kusha BS, Ranjeet D, Ganesh N. *Curcuma longa* leaves exhibits a potential antioxidant, antibacterial and immunomodulating properties. *International Journal of Phytomedicine*. 2011; 3(2011):270-278
  49. Ancy AR, Salini P, Antony S. Phytochemical screening and comparative study of antimicrobial activity of leaves and rhizomes of turmeric varieties. *International Journal of Research in Plant Science*. 2017;7(1):7-11.
  50. Sawant RS, Godghate AG. Qualitative phytochemical screening of rhizomes Of *Curcuma longa* Linn. *International Journal of Science, Environment and Technology*. 2013;2(4):634–641.
  51. Sindhu S, Chempakam B, Leela NK, Suseela BR. Chemoprevention by essential oil of turmeric leaves (*Curcuma longa* L.) on the growth of *Aspergillus flavus* and aflatoxin production. *Food Chem Toxicol*. 2011;49(5):88-92.
  52. Imma C, Vitulano M, Esterina C, D'Onofrio F, Lorenzo A. Extraction and curcuminoids activity from the roots of *Curcuma longa* by rslda using the naviglio extractor. *European Scientific Journal Special edition*. 2016;1857–7881.
  53. Anbujeba J, Suraj SR, Renitha G, Anandrarajagopal K, Gnamakumari AAV, Promwichit P. *In vitro* antimicrobial evaluation of Zingibera ffinale, *Curcuma longa* and *Alpinia galangal* extracts as natural food preservatives. *America Journal of Food Technology*. 2002;4(5):192-200.
  54. Araujo CC, Leon LL. Biological activities of *Curcuma longa* L. *Mem Inst Oswaldo Cruz*. 2001;96(5):723-728.
  55. Bruneton J. *Pharmacognosie, phytochimie, plantes médicinales*, 4er éd., Revue et Augmentée, T 6 et Doc. Editions médicinales internationales, Paris. 2009; 1288.
  56. Cheikh ZA. *Etudes chimiques et biologiques d'Aframomum sceptrum* (Zingibéraceae) et de la curcumine. Université paris-sud, Faculté de pharmacie de chatnay-Melabry. 2013;161. Available:<http://www.tel.archives-ouvertes.fr>
  57. Goel A, Kunnumkkara B, Aggarwal BB. Curcumin as «Curecumin: From kitchen to clinic. *Biochem.Pharmacol*. 2008;75:787–809.
  58. Nada GAL, Talet Y, Nayyarjehan M, Samia A. Studies of the antibacterial effect of different fraction of *Curcuma longa* against urinary tract infections Isolation. *Pakistan Journal of Biological Sciences*. 2004;7(12): 2055-2060. Available:<http://www.Scialert.net>
  59. Niamsa N, Sittwet C. Antimicrobial activity of *Curcuma longa* Aqueous Extract. *Journal of pharmacology and toxicology*. 2009;4(4):173-177. Available:<http://www.Scialert.net>
  60. Perry MC. Evaluation de la Curcumine comme agent anticancéreux dans le traitement des tumeurs cérébrales. *Mémoire De Master en Chimie, Université Du Québec à Montréal*. 2008;97.
  61. Sivabalan S, Anuradha CV. A comparative study on the antioxidant and glucose lowering effects of Curcumin Analogue trough in vitro essay. *International Journal of Pharmacology*. 2010;6(5):664-669. Available:<http://www.Scialert.net>
  62. Gupta SK, Lather A, Jaiswall V, Garg S, Kumar JA. Phytochemistry of *Curcuma longa* - An overview. *Journal of Pharmaceutical and Biomedical Sciences*. 2010;4(1):1-9.
  63. Awasthi PK, Dixit SC. Chemical composition of *Curcuma longa* leaves and rhizome oil from the plains of Northern India. *J Young Pharm*. 2009;1(4):312-316.
  64. Shawket DS. Screening the antibacterial potency of *Curcuma longa* L. essential oil extract against boils causing

- Staphylococcus* species. International Journal of Advanced Biological Research. 2013;3(4):490-500.
65. Voravuthikunchai SP. Family zingiberaceae compounds as functional antimicrobials, antioxidants, and antiradicals. World Journal of Medicine and Medical Science. 2007;1(2):227-240.
  66. Jurenka JS. Anti-inflammatory properties of curcumin, a major constituent of *Curcuma longa*: A review of preclinical and clinical research. Altern. Med. Rev. 2009; 14(2):141–153.
  67. Lassa KL. Valorization of non-timber forest products (NTFPs) of plant origin sold in markets and their availability in the surroundings of Kinshasa city: "Case of CADIM". Unpublished Master Dissertation, Department of Biology, University of Kinshasa. 2012;285.
  68. Kapepula PM, Mbombo PM, Franck F, Mouithys-Mickalad A, Mumba ND, Dibungi KP, Kabamba NN, Tits M, Frédéric M, Tamfum MJJ. Antioxidant potentiality of three herbal teas consumed in bandundu rural areas of Congo. Natural Product Research. 2017;31(16):1-4.
  69. Adedapo AA, Jimoh FO, Koduru S, Afolayan AJ, Masika PJ. Antibacterial and antioxidant properties of the methanol extracts of the leaves and stems of *Calpurnia aurea*. BMC Complementary and Alternative Medicine. 2008;8(1):1–8.
  70. Tula YM. Anti-sickling activity and stability of anthocyanins extracted from *Hypoxis angustifolia* bulbs and leaves of *Ipomoea batatas*. Unpublished BSc dissertation, Department of Chemistry, Faculty of Sciences, University of Kinshasa; 2016.
  71. Gayatri N, Rajani KS. Evaluation of antioxidant activity in ethanolic extracts of five *Curcuma* species. International Research Journal of Pharmacy. 2011; 2(12):243-248.
  72. Chang CH, Lin HY, Chang CY, Liu YC. Comparisons on the antioxidant properties of fresh, freeze-dried and hot-air-dried tomatoes. Journal of Food Engineering. 2006;77:478-485.
  73. Nilanjana D, Purba M, Ajoy KG. Pharmacognostic and phytochemical evaluation of the Rhizomes of *Curcuma longa* Linn. Journal of Pharma Sci Tech. 2013;2(2):81-86.
  74. Tiwari NB, Jadhav SK. Comparative phytochemical screening of bioactive compounds in *Curcuma caesia* Roxb. and *Curcuma longa*. Research Journal of Medical Plant. 2013;7(2):113-118.
  75. Pawar MA, Patil SS, Nagrik DM. Phytochemical and physicochemical investigation of *Curcuma longa* Linn Rhizome. International Journal of Chemical and Physical Sciences. 2015;4:458-463.
  76. Cousins M, Aldelberg J, Chen J, Rick F. Antioxidant capacity of fresh and dried rhizomes from four clones of turmeric (*Curcuma longa* L.) grown in vitro, 2007;2(2):129-135.
  77. Walker V. Impact of inoculation of phytobeneficial microorganisms on the secondary metabolism of *Zea mays*. Unpublished PhD dissertation, University of Claude Bernard Lyon 1; 2013.
  78. Yizhong C, Qiong L, Mei S, Harold C. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. Life Sciences. 2003;74:2157-2184.
  79. Akram M, Shahab U, Afzal A, Khan U, Abdul H, Mohiuddin E, Asif M. *Curcuma longa* and curcumin: A review article. J. Biol. – Plant Biol. 2011;55(2):65–70.
  80. Singh RP, Jain DA. Evaluation of antimicrobial activity of curcuminoids isolated from turmeric. Int. J. of Pharm. & Life Sci. (IJPLS). 2012;3(1):1368-1376.
  81. Lawhavinit O, Kongkathip N, Boonsong K. Antimicrobial activity of curcuminoids from *Curcuma longa* L. on Pathogenic Bacteria of Shrimp and Chicken. Kasetsart J. 2010;44:364–371.
  82. Shanmugam MK, Rane G, Madhu MK, Arfuso F, Arunachalam C, Zayed ME, Sulaiman AA, Benny KHT, Kumar AP, Gautam S. The multifaceted role of curcumin in cancer prevention and treatment. Molecules Journal. 2015;20:2728-2769.
  83. Virendra PV, Shalini T, Nirmala K, Nimgulkar C, Kalpagam P. *In vitro* evaluation on antioxidant and antimicrobial activity of spice extracts of ginger, turmeric and garlic. Journal of Pharmacognosy and Phytochemistry. 2013;2(3):143-148.
  84. Kumar S, Singh NN, Singh A, Singh N, Sinha RK. Use of *Curcuma longa* L. extract to stain various tissue samples for histological studies. Ayu Journal. 2014;35(4):447-451.
  85. Naama JH, Temimi AA, Al-Amiery AHA. Study the anticancer activities of ethanolic curcumin extract. African Journal of

- Pure and Applied Chemistry. 2010;4(5):68-73.
86. Seggani S, Boukehil H. Correlation between the polyphenolic content and the antioxidant and antimicrobial activity *in vitro* of *Curcuma longa* L. Unpublished MSc dissertation, Université des Frères Mentouri Constantine, Faculty of Natural Sciences and Life, Democratic Republic and Popular of Algeria. 2017;108.
87. Boutamani M. Etude de la variation du rendement et de la composition chimique du *Curcuma longa* et *Myristica fragrans* en fonction du temps et de la technique utilisée. Mémoire de Master, Université Des Sciences et De La Technologie Houari Boumediene Alger, Chimie De Médicament; 2013. (Mémoire Online.net).
88. Shan B, Cai YZ, Sun M, Corke H. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *Journal of the Agricultural and Food Chemistry*. 2005;53:7749–7759.
89. Wu CQ, Chen F, Wang X, Kim HJ, He GQ, Haley- Zitlin V, Huang G. Antioxidant constituents in feverfew (*Tanacetum parthenium*) extract and their chromatographic quantification. *Food Chemistry*. 2006;96:220–227.
90. Velioglu YS, Mazza G, Gao L, Oomah BD. Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *Journal of Agricultural Food Chemistry*. 1998;46:4113-4117.
91. Kaur C, Kapoor HC. Antioxidant activity and total phenolic content of some Asian vegetables. *International Journal of Food Science and Technology*. 2002;37:153-161.
92. Naama JH, Temimi AA, Al-Amiery AHA. Study the anticancer activities of ethanolic curcumin extract. *African Journal of Pure and Applied Chemistry*. 2010;4(5):68-73.
93. Omodamiro OD, Ekeleme CM. Comparative study of *in vitro* antioxidant and antimicrobial activities of *Piper guineense*, *Curcuma longa*, *Gongronemalati folium*, *Allium sativum*, *Ocimum gratissimum*. *World Journal of Medicine and Medical Science*. 2013; 1(4):51-69.
94. Jayaprakasha GK, Jagan MRAOL, Sakariah KK. Chemistry and biological activities of *C. longa*. *Trends Food Sci. Tech*. 2005;16(12):533–548.
95. Anand P, Kunnumakkara AB, Newman RA, Aggarwal BB. Bioavailability of curcumin: Problems and promises. *Mol. Pharm*. 2007;4(6):807–818.
96. Camacho-Barquero L, Villegas I, Sanchez-Calvo JM, Talero E, Sanchez-Fidalgo S, Motivla V, Catalina A. Curcumin, a *Curcuma longa* constituent, acts on MAPK p38 pathway modulating COX-2 and iNOS expression in chronic experimental colitis. *Int. Immunol*. 2007;7(3):333-342.

© 2019 Mbadiko et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sdiarticle3.com/review-history/46718>