

Risk Assessment of Heavy Metal Contaminants in Some Common Ready to Eat Meals Sold at Petrol Stations in Calabar Metropolis

**G. E. Udosen^{1*}, F. E. Uboh², Dennis Amaechi³, M. O. Odey⁴
and J. E. Mgbang²**

¹Department of Nursing Sciences, Faculty of Allied Medical Sciences, University of Calabar, Nigeria.

²Department of Biochemistry, Faculty of Basic Medical Sciences, University of Calabar, Nigeria.

³Department of Biochemistry, College of Natural and Applied Sciences, Veritas University, Bwari, Abuja, Nigeria.

⁴Department of Biochemistry, Faculty of Basic Medical Sciences, Cross River State University of Science and Technology, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Carla Maria Ferreira Guerreiro da Silva Mendes, Nurse Director at Centro Hospitalar de Setúbal, Portugal.

Reviewers:

(1) Felagha Iniebiyo, University of Port Harcourt, Nigeria.

(2) C. I. A. Nwoko, Federal University of Technology Owerri, P.M.B. 1526, Nigeria.

(3) Dr. Selina Ama Saah, University of Energy and Natural Resources, Ghana.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48943>

Received 20 February 2019

Accepted 02 May 2019

Published 09 May 2019

Original Research Article

ABSTRACT

Aim: In this study, the concentrations of Cu, Fe and Ni, long term exposure and daily consumption of ready-to-eat food foods sold at Petrol station's Atmospheric conditions (AF) in Calabar Metropolis was evaluated.

Methods: Foods samples, including such prepared ready-to-eat foods as garri, meat pie, stew, rice, beans, afang and melon soups were collected at the point of sale at the fuel stations, about 7:00 am in the morning before they were opened for sale (and exposed to the environment of the filling stations). These were labelled "Before". At about 2:00 pm to 3:00 pm same day, the same ready-to-eat food samples were collected again at the same spots (after they have been exposed to

the filling stations atmospheric environment). These were labelled "After". The food samples were collected in such a way that all the Garri purchased before, across the different filling stations were mashed together as a single sample, while the ones purchased after were equally collected to form a single sample. The other food samples were handled in like manner.

Results: The levels of Cu, Fe and Ni recorded for garri, afang soup, melon soup, white rice, beans, stew and meat pie (ready-to-eat meals) . The results showed that the level of Ni was significantly high in garri, afang soup, melon soup, white rice, beans and stew, while the level of Fe was significantly ($p < 0.05$) increased in garri, melon soup, white rice, beans, stew and meat pie after 6 hours of exposure to petrol station's atmospheric conditions. It was also observed that the level of Cu was significantly high in garri and white rice after 6 hours of exposure to petrol station's atmospheric conditions. This also gave an indication exposure of most ready-to-eat foods to the petrol stations atmospheric conditions for 6 hours, during sales, are likely to results in the contamination of the foods with Ni.

Conclusion: In this study, such ready-to-eat foods as garri, afang soup, melon soup, white rice, beans, stew and meat pie displayed for sale at the petrol refueling stations in Calabar were observed to be contaminated with Cu, Fe and level of Ni.

Keywords: Copper; iron; nickel; environmental and contaminants.

1. INTRODUCTION

Ready-to-Eat foods are important because they are readily available to both high and low income workers in most Nigerian cities. They are very economical and are sold on the streets of the major urban cities around the country [1]. The fact they are sold in the open on street sides, exposes these foods to certain contaminants in the course of preparation, processing and sales of these foods. It has been reported that hundreds of chemical pollutants, including heavy metals, are discharged into the environment in the course of various human activities [1]. The presence of heavy metals in ready-to-eat foods could make these foods to become environmental hazards to human health. Consumption of these foods over time usually leads to build-up of heavy metals which may therefore lead to retention of the metals in the body, thereby producing some kind of hypertoxicity effects [1]. Various human activities are known to release heavy metals to the different environments, including the atmosphere. Heavy metals are therefore chemical substances that occur naturally, or are released into the environments by several human and industrial activities. Such metals as arsenic, cadmium, lead and mercury can be present at various levels in environments, e.g. soil, water and atmosphere. The metallic load of food crops depends mainly on their uptake by the plant, ability to induce toxicity, and bio-accumulate in the food chain [2]. Food has been reported to constitute the principal source of cadmium in non-smokers. Different mean daily cadmium intake from food have been reported in such

places like Croatia, Czech Republic, France, Poland, Germany and Spain [3,4].

Some of the ready-to-eat foods are commonly sold along petrol stations, motor parks, mechanic workshops and roadsides of most cities in Nigeria include roasted plantain, stick meat, rice, beans, porridge yams and soups. Various occupational activities at the petrol stations, motor parks, mechanic workshops and traffic congested roadsides release some chemical pollutants, including heavy metals and hydrocarbons into the environment. Exposure of these ready-to-eat foods to the atmosphere at the petrol stations, motor parks, mechanic workshops and traffic congested roadsides is likely to introduce atmospheric chemical contaminants, such hydrocarbons and heavy metals pollutants, into these foods. The deposition of heavy metals are potentially poisonous and toxic with its accumulation in human system [4].

Copper is to play important role in maintenance of several metabolic activities in the body. However, excessive intake of copper may produce adverse effect in the body. Hence, the recommended dietary intake of copper had been documented by the various National and International Regulatory Bodies. [5], 0.2 mg/kg/bw/day is recommended as the upper limit, a safe range of adults' standard consumption for copper. Therefore, the acceptable safe limit of copper recommended for adults by WHO is about 0.2mg/kg body weight/day.

High level of nickel has been reported in foods such as oatmeal and some legumes such as soybean and cocoa [6]. A comparatively higher level of about 900 µg/day of nickel has been reported in Denmark probably due to high dietary intake from these foods. In UK, Finland, US and Canada, "the mean total dietary intake of nickel has been reported to be between 0.12 - 0.21 mg, 0.13 mg, 0.17 mg, 0.207 - 0.406 mg respectively" [7,8,9,10]. These reports indicate that daily dietary intake of nickel varies on the basis of food type and quantity of food eaten. It can therefore be deduced the amount of nickel vary considerably with food type and from place to place.

2. MATERIALS AND METHODS

2.1 Materials

The following equipment and glass wares were used in the course of this research: Laboratory mortar (model EW-63100-60, from Cole-parmer company Ltd, USA), Evaporating plate (model SER-No.62, from Gallenkamp company Ltd, UK), Atomic Absorption Spectrophotometer (model AA6800, Schemadzu company, Japan).

2.1.1 Reagents/chemicals

Standard reagents and chemicals were used and include: Lichens coded International Atomic Energy Agency (IAEA-336), from Sigma, USA, Nitric acid (Riedel-deHaën, Germany), Perchloric acid (Sigma-Aldrich, Germany), Hydrofluoric acid, Ethanol, Methanol, Ethyl-acetate, (British Drug House Chemicals Ltd, Poole, England).

2.2 Methods

2.2.1 Collection of food samples

Foods samples, including such prepared ready-to-eat foods as garri, meat pie, stew, rice, beans, afang and melon soups were collected at the point of sale at the fuel stations, about 7:00 am in the morning before they were opened for sale (and exposed to the environment of the filling stations). These were labelled "Before". At about 2:00 pm to 3:00 pm same day, the same ready-to-eat food samples were collected again at the same spots (after they have been exposed to the filling stations atmospheric environment). These were labelled "After".

2.2.2 Preparation of food samples for heavy metal analysis

One gram each of the pulled samples was weighed into a beaker. 20ml of aqua-rega (a solution of Nitric acid and perchloric acid (3:1) was added and the beaker was covered with glass for the initial effervescence to subside. Thereafter, the beaker was placed on a hot plate and heated to near dryness at about 80-90°C. The aqua-rega was added as required in the course of digestion, to avoid drying. After the sample was fully digested, giving light coloured solution, the beaker was transferred onto a work bench and allowed to cool. The cooled sample was filtered into a beaker and made up to the mark with distilled water. This was transferred into a sample container in preparation for heavy metal (elemental) determination using Atomic Absorption Spectrophotometer [11].

2.2.3 Analytical quality assurance

To make sure that the analytical methods used for heavy metal determination are reliable, standard reference materials, Lichens coded (International Atomic Energy Agency; IAEA-336) were also digested and then analyzed using same procedure. Comparison of determined values with certified elemental values was carried out to ensure reliability of the analytical method used [12].

3. RESULTS

Results of different concentrations of Cu, Fe and Ni in some ready-to-eat meals before and after 6 hrs of exposure to petrol station's atmospheric conditions (AF).

4. DISCUSSION

The presence of heavy metals in the air, water and food forms a major health threat globally [13]. Such human activities as use of agricultural pesticides, increase in industrialization and mining are known to release high amount of heavy metals into the environment, thereby increasing the levels of heavy metal pollution in the ecosystem [14]. Consumption of food items displayed at these environments is therefore likely to expose the consumers to the risk of these heavy metals toxicity. Some heavy metals are known to cause various health hazards to




Fig. 1. Comparison of concentration of copper before and after exposure in the different food samples

Values are expressed as mean + SEM, n = 3
* = significantly different from before exposure at p < 0.05




Fig. 2. Comparison of concentration of iron before and after exposure in the different food samples

Values are expressed as mean + SEM, n = 3.
* = significantly different from before exposure at p < 0.05




Fig. 3. Comparison of concentration of nickel before and after exposure in the different food samples

Values are expressed as mean + SEM, n = 3.

* = significantly different from before exposure at $p < 0.05$

individuals that consume those foods that are enormously contaminated with the metals. Generally, heavy metals get into human systems following consumption of foods and drinking of water that are contaminated with heavy metals. This may also play a key role in evaluating food safety and the consequent effects of heavy metals on the consumers.

This study assessed the level of some heavy metals in ready-to-eat foods such as prepared garri, meat pie, stew, rice, beans, afang and melon soups that are sold at the petrol refueling stations in Calabar. According to the “Joint FAO/World Health Organization Expert Committee on Food Additives (JEFCA) the established provisional tolerable weekly intake for lead is 0.025 mg/kg body weight”. Also, the report of “WHO provisional guideline records 0.01 mg/L as the adopted standard for drinking water” [15]. According to the FAO/WHO standard, “the permissible level of cadmium and lead is 0.05 and 0.2mg/kg, respectively”.

The levels of Cu, Fe and Ni recorded for garri, afang soup, melon soup, white rice, beans, stew and meat pie (ready-to-eat meals) are presented in Figs. 1-3. The results showed that the level of

Ni was significantly high in garri, afang soup, melon soup, white rice, beans and stew, while the level of Fe was significantly ($p < 0.05$) increased in garri, melon soup, white rice, beans, stew and meat pie after 6 hours of exposure to petrol station’s atmospheric conditions. It was also observed that the level of Cu was significantly high in garri and white rice after 6 hours of exposure to petrol station’s atmospheric conditions. This also gave an indication exposure of most ready-to-eat foods to the petrol stations atmospheric conditions for 6 hours, during sales, are likely to results in the contamination of the foods with Ni.

5. CONCLUSION

In this study, such ready-to-eat foods as garri, afang soup, melon soup, white rice, beans, stew and meat pie displayed for sale at the petrol refueling stations in Calabar were observed to be contaminated with Cu, Fe and level of Ni. The results of this study therefore suggest that most ready-to-eat foods that exposed to prevailing environmental conditions at filling stations are likely to be contaminated with some heavy metals (such as Cu, Fe and Ni).

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Altindağ A, Yiğit S. Assessment of heavy metal concentrations in the food web of lake Beyşehir, Turkey. *Chemosphere*. 2005;60(4):552-6.
2. Hussain RH, Ebraheem MK, Moker HM. Assessment of heavy metals (Cd, Pb and Zn) contents in livers of chicken available in the local markets of Basrah City, Iraq. *Basrah Journal of Veterinary Resesearch*. 2012;11:43-51.
3. Sapunar-Postruznik J, Bazulic D, Kubalab H, Balint I. Estimation of dietary intake of lead and cadmium of the general population of Republic of Croatia, *Science of the Total Environment*. 1996;177:31-35.
4. Puklová V, Batáriová A, Černá M, Kotlík B, Kratzer K, Melicherčík J, Ruprich J, Řehůřková I, Spěváčková V. Cadmium exposure pathways in the Czech Urban population. *Central European Journal of Public Health*. 2005;13(1):11-19.
5. FAO/WHO. Report of the 32nd Session of the codex committee of the food additives Contaminants. Beijing People's Republic of China. 2000;20-24.
6. Han HJ, Lee BH, Park CW, Lee CH, Kang YS. A study of nickel Content in Korean Foods. *Korean Journal of Dermatology*. 2005;43:593-598.
7. Ysart G, Miller P, Crews H, Robb P, Baxter M, De L'Argy C. Dietary exposure estimates of 30 elements from the UK Total Diet Study. *Food Additives and Contaminations*. 1999;16:391-403.
8. Varo P, Koivistonen P. Mineral composition of Finnish foods XII. General discussion and nutritional evaluation. *Acta Agricultura Scandinavica*. 1980;S22:165-170.
9. Nielsen FH. Fluoride, vanadium, nickel, arsenic and silicon in total parenteral nutrition. *The Bulletin of the New York Academy of Medicine*. 1984;60:177-195.
10. Dabeka RW, MacKenzie AD. Survey of lead, cadmium, fluoride, nickel and cobalt in food composites and estimations of dietary intakes of these elements by Canadians. *Journal of AOAC International*. 1995;78:897-909.
11. Awofolu OR, Mbolekwa Z, Mtshemla V, Fatoki OS. Levels of trace metals in water and sediments from Tyume River and its effects on an irrigated farmland. *Water SA*. 2005;31(1):87-94.
12. Udiba UU, Gauje B, Ashade NO, Ade-Ajayi FA, Okezie VC, Aji BM, Agboun TDT. An assessment of the heavy metal status of River Galma around Dakace industrial layout, Zaria, Nigeria. *Merit Research Journal of Environmental Science and Toxicology*. 2014;2(8):176-184.
13. Matthew MM, Henke R, Atwood A. Effectiveness of commercial heavy Metal chelators with new insights for the future in chelate design. *Journal of Hazardous Materials*. 2002;92(2):129-142.
14. Srikanth R, Rao AM, Kumar CHS, Khanum A. Lead, cadmium, nickel, and zinc contamination of ground water around Hussain Sagar Lake, Hyderabad, India. *Bulletin of Environmental Contamination and Toxicology*. 2004;50(1):138-143.
15. WHO. Evaluation of certain food additives and contaminants. Sixty-first report of the Joint FAO/WHO Expert Committee on Food Additives, WHO Technical Report Series. 2004;922.

© 2019 Udosen et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sdiarticle3.com/review-history/48943>