



First Report of *Tetragonula calophyllae* Shanas & Faseeh 2019 (Apidae: Apinae: Meliponini) from Goa State, India

Arya Krishna ^a, Anitha N. ^a, Santhosh Kumar T. ^a,
Swapna Alex ^a, Pratheesh P. G. ^a and Shanas S. ^{b*}

^a College of Agriculture, Vellayani, Kerala Agricultural University, Kerala 695522, India.

^b Integrated Farming System Research Station, Kerala Agricultural University, Karamana, Thiruvananthapuram, Kerala 695002, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i13804

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111420>

Original Research Article

Received: 28/10/2023

Accepted: 04/01/2024

Published: 04/01/2024

ABSTRACT

The occurrence of *Tetragonula calophyllae* Shanas & Faseeh 2019, in Goa state, India, is reported for the first time. Morphometric comparison between *T. calophyllae* workers from Goa and Type series are provided along with comments on the external nest entrance. The need for developing baseline database on species distribution is stressed for developing policies for conservation planning and climate adaptation strategies for future.

Keywords: Stingless bees; nomenclature; taxonomy; distribution; diversity.

*Corresponding author: E-mail: shanassudheer@gmail.com;

1. INTRODUCTION

The stingless bees (Meliponini) are highly eusocial bees restricted to the tropical and subtropical areas of the world [1], represented by 605 extant species in 45 genera [2]. They are particularly important pollinators of tropical plants, visiting approximately 90 crop species [3]. They are major pollinators in tropical areas and their use in managed pollination, to produce high-value honey, and as recreation is increasingly popular worldwide [4].

The availability of different types of habitats affects the diversity value, abundance, and distribution patterns of stingless bees and temperature is one of the environmental parameters with the greatest impact [5]. A model study on habitat suitability for stingless bees in Colombia predicted that seven of the nine species of stingless bees used in meliponiculture would experience a significant reduction in their climatically suitable areas, and thus will likely influence agriculture and rural livelihoods [4].

In India, stingless bees are represented by three genera viz., *Lepidotrigona*, *Lisotrigona* and *Tetragonula* among which, *Tetragonula* is the most speciose with 12 species described till date Rasmussen [6] Shanas and Faseeh, [7] Viraktamath and Rojeet, [8] Viraktamath and Roy, [9].

Among the *Tetragonula* spp., *T. calophyllae* was categorised as endangered due to its rarity in the wild as well as in beekeeper managed colonies and unknown breeding behavior [7]. *T. calophyllae* currently known only from southern Kerala, is reported for the first time from Goa state, indicating a need for proper surveys to map species distribution for achieving long term conservation.

2. MATERIALS AND METHODS

Workers of stingless bee *T. calophyllae* were collected from live colonies and preserved in 98-100% ethyl alcohol. The specimens were card mounted and labeled. Card mounted samples were examined under a stereoscopic binocular microscope and morphometric parameters were recorded. Species identifications were confirmed using morphological characters, and by using published records.

3. RESULTS AND DISCUSSION

The specimen observed by us collected from Goa differed in respect of their size when compared to the Type series. The identity of *T. calophyllae*, obtained from Goa, was confirmed by the presence of a yellow band on clypeus (Fig. 1A) and presence of weakly curved median vein on the hindwing (Fig. 1B). The colouration and pilosity of the specimens were similar to the type species. However, few morphometric variations in size and character ratios observed are listed (Table. 1).

Hive entrance structure: The external hive entrance of *T. calophyllae* was found shorter, thicker and smoother, with patches of spread cerumen (Fig. 1D), probably to deter enemies, compared to the longer, thinner and drier nest entrance of *T. travancorica* Shanas and Faseeh, 2019, which had debris spread all over (Fig. 1C).

Material Examined: *T. calophyllae*: INDIA, GOA, 10 (workers): Colvale: North Goa, Arya, K. coll. 22-II-2022.

Distribution: INDIA (Kerala, Goa)

Table 1. Morphometric difference between *T. calophyllae* workers from Goa and Type series

Sl.No	Character	<i>T. calophyllae</i> Workers (Goa)	<i>T. calophyllae</i> (Type series)
1	Length of body	3.10 mm (3.06- 3.15 mm)	3.4mm (3.35-3.45 mm)
2	Head length	1.25 mm (1.22-1.26 mm)	1.55mm(1.52-1.85 mm)
3	Length of scape	0.54 mm	0.62mm(0.62-0.64 mm)
4	Length of 2 nd flagellomere	0.09 mm	0.11mm
5	WL2	0.95 mm (0.92-0.95 mm)	1.12mm(1.01-1.12 mm)
6	IOD/OD	2.50 mm	2.22-2.38 mm
7	Length /width of head ratio	0.83 mm (0.82-0.84 mm)	0.86-1.19 mm
8	Malar space /F3	0.29 mm	0.36-0.44 mm




Fig. 1. (A) Worker of *T. calophyllae*, head, frontal view (B) Worker of *T. calophyllae*, hind wing (C) Nest entrance of *T. travancorica*, (D) Nest entrance of *T. calophyllae*

Although information is limited, few studies have addressed the possible impact of climate change on the spatial distribution of stingless bees [4]. Land use changes and climate change pose a serious threat to the conservation of the stingless bees [10]. Some species can gain suitable habitats or experience relatively small habitat reductions [11,12] while others like *Melipona quadrifasciata* Lepeletier, an endemic species to the Brazilian Atlantic Forest Marchioro *et al.*, [13] and other species from the eastern Amazon of Brazil Giannini *et al.*, [12] will have higher loss of occurrence area.

There exists only few reliable records on the occurrence of stingless bee species in India Rasmussen, [6] Shanasa and Faseeh, [7] Viraktamath and Rojeet, [8] Viraktamath and Roy, [9] which is insufficient to truly map species distribution.

4. CONCLUSION

Concluding, in this study, our results obtained indicate that, the rare stingless bee species, *T. calophyllae* has an extended distribution range of about 1000 km north to the known type locality, in Kerala. Hence, a baseline distribution data on

stingless bees used in meliponiculture is essential for developing conservation policies and climate adaptation strategies under present and future climate scenarios. It is also essential that, occurrence records of different species of stingless bees have to be recorded in order to assess the threats to their conservation.

ACKNOWLEDGEMENTS

This work was supported by a postgraduate scholarship from Kerala Agricultural University for which the authors are extremely grateful.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ayala R, Gonzalez VH, Engel MS. Mexican stingless bees (Hymenoptera: Apidae): Diversity, distribution, and indigenous knowledge. In: Vit P, Pedro S, Roubik D (eds) Pot-Honey: A Legacy of Stingless Bees. Springer Science &

- Business Media, New York. 2013;135–152. Available:https://doi.org/10.1007/978-1-4614-4960-7_9
2. Engel MS, Rasmussen C, Ayala R, de Oliveira FF. Stingless bee classification and biology (Hymenoptera, Apidae): A review, with an updated key to genera and subgenera. *ZooKeys*. 2023;1172:239–312.
 3. Heard TA. The role of stingless bees in crop pollination. *Annual Review of Entomology*. 1999;44:183-206.
 4. Gonzalez VH, Cobos ME, Jaramillo J, Ospina R. Climate change will reduce the potential distribution ranges of Colombia's most valuable pollinators. *Perspectives in Ecology and Conservation*. 2021;19:195–206.
DOI: 10.1016/j.pecon.2021.02.010
 5. Trianto M, Purwanto H. Diversity, abundance, and distribution patterns of stingless bees (Hymenoptera: Meliponini) in Yogyakarta, Indonesia. *Biodiversitas Journal of Biological Diversity*. 2022;23(2):695-702.
DOI: 10.13057/biodiv/d230214
 6. Rasmussen C. Stingless Bees (Hymenoptera: Apidae: Meliponini) of the Indian subcontinent: Diversity, taxonomy and current status of knowledge. *Zootaxa*. 2013;364(3):401–428.
 7. Shanas S, Faseeh P. A new subgenus and three new species of stingless bees (Hymenoptera: Apidae: Meliponini) from India. *Entomon*. 2019;44:33–48.
 8. Viraktamath S, Rojeet T. Two new species of *Tetragonula* (Hymenoptera: Apidae: Meliponini) from North-East India with notes on their nest structure. *Biologia*. 2021;76:1691–1704.
Available:<https://doi.org/10.2478/s11756-020-00662-0>
 9. Viraktamath S, Roy J. Description of five new species of *Tetragonula* (Hymenoptera: Apidae: Meliponini) from India; *Biologia*. 2022;77(7):1769–1793.
 10. Lima VP, Marchioro CA. Brazilian stingless bees are threatened by habitat conversion and climate change. *Regional Environmental Change*. 2021;21:1-12.
Available:<https://doi.org/10.1007/s10113-021-01751-9>
 11. Giannini, TC, Maia-Silva, C, Acosta, AL, Jaffé, R, Carvalho, AT, Martins, CF, Zanella, FCV, Carvalho, CAL, Hrcir, M, Saraiva, AM, Siqueira, JO, Imperatriz-Fonseca, VL. Protecting a managed bee pollinator against climate change: strategies for an area with extreme climatic conditions and socio economic vulnerability. *Apidologie*. 2017;48:784–794.
 12. Giannini, TC, Costa, WF, Borges, RC, Miranda, L, Wanzeler da Costa, CP, Saraiva, AM, Imperatriz-Fonseca, VL. Climate change in the Eastern Amazon: crop-pollinator and occurrence-restricted bees are potentially more affected. *Regional Environmental Change* 2020;20, 9.
 13. Marchioro, CA, Lima, VP, Sales, CR, 2020. Climate change can affect the spatial association between stingless bees and *Mimosa scabrella* in the Brazilian Atlantic Forest.

© 2024 Krishna et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111420>