



Seasonal Incidence of Shoot and Fruit Borer (*Earias vittella*) in Okra

Diplal Chaudhary^{a++}, Saroj Chauhan^{b#}, Ritesh Kumar^{b†*}
and Jatin Kumar Singh^{c‡}

^a Institute of Agriculture and Natural Sciences, Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur, U.P., Pin-273009, India.

^b Department of Entomology, Institute of Agriculture and Natural Sciences, Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur, U.P., Pin-273009, India.

^c Department of Entomology, GBPUAT, Pantnagar, Uttarakhand, Pin-263145, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i84859>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120357>

Original Research Article

Received: 22/05/2024
Accepted: 24/07/2024
Published: 27/07/2024

ABSTRACT

The present investigation carried out during the *Kharif* season 2023 at Deeksha Bhawan Entomology Field, Institute of Agriculture and Natural Sciences, Deen Dayal Upadhyaya Gorakhpur University, (Uttar Pradesh). During the investigation it has been revealed that the infestation of the shoot and fruit borer, *Earias vittella* on shoots of okra commenced in the 4th week of August which gradually increased and peaked in the last week of September. The infestation of pests on fruits started in the last week of August which gradually increased and peaked in the 43rd standard

⁺⁺ M.Sc. Ag. Entomology Student;

[#] Head of Department;

[†] Assistant Professor;

[‡] Ph.D. Scholar;

*Corresponding author: E-mail: riteshskuast@gmail.com;

Cite as: Chaudhary, Diplal, Saroj Chauhan, Ritesh Kumar, and Jatin Kumar Singh. 2024. "Seasonal Incidence of Shoot and Fruit Borer (*Earias Vittella*) in Okra". *International Journal of Plant & Soil Science* 36 (8):309-15. <https://doi.org/10.9734/ijpss/2024/v36i84859>.

metrological week. The correlation studies showed that the maximum temperature, minimum temperatures, relative humidity morning, and relative humidity evening had a non-significant positive correlation while rainfall showed a negative non-significant effect on shoot infestation. However, the infestation of *E. vittella* on fruits of okra showed a negative non-significant correlation with maximum temperature, relative humidity morning and rainfall while minimum temperature and relative humidity evening had a negative significant correlation.

Keywords: Population; okra; shoot and fruit borer; correlation; weather parameters damage.

1. INTRODUCTION

Okra, *Abelmoschus esculentus* (L.) or Ladies finger, which is also known as 'Bhindi', is one of the important vegetables of India. It is one of the most widely known and utilized species of the family Malvaceae [1] and an economically important vegetable crop grown in tropical and sub-tropical parts of the world [2,3,4]. This crop is one of the most widely known and utilized species of the family Malvaceae [1]. "Okra plant was previously included in the genus *Hibiscus*. Later, it was designated to *Abelmoschus*, which is distinguished from the genus *Hibiscus*" [5]. Okra originated in Ethiopia [6] and was then propagated in North Africa, in the Mediterranean, in Arabia and India by the 12th century BC [7]. "Considering the little contact between Ethiopia and the rest of the world within historic times, it is not surprising that little is known about the early history and distribution of okra. The routes by which okra was taken from Ethiopia to North Africa, the eastern Mediterranean, Arabia, and India, and when, are by no means certain" [8]. "Okra is suitable for cultivation as a garden crop as well as on large commercial farms" [9]. "Okra plants are grown commercially in many countries such as India, Japan, Turkey, Iran, Western Africa, Yugoslavia, Bangladesh, Afghanistan, Pakistan, Myanmar, Malaysia, Thailand, India, Brazil, Ethiopia, Cyprus and in the Southern United States" [10]. Among vegetable crops, this crop occupies an important position and is grown extensively throughout India mainly for its immature fruits and an area of 5.31 lakh hectares with a production of 63.50 lakh tonnes a productivity of 11.93 tonnes per hectare. "Uttar Pradesh, Gujarat, Maharashtra, Andhra Pradesh, Tamil Nadu, Karnataka, Haryana and Punjab are the most prominent okra-growing states in India, where it is grown as a kharif and summer season crop" [11]. "Okra plays an important role in the human diet and is a good source of protein, carbohydrates, vitamins, calcium, potassium, enzymes, and total minerals which are often lacking in the diet of developing countries" [12,13]. "The composition of okra pods per 100 g

edible portion is water 88.6 g, energy 144.00 kJ (36 kcal), protein 2.10 g, carbohydrate 8.20 g, fat 0.20 g, fiber 1.70 g, Ca 84.00 mg, P 90.00 mg, Fe 1.20 mg, β -carotene 185.00 μ g, riboflavin 0.08mg, thiamin 0.04 mg, niacin 0.60 mg, ascorbic acid 47.00 mg. The fruits of okra have reawakened beneficial interest in bringing this crop into commercial production. Its medicinal value has also been reported in curing ulcers and relief from haemorrhoids. It has also been found to be medically beneficial as a plasma replacement or blood volume expander and also useful in genitourinary disorders, spermatorrhea and chronic dysentery" [14,15].

Okra is a better source of livelihood to the farmers but the infestation of different insect pests at various stages of its growth limits its successful cultivation and yield. "According to Dhamdhare et al. [16] the crop is attacked by several species of insect pests causing considerable damage. All the plant stages are susceptible to insect pests and more than seventy-two species of insect pests have been recorded on okra crop" [17]. The major pests reported are jassid, *Amrasca biguttula biguttula* (Ishida), aphid, *Aphis gossypii* (Glover), whitefly, *Bemisia tabaci* (Genn.), shoot and fruit borer, *Earias insulana* (Boisd) and *E. vittella* (Fab.), fruit borer, *Helicoverpa armigera* (Hb.), while minor pests include leaf roller, *Sylepta derogata* (Fab), mite, *Tetranychus cinabarinus* (Boisd.), green plant bug, *Nezara viridula* (Linn.) and green semilooper, *Anomis flava* (Fab.) [18,19,20].

"Among all pests, shoot and fruit borer (*Earias insulana* and *E. vittella*) the most destructive pests of okra, as young larva bores into tender shoots in early vegetative growth of plants and grown-up larvae damage fruits resulting in serious loss in yield. *E. vittella* is an oligophagous pest and its main hosts are okra and cotton" (Rajendran et al., 2018). It is also found feeding on a large number of cultivated and wild malvaceous plants species. The pest is active almost the year-round and prefers high humidity and high temperature for its growth and

development. The affected fruits are rendered unfit for human consumption, as well as for procurement of seed. The borer has been reported to cause 24.6 to 26.0 per cent damage to okra shoots [21,22], and 40 to 100 per cent loss to fruits [23,24,25]. "The pest is active almost year-round and prefers high humidity and high temperatures for its growth and development. In India, an estimated loss of 69.00 per cent in marketable yield was due to the attack of this insect on okra alone" [26].

The incidence of *E. vittella* shows violent fluctuations due to changes in climatic conditions. Information on the impact of abiotic factors on the incidence of *E. vittella* is very useful for the management of this pest in eastern Uttar Pradesh. The seasonal incidence study will help to determine the relation between the weather factors and the population of this pest, this research study also helps in okra to understanding pest dynamics, developing effective management strategies, and reducing crop losses. The study helps identify peak infestation periods, informing targeted pest management interventions and contributing to sustainable agriculture practices. Keeping in view the importance of okra and the incidence of shoot and fruit the present work was carried out to know the seasonal incidence of pest and results obtained were documented here under.

2. MATERIALS AND METHODS

The field experiment was carried out during the *Kharif* season 2023 at Deeksha Bhawan Entomology Field, Institute of Agriculture and Natural Sciences, Deen Dayal Upadhyaya Gorakhpur University, (Uttar Pradesh). Geographically Gorakhpur is situated within latitude 26°46' N and Longitude 82°2' E with an altitude of 75 meters above the mean sea level. To study the seasonal incidence of shoot and fruit borer infesting okra in relation to key abiotic factors, *Basanti* cultivar a commonly grown and recommended variety in this area was sown on the first week of August 2023 in plots of 3.0 X 3.0 m² size keeping row to row and plant to plant distance of 45 and 30 cm, respectively. For recording the observations, the crop was left for the natural infestation. The data on the shoot and fruit borer, *Earias* spp. were recorded on five randomly selected and tagged plants throughout the crop period by a visual count of the plant in which the top portion was damaged, starting after two weeks of sowing to last picking of the fruits. The per cent shoot infestation was calculated by

counting the total number of shoots and the number of damaged shoots. The per cent infestation of fruits on a number basis was calculated by counting the infested and healthy fruits separately from selected tagged plants at each picking till the last picking of fruits. The data recorded on shoot and fruit damage and meteorological parameters were used for statistical analysis [27]. The simple correlation was computed between shoot and fruit damage and abiotic factors, viz.; maximum and minimum temperature, relative humidity and rainfall. The following formulae were used for calculating the correlation coefficient is given below [28].

$$\text{Per cent shoot damage} = \frac{\text{Total number of damage shoots}}{\text{Total number of shoots}} \times 100$$

$$\text{Per cent fruit damage} = \frac{\text{Total number of damage fruit}}{\text{Total number of fruits}} \times 100$$

$$\text{Correlation coefficient} = \frac{\text{Cov}(XY)}{\sqrt{\text{var}(x)\text{var}(y)}}$$

Where,

r = correlation coefficient,
Cov(XY) = Covariance of the characters X and Y,
Var (x) and Var (y) = Variance of the characters of X and Y, respectively.

3. RESULTS AND DISCUSSION

The data on seasonal incidence of a pest provide useful information on the population buildup of pests concerning the abiotic factors of the environment. Such information can effectively be utilized in predicting the buildup of pest population and is thus helpful in revalidation of integrated pest management programmes.

Weekly observations on shoot damage were recorded as soon as infestation started, while fruit damage was recorded at each picking till the last picking of the crop. Shoot and fruit borer, *E. vittella* were recorded infesting okra shoots and fruits during the study. The shoot damage was noticed when the crop was in the vegetative stage. The data presented in Table 1 revealed that the infestation of the shoot and fruit borer on shoots of okra commenced in the 4th week of August (2.10%) i.e. four weeks after sowing (WAS) which gradually increased and reached to peak (13.32%) in the last week of September (39 SMW) at 33.35 °C and 25.41 °C maximum and minimum temperatures, 77.28 per cent relative

humidity Morning and 71.28 per cent relative humidity evening and 0.68 mm rainfall. The maximum temperature ($r = 0.194$), minimum temperatures ($r = 0.317$), relative humidity morning ($r = 0.0372$), and relative humidity evening ($r = 0.155$) had a non-significant positive correlation while rainfall ($r = -0.362$) showed a negatively non-significant effect on shoot infestation. As soon as the fruiting started, the incidence of this insect pest on the shoots started to decline and disappeared from shoots during 43 SMW. The present findings are in accordance with Meena and Kanwat [29]. Meena et al. [29], Sharma and Jat [30] who reported that the pest infestation was present during the full cropping season. Sharma and Jat [30] also reported that the shoot infestation reached to peak in the last week of September. The minimum temperature and relative humidity had a non-significant correlation with shoot damage is also in full conformity with the findings of Yadav [20]. Chaudhary [31] reported that minimum temperature, relative humidity and rainfall showed a non-significant correlation with the shoot damage. Rathore et al. [32] reported that the infestation has a non-significant correlation with maximum and minimum temperatures, relative humidity morning, relative humidity in the evening and negative non-significant with the

rainfall. Nandaniya and Chawda [33] reported a non-significant correlation between shoot infestation with minimum temperature and relative humidity evening. Ankur et al., [34] reported non-significant correlation with the minimum temperature.

The infestation of pests on fruits started in the last week of August *i.e.* 38 SMW (4.75%) which gradually increased and reached to peak (18.75 %) in the 5th week of October *i.e.* 44 SMW at 31.6 ° C maximum temperature and 17.42 ° C minimum temperature, 74.57 per cent relative humidity morning, 60.28 per cent relative humidity evening and 00.0 mm rainfall. The infestation of *Earias vitella* on fruits of okra showed a negative non-significant correlation with maximum temperature ($r = -0.481$), relative humidity morning ($r = -0.471$) and rainfall ($r = -0.496$) while minimum temperature (-0.883) and relative humidity evening (-0.663) had negative significant correlation. Yadav [20] reported that minimum temperature and relative humidity showed a significant negative correlation while rainfall had a non-significant correlation with fruit damage. Chaudhary [31] reported that minimum temperature and relative humidity showed a negatively significant correlation with fruit infestation.

Table 1. Incidence of the shoot and fruit borer in Okra during *Kharif* Season 2023

S. No.	SMW	Temperature		Relative Humidity (RH)		Rainfall (mm)	Per. shoot infestation (%)	Per. Fruit infestation (%)
		Max.temp.	Min. temp.	Morning	Evening			
1.	34	31.10	25.65	84.28	76.57	23.55	2.10	0
2.	35	35.10	26.94	71.42	62.00	0.31	3.01	0
3.	36	33.42	26.12	77.42	75.14	8.28	4.45	0
4.	37	34.75	26.88	75.28	69.28	1.18	6.75	0
5.	38	33.14	25.91	73.57	71.71	0.32	9.15	4.75
6.	39	33.35	25.41	77.28	71.28	0.68	13.32	9.15
7.	40	31.61	24.37	78.71	73.42	1.72	11.12	11.54
8.	41	34.42	23.44	71.85	55.00	0.00	9.47	12.78
9.	42	31.60	19.74	70.14	53.28	0.45	7.51	14.1
10.	43	31.90	17.65	65.00	52.71	0.00	4.12	15.75
11.	44	31.60	17.42	74.57	60.28	0.00	0	18.75

Table 2. Correlation coefficient (r) of shoot and fruit borer with weather parameters in okra crop during *Kharif*, 2023

Metrological parameters	Correlation coefficient (r)	
Merge the cell with the one above	Shoot infestation	Fruit infestation
Max. temp.	0.194	-0.481
Min. Temp.	0.317	-0.883
RH morning	0.0372	-0.478
RH evening	0.155	-0.663
Rainfall	-0.362	-0.496




Fig. 1. Shoot and fruit infestation by shoot and fruit borer in relation with the weather parameters during the study period, Kharif, 2023

There is need for better discussion. Results can be compared with the changes in other insects' correlation with weather also. This would throw some light on climate change impact and also add weight to your study and the results.

4. CONCLUSION

From the present study it can be concluded that the infestation of the shoot and fruit borer on shoots of okra commenced in the 4th week of August (2.10%) i.e. four weeks after sowing which gradually increased and reached to peak (13.32%) in the last week of September. The infestation of pests on fruits started in the last week of August i.e. 38 SMW (4.75%) which gradually increased and reached to peak (18.75%) in the 5th week of October i.e. 43 SMW. The correlation studies showed that the maximum temperature ($r = 0.194$), minimum temperatures ($r = 0.317$), relative humidity morning ($r = 0.0372$), and relative humidity evening ($r = 0.155$) had a non-significant positive correlation while rainfall (r

$= -0.362$) showed a negatively non-significant effect on shoot infestation. However, the infestation of *Earias vitella* on fruits of okra showed a negative non-significant correlation with maximum temperature ($r = -0.481$), relative humidity morning ($r = -0.471$) and rainfall ($r = -0.496$) while minimum temperature (-0.883) and relative humidity evening (-0.663) had negative significant correlation.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

I am thankful to the director professors and whole staff of IANS for providing support and materials during the period of the investigations.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Naveed A, Khan AA, Khan IA. Generation mean analysis of water stress tolerance in okra (*Abelmoschus esculentus* L.). Pakistan Journal of Botany. 2009;41:195-205.
2. Oyelade OJ, Ade-Omowaye Blo, Adeomi VF. Influence of variety on protein, fat contents and some physical characteristics of okra seeds. Journal of Food Engineering. 2003;57:111-114.
3. Andras CD, Simandi B, Orsi F, Lambrou C, Tatla DM, Panayiotou C, Domokos J, Doleschall F. Supercritical carbon dioxide extraction of okra (*Hibiscus esculentus* L.) seeds. Journal of the Science of Food and Agriculture. 2005;85:1415-1419.
4. Saifullah M, Rabbani MG. Evaluation and characterization of okra (*Abelmoschus esculentus* L. Moench.) genotypes. SAARC Journal of Agriculture. 2009;7:92-99.
5. Aladele SE, Ariyo OJ, Lapena R de. Genetic relationships among West African okra (*Abelmoschus caillei*) and Asian genotypes (*Abelmoschus esculentus*) using RAPD. Indian Journal of Biotechnology. 2008;7(10):1426-1431.
6. Sathish D, Eswar A. A Review on: *Abelmoschus esculentus* (Okra). Int. Res J Pharm. App Sci., 2013; 3(4):129-132.
7. Nzikou J, Mvoula-Tsieri M, Matouba E, Ouamba JM, Kapseu C, Parmentier M, Desobry S. A study on gumbo seed grown in Congo Brazzaville for its food and industrial applications. African Journal of Biotechnology. 2006;5(24):2469-2475.
8. Tindall HD. Vegetables in the tropics. Macmillan Education Limited, London, UK.; 1983.
9. Rubatzky VE, Yamaguchi M. World Vegetables: Principles, Production, and Nutritive Values. New York, USA: Springer New York, NY; 1997.
10. Qhureshi Z. Breeding Investigation in Bhendi (*Abelmoschus esculentus* (L.) Moench). Master Thesis, University of Agricultural Sciences, GKVK, Bangalore; 2007.
11. Anonymous. Agricultural Statistics at a Glance, Ministry of Agriculture and Farmers' Welfare, GOI. 2021;159.
12. Saifullah M, Rabbani MG. Evaluation and characterization of okra (*Abelmoschus esculentus* L. Moench.) genotypes. SAARC Journal of Agriculture. 2009;7:92-99.
13. Gopalan C, Sastri SBV, Balasubramanian S. Nutritive value of Indian foods. National Institute of Nutrition (NIN), ICMR, India; 2007.
14. Gemedede HF, Ratta N, Haki GD, Woldegiorgis AZ, Beyene F. Nutritional quality and health benefits of okra (*Abelmoschus esculentus*): A review. J Food Process Technol. 2015;6(458):2.
15. Basnet S, Lamichane P, Meheta A, Rajbanshi Y. A Review on Biochemical, Nutritional and Medicinal Properties of Okra. International Congresses of Turkish Science and Technology Publishing. 2023; 46-53.
16. Dhamdhare SV, Bhadur I, Mishra US. Studies on occurrence and succession of pests of okra at Gwalior. Indian Journal of Plant Protection. 1984;12:9-12.
17. Naik RH, Devakumar N, Rao GE, Vijaya N, Khan HI, Subha S. Performance of botanical and fungal formulation for pest management in organic okra production system. Journal of Biopesticides. 2012;5: 12-16.
18. Meena NK, Kanwat PM. Broad reconnaissance of insect pests of okra in semi-arid region of Rajasthan. National Conference on applied Entomology. 2005;261-262.
19. Mandal SK, Sah SB, Gupta SC. Management of insect pests on okra with biopesticides and chemicals. Annals of Plant Protection Sciences. 2007;15(1):87-91.
20. Yadav SS. Management of major insect pests of okra, *Abelmoschus esculentus* (L.) Moench with special reference to biorational practices. PhD Thesis submitted to S.K.N. Agriculture University, Jobner; 2015.
21. Pareek BL, Sharma GR, Bhatnagar KN. Seasonal incidence of major insect pests of okra in semi-arid region of Rajasthan. Annals of Arid Zone. 1986;25(3):222-224.
22. Zala SP, Patel JR, Patel NC. Impact of weather on magnitude of *Earias vitella* infesting okra. Indian Journal of Entomology. 1999;61(4):351-355.

23. Pareek BL, Bhargava MC. Estimation of avoidable losses in vegetable crops caused by borers under semi-arid conditions of Rajasthan. *Insect Environment*. 2003;9:59-60.
24. Shinde BD, Sarkate MB, Memade PW, Sable YR. Bioefficacy of botanical microbial and synthetic insecticides against okra fruit borer. *Pestology*. 2007;31(3):19-22.
25. Rai AB, Halder J, Kodandaram MH. Emerging insect pest problems in vegetable crops and their management in India: an appraisal. *Pest Management in Horticultural Ecosystems*. 2014;20(2):113-122.
26. Ray A, Kerketta A, Awasthi AK, Tomar RKS, Chandan M, Chandravanshi D. Impact of weather aspects on the prevalence and correlation of okra shoot and fruit borer (*Earias vittella* Fab.) during summer season. *Journal of Entomology and Zoology Studies*. 2019;7(5):773-776.
27. Panse VG, Sukhatme PV. *Statistical methods for agricultural workers*. ICAR, New Delhi. 1967.
28. Gupta SC. Correlation. In: *Fundamentals of statistics*. Himalaya Publishing House, Mumbai. 1996;510-587.
29. Meena NK, Meena BL, Kanwat PM. Seasonal occurrence of shoot and fruit borer on okra in semi-arid region of Rajasthan. *Annals of Plant Protection Sciences*. 2010;18(2):504-506.
30. Sharma R, Jat BL. Effect of abiotic factors on the incidence of shoot and fruit borer, *Earias* spp. on okra [*Abelmoschus esculentus* (Linn.) Moench]. *National Conference on Plant Protection in Agriculture*. 2010;44.
31. Chaudhary S. *Population Dynamics and Management of Shoot and Fruit Borer, Earias spp. on Okra [Abelmoschus esculentus (L.) Moench]*. (link unavailable) Thesis submitted to Sri Karan Narendra Agriculture University, Jobner; 2016.
32. Rathore BS, Baghla K, Thakur S. Effect of Abiotic Factors on Seasonal Incidence of Shoot and Fruit Borer [*Earias vittella* (Fabricius)] on Okra. *International Journal of Plant and Environment*. 2021;7(03):240-242.
33. Nandaniya MG, Patel HV, Chawda SK. Seasonal incidence and management of okra shoot and fruit borer, *Earias vittella* (Fabricius). *Pharm Innov J*. 2022;11(11): 2339-2346.
34. Ankur PR, Bhoi TK, Pensiya M, Tatarwal ML, Samal I, Majhi PK. Impact of weather parameters on okra shoot and fruit borer infestation. *The Pharma Innovation Journal*. 2022;11(4): 1736-1739.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/120357>