



# Impact of Long-term Application of Fertilizers and FYM on Nutrient Uptake by Soybean Crop and Properties of Black Soil

Chandrabhan Bajoliya <sup>a</sup>, B. S. Dwivedi <sup>a</sup>,  
Rishikesh Tiwari <sup>a\*</sup>, Prashant Kurmi <sup>a</sup>, Yagni Tekam <sup>a</sup>  
and Aruna Meravi <sup>a</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur-482004, Madhya Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i95013>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119233>

Original Research Article

Received: 08/05/2024

Accepted: 10/07/2024

Published: 18/09/2024

## ABSTRACT

Long Term Fertilizer Experiment (LTFE) play an important role in understanding the complex interactions involving plant, soil, climate, management practices and their effect on soil productivity over a long period of time. The present investigation is a part of ongoing All India Coordinated Research Project on Long Term Fertilizer Experiment with Soybean-Wheat Cropping Sequence, which was initiated since 1972. The experiment was carried out in a randomized block design

\*Corresponding author: E-mail: [rktiwari0761@gmail.com](mailto:rktiwari0761@gmail.com);

**Cite as:** Bajoliya, Chandrabhan, B. S. Dwivedi, Rishikesh Tiwari, Prashant Kurmi, Yagni Tekam, and Aruna Meravi. 2024. "Impact of Long-Term Application of Fertilizers and FYM on Nutrient Uptake by Soybean Crop and Properties of Black Soil". *International Journal of Plant & Soil Science* 36 (9):650-59. <https://doi.org/10.9734/ijpss/2024/v36i95013>.

consisting of eight treatments, viz. 50% NPK, 100% NPK, 150% NPK, 100% NP, 100% N, FYM (100% NPK+FYM), 100% NPK (-S) and Control. The significant nutrient content with nutrient uptake by soybean was noted in 100% NPK+FYM treatment over the control plot. It was also observed that nutrient content with nutrient uptake increased significantly with consecutive doses of fertilizers. On the other hand, 100% N and in the control group had the lowest value. However, adding fertilizer P in addition to N (100% NP) increased nutrient content and nutrient uptake by soybean crop. The highest. Soil organic carbon content was recorded with 100% NPK dose+FYM followed by 150% NPK treatment combination which could apparently due to the synergistic contribution of integration of fertilizer and FYM application. Similarly, the available N,P, K and S content in soil were found highest with 100% NPK + FYM followed by 150% NPK fertilizer dose. However, the lowest content was noted in control where crop was grown without fertilizers. Available sulphur was significantly higher over control in all the treatments receiving single super phosphate.

**Keywords:** Soil properties; nutrient content; nutrient uptake; soybean.

## 1. INTRODUCTION

Soybean (*Glycine max* L.) is one of the leading oil and protein-containing crops in the world. "Long-term fertilizer experiments give valuable information on the effect of continuous application of different levels of fertilizer nutrients alone with and without organic manure under intensive cropping on soil fertility and crop productivity. These experiments can be used for precise monitoring of changes in soil fertility and could be of paramount help in solving the complex problems related to soil fertility management" [1,2]. It has been discovered that a long-term organic amendment significantly affects soil parameters including pH, accessible phosphorus, inorganic nitrogen, organic carbon, and other organic matter [3]. Crop production requires balanced fertilization; however, applying manure in combination with other fertilizers may minimize the requirement for artificial fertilizers. According to studies by [4,5,6], applying chemical fertilizers in addition to FYM may increase the yield and yield contributing characteristics like grain yield, straw yield, nutrient content, and nutrient uptake. "The soil available N, P, K and S were also found higher under the sole application of balanced fertilization and its combined application with FYM. The long-term fertilizer application with or without organic manure influences the soil nutrient status" [7]. "observed that modern soybean nutrient uptake compared to cultivars planted in the country". "The maximum N content in grain and straw was observed with the application of super optimal dose and the lowest content in grain and straw was recorded in the control treatment" [8,9]. In view of this, the present investigation was undertaken to study the Impact of long-term application of fertilizer

and FYM on nutrient uptake by soybean crops and properties of a black soil.

## 2. MATERIALS AND METHODS

### 2.1 Location Description

The present investigation is a part of an ongoing All India Coordinated Research Project on Long-Term Fertilizer Experiment with the soybean-wheat cropping system, which was initiated in 1972 at the Research Farm of Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh, India. The experimental site (23°10" N latitude and 79°57" E longitude) has a semi-arid and subtropical climate with a characteristic feature of dry summer and cold winter. During winter, i.e., from November to February, the temperature ranges from 4 to 33°C and the relative humidity varies from 70 to 90%. Dry and warm weather usually prevails during March to June. The temperature in the month of May attains a value as high as 46°C. The monsoon season extends from mid-June to mid-September. The temperature during this period varies from 25 to 35°C, and the relative humidity ranges from 70 to 80%. The total annual rainfall varies from 1000 to 1500 mm with a mean value of around 1350 mm. The length of the growing period ranges from 150 to 180 days.

### 2.2 Experimental Details

The experiment was designed and conducted with 8 treatments having four replications arranged in the randomized block design. Two blocks were separated with a gap of 2.0 m, whereas individual plots (17.0 m x 10.8 m) were separated with a permanent bund. The 100% optimal NPK doses based on initial (1972) soil

test values were 20:80:20 and 120:80:40 kg ha<sup>-1</sup> for soybean and wheat, respectively. The treatment details and quantity of nutrients added are given in Table 1.

“The sources of N, P and K used were urea, single superphosphate and muriate of potash. In sulphur-free treatment, diammonium phosphate (DAP) was used instead of SSP as a source of phosphorus. Farmyard manure @ 5 t ha<sup>-1</sup> year<sup>-1</sup> was applied only to soybean crop during *kharif* season. During the rainy season, all the nutrients, viz. N, P and K were applied as basal before last harrowing, whereas in wheat half of the nitrogen was applied at the time of sowing and the remaining half was applied in two splits first half at 21–25 days (after first irrigation) and the rest at 51-55 days after sowing” [8].

### 2.3 Soil Sampling and Analysis

Initial soil samples (0-15 cm depth) were taken from 8 different locations in the study area and analyzed for pH, EC, organic carbon, available

N, P and K by following standard procedures. “Soil pH and EC were determined by preparing 1:2.5 soil water suspension and measured by glass electrode pH meter and digital electrical conductivity meter [10]. Organic carbon content in soil was determined using the potassium dichromate rapid titration method [11] and soil available N was determined by [12]. Available phosphorus was extracted with 0.5 M NaHCO<sub>3</sub> (pH 8.5) and colour was developed by ascorbic acid [13] and available potassium was extracted with neutral normal ammonium acetate and estimated using a flame photometer [10]. Available sulphur was determined by [14]. All soybean plants were harvested at crop maturity and grain yield was obtained. Formula: Nutrient absorption (kg ha<sup>-1</sup>) = Nutrient content (%) x Yield (kg ha<sup>-1</sup>)” [8].

### 2.4 Statistical Analysis

The data generated on soil analysis, grain and straw yields and their P uptake were statistically analyzed to draw suitable inference as per standard method [15].

**Table 1. Details of treatment and nutrient rates (kg ha<sup>-1</sup>) in soybean and wheat**

Treatments	Soybean			Wheat		
	N	P	K	N	P	K
50 % NPK	10	17.6	8.3	60	17.6	16.6
100 % NPK	20	35.2	16.6	120	35.2	33.2
150% NPK	30	52.8	24.9	180	52.8	49.8
100 %NP	20	35.2	NA	120	35.2	NA
100 % N	20	NA	NA	120	NA	NA
100 % NPK+FYM*	46.5	41.8	42.7	120	35.2	33.2
100 % NPK (-S)#	20	35.2	16.6	120	35.2	33.2
Control	NA	NA	NA	NA	NA	NA

\*FYM (contains 0.53% N, 0.30% P<sub>2</sub>O<sub>5</sub> & 0.63% K<sub>2</sub>O) was applied 5 t ha<sup>-1</sup> to soybean every year 15–20 days before sowing;  
# DAP was used instead of SSP as source of P.

**Table 2. Initial properties of the experimental soil (1972)**

S.NO.	Soil properties	value
1.	pH (1:2.5)	7.6
2.	Electrical conductivity (1:2.5)	0.18 dSm <sup>-1</sup>
3.	Organic. carbon	5.7 g kg <sup>-1</sup>
4.	Calcium carbonate	46.0 g kg <sup>-1</sup>
5.	Available nitrogen	193 kg ha <sup>-1</sup>
6.	Available phosphorus	7.60 kg ha <sup>-1</sup>
7.	Available potassium	370 kg ha <sup>-1</sup>
8.	Available sulphur	15.6 kg ha <sup>-1</sup>

### 3. RESULTS AND DISCUSSION

#### 3.1 Impact of Treatments on Properties of a Black Soil

**Soil pH:** “The data presented in Table 3, ranged between 7.46 (100% N) to 7.60 (150% NPK) in various treatments. There was not any significant effect of continuous use of fertilizer and manure on soil pH. *This could be due to the high buffering capacity of the soil and the presence of appreciable content of free calcium carbonate*” [1].

**Soil Electrical conductivity:** “The EC values in soil as influenced by the long term application of fertilizer and FYM are presented in Table 3. It ranged between 0.14 (control) to 0.19 dSm<sup>-1</sup> (150% NPK) in various treatments and it was noticed that the continuous application of different doses of fertilizers and manure did not resulted in any significant effect on soil EC” [16]. This could also be due to the inherent high buffering capacity of soil [5].

**Organic carbon in soil:** The organic carbon significantly increased with an increase in the doses of fertilizers. The lowest value was noticed in control (4.23 g kg<sup>-1</sup>) which was increased to 5.75, 7.48 and 8.34 g kg<sup>-1</sup> due to the use of 50% NPK, 100% NPK and 150% NPK of RDF, respectively. The highest content 8.96 g kg<sup>-1</sup> was observed in 100% NPK+FYM treatment (Table 3). Further, 100% NPK+FYM dose showed a significantly higher content of organic carbon over optimal and sub-optimal doses. “This could be ascribed to the organic manure (5 t FYM ha<sup>-1</sup>) application combination with fertilizers that increased total N and soil organic matter contents compared with sole fertilizer treatments. Increasing levels of fertilizer application helped in increasing the organic carbon content, which may be ascribed to an increase in productivity and incorporation of larger residual biomass through root, leaves, stable, and rhizodeposition” [17,18]. “Further, it may also be attributed to the addition of FYM which stimulated the growth and activity of microorganisms and better root biomass” [19,20]. Soil organic matter is vital for enhancing soil fertility and sustaining crop productivity.

**Content of available N in soil:** The data on Table 3 indicated that the highest available N content (315 kg ha<sup>-1</sup>) was recorded in 100% NPK+FYM could be a result due to better

biological activities in the presence of FYM [21,22]. Whereas the lowest value was noticed as 183 kg ha<sup>-1</sup> in control. However, due to the addition of fertilizer dose (suboptimal, optimal and super optimal), N content was correspondingly improved indicating an impact of fertilizer application on the enrichment of available nitrogen in soil. Available nitrogen in the soil was significantly higher in all the treatments over control except 100%N and 50%NPK treatments [23].

**Content of available P in soil:** The result showed that the available P content increased successively and significantly from 9.38 kg ha<sup>-1</sup> (control) to 21.60, 32.86 and 38.55 kg ha<sup>-1</sup> in 50%NPK, 100%NPK and 150%NPK treatments, respectively (Table 3).

The content of available P was higher in 100% NPK (32.86 kg ha<sup>-1</sup>) than 100% NPK-S (30.93 kg ha<sup>-1</sup>). Because continuous use of balanced fertilizer is conducive for maintaining the soil available P. In black soil, the applied phosphorus gets fixed (80-85%) and only a small part (15–20%) of it becomes available to the plants [24,22]. However, the build-up of available P in 100% NPK+ FYM was higher 39.42 kg ha<sup>-1</sup> in comparison to 32.86 kg ha<sup>-1</sup> in 100% NPK treatments indicating the beneficial effect of FYM on mineralization of P to a greater extent in soil. [25,26]. “Further, the imbalanced use of fertilizers reduced the available P content in the soil. A significant reduction in available P content was observed under nitrogen alone (100% N) and unfertilized treatments due to removal of P by the crops in the absence of P supplementation through an external source. Use of 100% NP over 100% N significantly increased the available P status of soil. A similar trend on available P was also reported” [27].

**Content of available K in soil:** The perusal of data indicated a declining trend (228 to 328 kg ha<sup>-1</sup>) from its initial level (370 kg ha<sup>-1</sup>) of available K status which indicates considerable soil mining of available K after forty five years of intensive cropping (Table 3). The decline was observed maximum in case of control followed by 100% N alone. The magnitude of decline decreased with increasing levels of NPK applications. Among the inorganic fertilizers, continuous application of N or NP had a depressive effect on available K content of the soil, which may be due to nutrient imbalance in soil. Continuous omission of K in crop nutrition caused mining of its native pools caused reduction in the crop yields” [18]. The

available K content was significantly higher in 100%NPK-S, 100%NPK, 150%NPK and 100%NPK+FYM treatments. The highest value of available K (328 kg ha<sup>-1</sup>) was recorded in 100% NPK+FYM followed by 150% NPK (305 kg ha<sup>-1</sup>) whereas; the lowest value (228 kg ha<sup>-1</sup>) was observed in control. Available K content in soil was significantly higher in 150%NPK and 100%NPK+FYM over 50%NPK treatments. Similar results have also been reported [28,29].

**Content of available sulphur in soil:** The maximum build up of available sulphur was observed in 100% NPK+FYM (37.03 kg ha<sup>-1</sup>) followed by 150% NPK (35.20 kg ha<sup>-1</sup>). The magnitude of buildup corresponds to the proportion to level of applied fertilizer [30]. The highest value was found associated with FYM addition that provide the favourable condition for better nutrient use efficiency [31]. However, the minimum content of available sulphur was observed in control (10.90 kg ha<sup>-1</sup>). The content of available sulphur was found higher in 100% NPK (31.05 kg ha<sup>-1</sup>) as compared to 100% NPK-S (11.10 kg ha<sup>-1</sup>). It was also found that the available sulphur content was higher in 50% NPK (22.10 kg ha<sup>-1</sup>) than 100% N (11.20 kg ha<sup>-1</sup>). Available sulphur was significantly higher over control in all the treatments receiving single super phosphate, respectively [30,16].

### 3.2 Nutrient Content in Soybean Grain and Stover

**Nutrient content in grain:** The data of Table 4 indicate that the highest N content (6.69%) was recorded in 100% NPK+FYM whereas the lowest value was noticed as 3.80% in control. Nitrogen content in grain was significantly higher in all the treatments over control except 50%NPK treatments. Whereas, P content in grain raised successively from 0.23% (control) to 0.25%, 0.33% and 0.35% in 50%NPK, 100%NPK and 150%NPK treatments respectively. Phosphorus content in grain was higher in 100% NPK+FYM (0.36%) than 150% NPK (0.35%). P content in grain was significantly higher in all the treatments over control except 50%NPK and 100%N treatments. The highest K content (1.68%) was recorded in 100% NPK+FYM whereas the lowest value was noticed as 1.18% in control. Potassium content in grain was significantly higher in all the treatments over control except 50%NPK, 100%NP and 100%N treatments. The S content in grain increased successively from 0.18% (control) to 0.22%, 0.30% and 0.32% in 50%NPK, 100%NPK and 150%NPK treatments,

respectively. Sulphur content of grain was higher in 100% NPK+FYM (0.33%) than 150% NPK (0.32%). S content in grain was significantly higher in all the treatments over control except 50%NPK, 100%N and 100%NPK-S treatments also reported [6,32].

**Nutrient content in stover:** The data of Table 4 indicate that the highest N content (2.81%) in stover resulted by 100% NPK+FYM whereas the lowest value was noticed as 1.51% in control. Nitrogen content in stover was significantly higher in all the treatments over control except 50%NPK treatment. As regards the P content in stover raised successively from 0.11% (control) to 0.12%, 0.15% and 0.16% in 50%NPK, 100%NPK and 150%NPK treatments, respectively. Phosphorus content in stover was higher in 100% NPK+FYM (0.17%) than 150% NPK (0.16%). P content in stover was significantly higher in all the treatments over control except 50%NPK and 100%N treatments. Further, the highest K content (2.56%) was observed in 100% NPK+FYM whereas the lowest value was obtained (1.89%) in control. Potassium content in stover was significantly higher in all the treatments over control except 50%NPK, 100%N and 100%NP treatments. The S content in stover raised successively from 0.14% (control) to 0.16%, 0.21% and 0.22% in 50%NPK, 100%NPK and 150%NPK treatments, respectively. Sulphur content of stover was higher in 100% NPK+FYM (0.23%) than 150% NPK (0.22%). Sulphur content in stover was significantly higher in all the treatments over control except 50%NPK, 100%N and 100%NPK-S treatments [1,8].

### 3.3 Nutrient Uptake by Soybean Grain and Stover

**Nutrient uptake by grain:** The effect of fertilizer and FYM addition on nutrient uptake by crop revealed that the wide variation has been obtained which clearly reflect the influence of varying fertilizer levels on crop growth and yield. In this regard, uptake of N, P, K and S by grain (Table 5.) synchronized with the biomass yield produced and the highest uptake was found in 100 % NPK + FYM treatment followed by 150 % NPK and 100 % NPK. The beneficial effect of FYM can be due to steady supply of all nutrients including the micronutrients and improvements in physical condition. The similar beneficial effects

FYM can be due to steady supply of all nutrients including the micronutrients and improvements in

physical condition. They observed that the continuous use of chemical fertilizers either singly or in combination with FYM had a marked effect on nutrient uptake by soybean [33,34,4]. The uptake of N, P, K and S by soybean grain were found 48.88, 2.23, 11.34 and 1.75 kg ha<sup>-1</sup>, respectively in 100 % N alone treatment, increased to 86.89, 4.55, 20.53 and 4.43 kg ha<sup>-1</sup> due to inclusion of P i.e. 100 % NP. However, inclusion of K with 100 % NP i.e. 100 % NPK resulted in slight increase as 104.98, 6.32, 26.93 and 5.75 kg ha<sup>-1</sup>, and minimum uptake of N, P, K and S in grain was observed in control plot, respectively. Total nutrient uptake in soybean was significantly higher in all the treatments over control except 100%N treatments [35,36].

**Nutrient uptake by stover:** The finding exhibited that successively increasing addition of fertilizer proportionately increased the uptake of

nutrients by the crop. In this regard, uptake of N, P, K and S by stover (Table 5) synchronized with the biomass yield produced and the highest uptake was found in 100% NPK + FYM treatment followed by 150% NPK and 100 % NPK. [16]. who the maximum was also fund with the treatment of integrated use of organic manure and recommended fertilizer dose as compared to control plot [37,16]. The uptake of N, P, K and S in soybean stover which was 39.75, 3.36, 40.12 and 3.13 kg ha<sup>-1</sup>, respectively in 100 % N alone treatment, increased to 56.23, 4.63, 49.98 and 4.67 kg ha<sup>-1</sup> due to inclusion of P i.e. 100 % NP. However, inclusion of K with 100 % NP i.e. 100 % NPK resulted in slight increase 74.38, 5.79, 70.82 and 6.34 kg ha<sup>-1</sup>, and minimum uptake of N, P, K and S in straw were found in the control plot, respectively. Total nutrient uptake in soybean was significantly higher in all the treatments over control except 100% N treatment [35,8].

**Table 3. Effect of long-term application of fertilizer and FYM on Soil properties**

Treatments	pH	EC (dsm <sup>-1</sup> )	OC (g kg <sup>-1</sup> )	Available Nutrients (kg ha <sup>-1</sup> )			
				N	P	K	S
50 % NPK	7.50	0.14	5.75	201	21.60	254	22.10
100 % NPK	7.55	0.17	7.48	278	32.86	280	31.05
150% NPK	7.60	0.19	8.34	302	38.55	305	35.20
100 % NP	7.54	0.17	6.68	248	31.39	230	29.30
100 % N	7.46	0.15	5.68	219	9.88	227	11.20
100%NPK+FYM	7.54	0.18	8.96	315	39.42	328	37.03
100 % NPK-S	7.56	0.17	7.25	263	30.93	270	11.10
Control	7.55	0.14	4.23	183	9.38	228	10.90
<b>CD (p=0.05)</b>	<b>NS</b>	<b>NS</b>	<b>1.05</b>	<b>40.10</b>	<b>3.97</b>	<b>39.24</b>	<b>4.00</b>

**Table 4. Effect of treatments on nutrient content of soybean grain and stover**

Treatments	Nutrients (%)							
	Nitrogen		Phosphorus		Potassium		Sulphur	
	Grain	Stover	Grain	Stover	Grain	Stover	Grain	Stover
50 % NPK	4.36	1.73	0.25	0.12	1.28	2.12	0.22	0.16
100 % NPK	5.51	2.48	0.33	0.15	1.42	2.37	0.30	0.21
150% NPK	6.38	2.65	0.35	0.16	1.64	2.48	0.32	0.22
100 % NP	5.41	2.17	0.28	0.14	1.26	1.92	0.27	0.18
100 % N	5.18	1.90	0.24	0.12	1.20	1.90	0.19	0.15
100%NPK+FYM	6.69	2.81	0.36	0.17	1.68	2.56	0.33	0.23
100 % NPK-S	5.40	2.20	0.29	0.15	1.39	2.30	0.19	0.15
Control	3.80	1.51	0.23	0.11	1.18	1.89	0.18	0.14
<b>S Em±</b>	<b>0.217</b>	<b>0.089</b>	<b>0.011</b>	<b>0.005</b>	<b>0.049</b>	<b>0.120</b>	<b>0.015</b>	<b>0.008</b>
<b>CD (p=0.05)</b>	<b>0.629</b>	<b>0.259</b>	<b>0.032</b>	<b>0.013</b>	<b>0.141</b>	<b>0.349</b>	<b>0.044</b>	<b>0.024</b>

**Table 5. Effect of treatment on nutrient uptake (kg ha<sup>-1</sup>) by soybean grain and stover**

Treatments	Nitrogen			Phosphorus			Potassium			Sulphur		
	Grain	Stover	Total	Grain	Stover	Total	Grain	Stover	Total	Grain	Stover	Total
50% NPK	64.30	43.42	107.72	3.74	3.74	7.48	18.97	53.11	72.07	3.23	3.96	7.20
100%NPK	104.98	74.38	179.36	6.32	5.79	12.11	26.93	70.82	97.75	5.75	6.34	12.08
150%NPK	139.70	92.57	232.27	7.75	7.30	15.04	35.84	86.90	122.74	6.95	7.80	14.76
100%NP	86.89	56.23	143.12	4.55	4.63	9.18	20.53	49.98	70.51	4.43	4.67	9.09
100%N	48.88	39.75	88.64	2.23	3.36	5.60	11.34	40.12	51.46	1.75	3.13	4.88
100%NPK+FYM	152.86	101.31	254.17	8.28	7.84	16.11	38.50	92.96	131.45	7.51	8.41	15.92
100%NPK-S	87.23	66.20	153.43	4.64	5.32	9.96	22.49	68.94	91.43	2.97	4.44	7.41
Control	34.98	30.17	65.15	2.08	2.75	4.83	10.94	38.26	49.19	1.63	2.82	4.45
<b>S Em±</b>	<b>5.253</b>	<b>3.167</b>	<b>7.528</b>	<b>0.317</b>	<b>0.483</b>	<b>0.642</b>	<b>1.598</b>	<b>4.134</b>	<b>4.935</b>	<b>0.246</b>	<b>0.309</b>	<b>0.440</b>
<b>CD (p=0.05)</b>	<b>15.244</b>	<b>9.189</b>	<b>21.847</b>	<b>0.921</b>	<b>1.403</b>	<b>1.864</b>	<b>4.639</b>	<b>11.998</b>	<b>14.322</b>	<b>0.727</b>	<b>0.898</b>	<b>1.278</b>

#### 4. CONCLUSIONS

Thus, it could be concluded that the long-term application of balanced and integrated use of nutrient (100% NPK, 150% NPK and 100% NPK+FYM) to soybean significantly improved the soil properties, nutrient content, and nutrients uptake. It has been noted that as fertilizer doses were added one after the other, the proportion of nutrients in the soil and plants grew. Consequently, the control and 100% N alone had the lowest concentration, and a growing trend was seen with higher fertilizer addition, ranging from suboptimal to optimal and super optimal doses. Generally, grain was shown to have a higher composition and absorption of nutrients than straw. Long-term prospects are thus shown to be good when organic and mineral fertilizers are used together and wisely.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Nagwanshi A, Dwivedi AK, Dwivedi BS, Anupama K. Effect of long term application of fertilizer and manure on soil properties of a Vertisol in soybean-wheat cropping sequence. *International Journal of Chemical Studies*. 2018;6(3):2946-2949.
2. Tiwari R, Dwivedi BS, Sharma YM, Thakur R, Sharma A, Nagwanshi A. 2024. Soil Properties and Soybean Yield as Influenced by Long Term Fertilizer and Organic Manure Application in a Vertisol under Soybean-Wheat Cropping Sequence. *Legume Research*; 2024;47(7): 1158-1164.
3. Ge S, Zhu Z, Jiang Y. Long-term impact of fertilization on soil pH and fertility in an apple production system. *J Soil Sci Plant Nutr*. 2018;18:282–293.
4. Dwivedi BS, Pandey AK, Tiwari RK, Jha AK, Khamparia NK. Performance of integrated nutrient management on yield and uptake direct seeded rice. *Progressive Agriculture*. 2012;12(2):381-385.
5. Dwivedi AK, Dwivedi BS. Impact of long term fertilizer management for sustainable soil health and crop productivity: Issues and challenges. *JNKVV Res. J*. 2015; 49(3):387-397.
6. Raghuwanshi C, Khamparia NK, Thakur RK, Sahare SK, Sharma Vaishali, and Dwivedi BS. Impact of long-term application of fertilizers and organic manure on yield attributes characters, yield of soybean and soil fertility in a Vertisol. *Eco. Env.& Cons*. 2016;22(2):745-753.
7. Khandagle A, Dwivedi BS, Aher SB, Dwivedi AK, Yashona DS, Jat D. Effect of long term application of fertilizers and manure on soil properties. *Journal of Soils and Crops*. 2019;29(1):97-104.
8. Tiwari R, Sharma YM, Dwivedi BS, Mitra NG, Kewat ML. Nutrient content and uptake by soybean as influenced by continuous application of fertilizer and manure in black soil, *Journal of Pharmacognosy and Phytochemistry*. 2019;8(4):140-144.
9. Dwivedi BS, Dwivedi AK, Sharma A, Nagwanshi A. Effect of continuous fertilizer and manure application on productivity of soybean-wheat cropping sequence. *Abst. AgriCon 2018 International Conference on "Sustainability of Smallholder Agriculture in Developing Countries under Changing Climatic Scenario" at C.S. Azad University of Agriculture & Technology, Kanpur during. 14-17 Feb. 2018*;156.
10. Jackson ML. *Soil chemical analysis*. Prentice Hall of India Pvt. Ltd., New Delhi; 1973.
11. Walkley A, Black IA. An examination of the Digtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*. 1934;37:29-33.
12. Subbiah BV, Asija GL. A rapid procedure for the estimation of available nitrogen in soils. *Current Science*. 1956; 25: 259-260.
13. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available P in soil by extraction with NaHCO<sub>3</sub>. *USDA, Cir*. 1954;939.
14. Chesnin L, Yien CH. Turbid metric determination of available sulphate. *Soil Science Society of America Proceedings*. 1950;15:149-151.
15. Panse VG, Sukhatme SV. *Statistical Methods for Agricultural Workers*. ICAR Publication; 1970.



16. Suman J, Dwivedi BS, Dwivedi AK, Pandey SK. Impact of continuous addition of fertilizers and manure on soil nutrient status, uptake, yield, protein and oil content of soybean in a Vertisol. *Research Journal of Agricultural Sciences*. 2017;8:159-163.
17. Parihar SK, Dwivedi BS, Khan IM, Tiwari RK. Effect of integrated nutrient management on yield and economics of little millet (*Panicum sumatrense* roth. Ex Roem. and Schult.). *Journal of Soils and Crops*. 2010;20(2):211-215.
18. Pathariya Priyanka, Dwivedi BS, Dwivedi AK, Thakur RK, Singh Muneshwar, Sarvade S. Potassium Balance under Soybean–wheat Cropping System in a 44 Year Old Long Term Fertilizer Experiment on a Vertisol. *Communications in Soil Science and Plant Analysis*. 2022; 53(2):214-226.
19. Dwivedi BS, Rawat AK. Nutrient management technology for niger (*Guizotia abssinica* L. F.) crop in tribal areas. *Plant Archives*. 2013;13(2):809-813.
20. Meshram MK, Dwivedi BS, Naik KR, Thakur RK, Keram KS. Impact of organic and inorganic sources of nutrients on yield, nutrient uptake, soil fertility and economic performance of rice in a Typic Haplustert. *Journal of Soils and Crops*. 2018;28(1):31-36.
21. Suman J, Dwivedi BS, Dwivedi AK, Pandey SK. Interaction effect of phosphorus and Sulphur on yield and quality of Soybean in Vertisol. *International Journal of Current Microbial Applied Sciences*. 2018;7(3):152-158.
22. Dwivedi BS, Sharma Abhishek, Dwivedi AK, Thakur RK. Response of phosphorus application on productivity of wheat at farmer field. *Universal Journal of Agricultural Research*. 2019;7 (1):20-24.
23. Bairwa J, Dwivedi BS, Rawat A, Thakur RK, Mahawar N. Long-Term Effect of Nutrient Management on Soil Microbial Properties and Nitrogen Fixation in a Vertisol under Soybean– Wheat Cropping Sequence. *Journal of the Indian Society of Soil Science*. 2021;69(2):171-178.
24. Dubey L, Dwivedi BS, Dwivedi AK, Thakur RK. Effect of long term application of fertilizers and manure on profile distribution of various phosphorus fractions in Vertisol. *Green Farming*. 2016;7(2):365-370.
25. Mishra AK, Ghosh PK, Hati KM. Effect of integrated use of farmyard manure and chemical fertilizers on soil physical properties and productivity of soybean. *Journal of the Indian Society of Soil Science*. 2010;38(2):416-418.
26. Bagde V, Dwivedi BS, Dwivedi AK, Thakur R, Nagwanshi A. Effect of Long-Term Fertilizer and Farm yard Manure Application on Soil Phosphorus Fractions in a Vertisol under Soybean–Wheat Cropping Sequence. *Journal of the Indian Society of Soil Science*. 2023;71(2):207-216.
27. Bairwa J, Dwivedi BS, Dwivedi AK, Thakur RK, Patidar N. Impact of long term fertilization on soil properties and soybean productivity in a Vertisol. *J. Curr. Microbiol. App. Sci*. 2020;9(9):527-536.
28. Patidar N, Dwivedi AK, Dwivedi BS, Thakur RK, Bairwa J, Sharma A. Impact of Long-Term Application of Inorganic Fertilizers and Organic Manure on Soil Fertility and Crop Productivity under Soybean-Wheat Cropping System in a Vertisol. *International Journal of Environment and Climate Change*. 2021; 11(8):24-30.
29. Inwati D, Kumar, Dixit BK, Thakur R, Dwivedi BS, Sharma A, Singh V. Effect of long term fertilizer and FYM application on soil fertility of a vertisol under soybean-wheat cropping system. *Journal of Experimental Agriculture International*. 2022;44(10):179-184.
30. Birla K, Khamparia NK, Thakur RK, Dwivedi, BS and Sawarkar SD. Profile Distribution of Various Fractions of Sulphur as Influenced by Long Term Application of Fertilizers and Manure in Vertisol. *Green Farming*. 2015;6(4):753-756.
31. Singh M, Wanjari RH, Dwivedi A, Ramdalal. Yield response to applied nutrient and estimates of N<sub>2</sub> fixation in 33-year-old soybean-wheat experiment on a vertisol. *Expl Agric*. 2012;48(3):311-325.
32. Solanki AC, Solanki MK, Nagwanshi A, Dwivedi AK, Dwivedi BS. Nutrient uptake and grain yield enhancement of soybean by integrated application of farmyard manure and NPK. *International Journal of Current Microbial Applied Sciences*. 2018;7(9):1093-1102.
33. Sharma GD, Dwivedi BS, Nema HR. Effect of graded level of integrated input resources on yield and nutrients concentration in soybean. *JNKVV Res. J*. 2006;40 (1&2): 9-14.

34. Tiwari RK, Dwivedi BS, Singh Nirmala, Pandey AK. Organic farming in India. JNKVV Res. J. 2012;46(2): 141-159.
35. Gupta A, Dwivedi AK, Nagwanshi A, Dwivedi BS, Vishwakarma AK. Impact of longterm application of inorganic fertilizer and farm yard manure on productivity of soybean in Vertisol. Bulletin of Environment Pharmacology and Life Sciences. 2019; 8(4):116-132.
36. Khandagle A, Dwivedi BS, Dwivedi AK, Panwar S, Thakur RK. Nitrogen fractions under long-term fertilizer and manure applications in soybean – Wheat rotation in a Vertisol. Journal of the Indian Society of Soil Science. 2020;68(2):186-193.
37. Dwivedi BS, Rawat AK, Dixit BK, Thakur RK. Effect of inputs integration on yield, uptake and economics of kado millet (*Paspalum scrobiculatum* L.). Economic Affirs. 2016: 61(3): 519-526.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/119233>