



# Evaluation of the Antioxidant Power and Secondary Metabolites Contents of the Crude Compounds of *Talinum triangulare* (Talinaceae)

Koffi A. Florence <sup>a\*</sup>, Koffi A. Jeanne <sup>b</sup>, Tano K. Dominique <sup>b</sup>,  
Konan Ange <sup>c</sup>, Okou O. Constantin <sup>a</sup> and Yavo William <sup>b</sup>

<sup>a</sup> Group of Excellence and Research on Traditional Pharmacopoeia Products (GeRProPhaT), Agrovalorization Laboratory, Agroforestry Training and Research Unit (UFR), Jean Lorougnon Guédé University (UJLoG), Côte d'Ivoire.

<sup>b</sup> Center for Research and Combating Malaria (CRLP) at the National Institute of Public Health (INSP) in Abidjan, Côte d'Ivoire.

<sup>c</sup> Laboratory of Biology and Health of the UFR Biosciences of the Félix Houphouët Boigny University of Abidjan (UFHB), Côte d'Ivoire.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijbcrr/2024/v33i6917>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124698>

Original Research Article

Received: 20/08/2024  
Accepted: 23/10/2024  
Published: 31/10/2024

## ABSTRACT

*Talinum triangulare* is a food plant recognized for its nutritional properties and therapeutic virtues. This plant is used in particular in Daloa (Central-West Ivory Coast) by naturotherapists, for the treatment of malaria. This study aimed to evaluate the antioxidant activity of extracts and fractions

\*Corresponding author: E-mail: [fleurlaureat@gmail.com](mailto:fleurlaureat@gmail.com);

**Cite as:** Florence, Koffi A., Koffi A. Jeanne, Tano K. Dominique, Konan Ange, Okou O. Constantin, and Yavo William. 2024. "Evaluation of the Antioxidant Power and Secondary Metabolites Contents of the Crude Compounds of *Talinum Triangulare* (Talinaceae)". *International Journal of Biochemistry Research & Review* 33 (6):340-49. <https://doi.org/10.9734/ijbcrr/2024/v33i6917>.

of *Talinum triangulare* leaves and to determine their content of secondary metabolites. Two methods *in vitro* were used for this evaluation: reduction of the 2,2'-diphenyl-1-picrylhydrazyl radical (DPPH) and reduction of the ABTS<sup>+</sup> cation radical (2,2'-azinobis-3-ethylbenzothiazoline-6-sulfonic acid).

The results show that the IC<sub>50</sub> could not be determined for the 70 % aqueous and hydro-ethanolic extracts with the DPPH test, while it was respectively 0.9 and 2.3 mg/mL for the fractions dichloromethane and aqueous. Regarding the ABTS<sup>+</sup> radical-cation reduction test, the dichloromethane fraction showed a higher percentage of inhibition (41.59 %) compared to the aqueous fraction (40.59 %), while the aqueous fraction was a little more important than the hydro-ethanolic extract at 70 % (38.29 %). Among the four compounds tested, only the aqueous extract presented a relatively low percentage of inhibition (25.51 %).

Spectrophotometric analysis revealed the presence of various phenolic compounds, such as flavonoids, polyphenols and total tannins, at varying concentrations. These results could support the traditional use of *Talinum triangulare* in the treatment of malaria.

**Keywords:** *Talinum triangulare*; antioxidant activity; phenolic compounds.

## 1. INTRODUCTION

Over the past twenty years, the incidence of parasitic diseases has increased. Malaria is among these parasitic diseases. Despite being entirely preventable and treatable, it continues to have devastating consequences on the health and livelihoods of populations across the world [1]. This state of affairs has led authors to affirm that this disease is “one of the rare public health scourges that has survived the centuries without ever losing its activity” [2,3]. Unfortunately, its management faces resistance from most antimalarial drugs [4]. It is therefore crucial to search for new active molecules and medicinal plants appear to be an important source to explore. Furthermore, natural therapies based on medicinal plants are of great benefit. They are better biodegradable and are suspected of having resistance-reversing properties. In addition, the best treatments recommended by the WHO for the therapeutic management of malaria are based on derivatives of molecules from medicinal plants [5,6].

According to the World Health Organization (WHO), more than 80 % of the African population uses traditional medicine for their primary health care due to their proximity and accessibility [7]. Ivory Coast is one of the third world countries where medicinal plants are commonly used to solve health problems, especially in rural areas. This is how a significant number of works relating to the biological, pharmacological and phytochemical properties of plants from traditional Ivorian medicine have been carried out. The data provided by these studies made it possible on the one hand to explain the therapeutic action and on the other hand to

confirm the use of these different plants in traditional medicine [8]. *Talinum triangulare*, a cosmopolitan weed of the *Talinum* genus, of the *Talinaceae* family (formerly *Portulacaceae*), fits into this theme. This plant has great therapeutic values in the traditional medicine system, but is not yet fully exploited [9]. Preliminary work carried out on the aqueous and hydro-ethanolic extracts (70%) of its leaves made it possible to demonstrate the antimalarial properties of the leaves of this plant on clinical isolates of *Plasmodium falciparum* [10]. According to the work of Tano [6] when a person has an attack of malaria, plasmodia can lead to oxidative stress through the massive lysis of red blood cells. Hemmer et al. [11] also demonstrated that neutrophils activated by parasites can trigger the destruction of endothelial cells. This can be prevented by *ex vivo* intake antioxidants and proteolytic enzyme inhibitors. In sum, antioxidants can attenuate endothelial cell damage due to parasite-induced leukocyte activation, supporting the importance of antioxidant activity in the context of malaria treatment [12]. The present study is part of continuing investigations into the antimalarial activity of *Talinum triangulare*. More specifically, the aim was to evaluate the antioxidant activity of extracts and fractions of *Talinum triangulare* leaves and to determine their contents of phenolic compounds, particularly total polyphenols, total flavonoids and total tannins.

## 2. MATERIALS AND METHODS

### 2.1 Materials

The plant material consists of leaves of *Talinum triangulare* (Jacq.) Willd., belonging to the

Talinaceae family and well known under the name wild spinach. This plant was identified by Professor AKE Assi, at the national floristic center located in Abidjan, Cocody University Campus, under the specimen UCJ014548.

## 2.2 Methods

### 2.2.1 Harvesting, drying and spraying *Talinum triangulare* leaves

The leaves of *Talinum triangulare* were harvested between July and August 2021 at the start of the morning (6:30 a.m.) in Koffikro, a camp located 12 km from the Daloa Department and in the Tazibouo district within the UEESO-CI city near COPRO college of Daloa Department (Haut-Sassandra Region, Ivory Coast). After harvest, they were freed of all other elements, dried in the shade and in the open air for ten weeks then crushed until a fine powder was obtained using a mechanical grinder. The resulting powder was stored at room temperature in glass jars to avoid mold.

### 2.2.2 Preparation of extracts and fractions of *Talinum triangulare* leaves

The extraction of the active ingredients was generally done by cold maceration in a given solvent using a blender for a specific time.

### 2.2.3 Preparation of the aqueous extract

The fine powder initially obtained was extracted using the method of Zirihi et al. [13]. To do this, a quantity of one hundred grams of the vegetable powder was cold macerated in one liter of distilled water using a blender. After three minutes of homogenization, the homogenate obtained was collected in a square of clean white tissue and then pressed by hand. The collected solution was filtered twice through hydrophilic cotton and then once through Whatman filter paper (3 mm). The filtrate obtained was dried in an oven at a temperature of 60 °C for (five) 5 days to obtain the total aqueous extract (Aq).

### 2.2.4 Preparation of the hydro-ethanolic extract

It followed the same principle as the previous one, with the only difference that instead of water, an ethanol-water mixture (70/30: v/v) was used. It made it possible to obtain the hydro-ethanolic extract (HOH) after drying for 72 hours in an oven at 60°C.

These two extracts (Aq, HOH) were stored separately in airtight jars at laboratory temperature.

### 2.2.5 Fractionation of the hydro-ethanolic extract

With a view to improving the activity of the crude extract having given good antiplasmodial activities (HOH), the latter underwent partition chromatography, using solvents of increasing polarity: hexane, dichloromethane, l ethyl acetate and butanol. This method consists of liquid-liquid partitioning with solvents that are immiscible with each other. The partition method is inspired by those of Bolou & al. [14]. One hundred grams of crude extracts (HOH) having given good antiplasmodial activities were dissolved in 500 mL of distilled water and transferred to a 1000 mL separatory funnel. And this aqueous solution was extracted with 3 x 250 mL of hexane. The combined hexanic phases were evaporated to constitute the hexanic fraction. The residual aqueous phase was successively extracted with dichloromethane, ethyl acetate and butanol three times for each solvent. The dichloromethane, ethyl acetate and butanolic phases as well as the residual aqueous phase after treatment with organic solvents were evaporated separately in order to obtain a dichloromethane fraction (DCM), an ethyl acetate fraction, a butanolic fraction and an aqueous fraction (FAq).

## 2.3 Antioxidant Activity

### 2.3.1 Measurement of antioxidant activity by the DPPH test

The DPPH method is a test for the reduction of the 2,2-diphenyl-1-picrylhydrazyl (DPPH<sup>•</sup>) radical to 2,2-diphenyl-1-picrylhydrazine (DPPH-H). It makes it possible to evaluate the antioxidant potential of a sample, by its capacity to trap the DPPH<sup>•</sup> radical used as a chromogen [15,16]. This organic radical is stable at room temperature and has a purple color. The antioxidants present in the sample, by transferring an H atom to the DPPH<sup>•</sup> radical, reduce it to DPPH-H or DPPH<sub>2</sub>. This reduction causes a change in the color of the solution from purple to pale yellow [17]. For its production, a quantity of 4 mg of DPPH powder was dissolved in 100 mL of methanol, homogenized then stored away from light to constitute a solution. At the same time, a quantity of 30 mg of each extract to be tested was diluted in 3 mL of sterile distilled water in order to have an extract concentration of

10 mg/mL. Then, a volume of 1.5 mL of methanol was distributed into ten sterile test tubes numbered from 1 to 10 to constitute the experimental series of a given extract. In tube 1 of the experimental series, a volume of 1.5 mL of the extract with a concentration of 10 mg/mL previously prepared was added and then mixed to constitute the stock. It consisted of taking 1.5 mL from tube 1, diluting it in tube 2 then homogenizing it. This operation was repeated up to tube 10. From tube 10, the excess volume was rejected. Thus, the various concentrations varied from 10 to 0.019 mg/mL from tube 1 to tube 10. After which, a volume of 1.5 mL of DPPH was added to every 10 tubes. Alongside this experimental series, a reference series was produced. In fact, it consisted of using vitamin C instead of the plant extract, and always in 10 tubes. The only difference between the experimental and reference series is that the concentrations of the reference product used ranged from 25 to 0.195 µg/mL from tube 1 to tube 10. However, in parallel with the experimental and reference series, a control control was prepared by soliciting only methanol and DPPH at equal volume (1.5 mL). All of the tubes were kept away from light for 30 minutes then their absorbance was determined distinctly with a spectrophotometer at 517 nm with methanol as the blank control and vitamin C as a positive control. The percentage of inhibition of the DPPH radical was calculated according to the following equation: solution. After this step, the double dilution technique of geometric ratio ½ in medium was applied :

$$\text{DPPH inhibition (\%)} = (A_0 - A_{\text{sample}}) / A_0 \times 100$$

*A<sub>0</sub>*: absorbance of the control (drug-free). *A*: absorbance of the sample after 30 min.

### 2.3.2 Measurement of antioxidant activity by ABTS

The method used was that described by Choong et al., [18]. It is based on the ability of compounds to reduce the ABTS<sup>+</sup> cation radical (2,2'-azinobis-3-ethylbenzothiazoline-6-sulfonic acid). It was produced by the reaction of 8 mM ABTS (87.7 mg in 20 mL distilled water) and 3 mM potassium persulfate (0.0162 g in 20 mL distilled water) in a 1:1 ratio (v/v). The mixture was then incubated in the dark at room temperature for 12 to 16 hours to obtain an ABTS<sup>+</sup> solution. It was diluted with methanol to obtain a solution whose absorbance was 0.7 ± 0.02 at 734 nm. Then, a volume of 3.9 mL of this

diluted ABTS<sup>+</sup> solution was added to 100 µL of each compound to be tested (aqueous and hydro-ethanolic extracts, dichloromethane and aqueous fractions) in different test tubes then shaken vigorously. The initial concentration of these test compounds is 10 mg/mL. After shaking, the mixture was incubated for 6 minutes in the dark ( $T = 30 \pm 2 \text{ }^\circ\text{C}$ ). The residual absorbance of the ABTS<sup>+</sup> radical was then measured at 734 nm with a UV-visible spectrophotometer and should be between 20%-80% of the absorbance of the white. The tests were carried out in triplicate and the results were expressed in µmol Trolox equivalent per liter of extract (µmol TE/L). The inhibition rate (%) of ABTS<sup>+</sup> was expressed as follows:

$$\% I = [(A_0 - \text{Compound Abs}) / A_0] \times 100$$

*A<sub>0</sub>* = diluted ABTS absorbance. *Compound Abs* = diluted ABTS absorbance + sample

## 2.4 Spectrophotometric Determination of Certain Secondary Metabolites

### 2.4.1 Determination of total flavonoids

The method of Marinova et al. [19] was used for the determination of total flavonoids. It consisted of putting a volume of 0.75 mL of sodium nitrite (NaNO<sub>2</sub>) at 5 % (m/v) into a 25 mL flask, adding 2.5 mL of extract then homogenizing it. To this mixture, 0.75 mL of aluminum chloride (AlCl<sub>3</sub>) at 10% (m/v) was added, then incubated for 6 minutes in the dark. After the incubation, a volume of 5 mL of sodium hydroxide (1N NaOH) was added and then the volume was made up to 25 mL with distilled water. The mixture was shaken vigorously before being measured with a UV-visible spectrophotometer. The reading was taken at 510 nm. The tests were carried out in triplicate. The flavonoid content was expressed in grams per liter of quercetin equivalent extract.

### 2.4.2 Determination of total polyphenols

The determination of total polyphenols was carried out according to the method described by Wood et al. [20] To do this, to a volume of 30 µL of extract or fraction, 2.5 mL of Folin-Ciocalteu reagent diluted 1/10 was added. The mixture obtained was kept for 2 minutes in the dark at room temperature ( $27 \pm 3 \text{ }^\circ\text{C}$ ) then 2 mL of sodium carbonate solution at 75 g/L was added and then homogenized. The solution obtained was then incubated at 50 °C for 15 minutes. The absorbance reading was carried out with a UV-

visible spectrophotometer at a wavelength of 760 nm against a blank consisting of distilled water. Gallic acid was used as a reference standard for the quantification of total polyphenol contents expressed in grams per liter of gallic acid equivalent extract (g.L-1, Eq AG). The tests were carried out in triplicate for each sample.

### 2.4.3 Determination of total tannins

The condensed tannin content was determined according to the method described by Julkunen-Titto [21]. To carry it out, a volume of 50 mL of each fraction or the hydro-alcoholic extract was added to 1500  $\mu$ L of the 4 % solution of vanillin in methanol then stirred vigorously. To this mixture, 750  $\mu$ L of concentrated hydrochloric acid was added and the whole mixture was then left to stand at room temperature for 20 minutes. The absorbance was measured at a wavelength of 550 nm against a blank consisting of the 4% solution of vanillin in methanol. The tests were carried out in triplicate for each sample. A stock solution of tannic acid was used as a reference standard for the establishment of the calibration curve and for the quantification of the contents of condensed tannins expressed in milligram equivalent of tannic acid per gram of dry matter (mg EAT /g of dry matter). The tests were carried out in triplicate for each sample.

### 2.5 Statistical Analyzes

The curves and histograms are drawn using Microsoft Excel 2013 and GraphPad Prism 5 software. The IC<sub>50</sub> values (50% inhibitory concentration) were calculated by the linear regression method from the curve [% inhibition = f (concentrations)].


## 3. RESULTS

### 3.1 Antioxidant Activity by the DPPH Test


The antioxidant activity of extracts ((aqueous (Aq) and hydro-ethanolic (HOH)) and fractions (dichloromethane (DCM) and aqueous (Faq)) of *Talinum triangulare* leaves and that of ascorbic acid (vitamin C) was represented by graphs showing the variation in the percentage of free radical inhibitory power as a function of the concentration of each compound (Figs. 1 and 2). The DCM and Faq fractions of *Talinum triangulare* leaves gave respective IC<sub>50</sub>s of 0.9 and 2.3 mg/mL As for the aqueous (Aq) and hydro-ethanolic (HOH) extracts, their IC<sub>50</sub> could not be defined from the linear regression equations of the graphs. Regarding vitamin C (reference molecule), it provided an IC<sub>50</sub> of 3.056  $\mu$ g/mL. It is also noted that the reduction of free radicals at the level of the two fractions (DCM and Faq) and the reference molecule (vitamin C) is a function of the concentration of the substance used.

### 3.2 Antioxidant Activity by ABTS<sup>+</sup>


Fig. 3 shows the proton trapping capacity of the cationic radical ABTS<sup>+</sup> of the various products used (plant extracts and fractions). These results generally reveal that the products requested inhibited the absorbance of the ABTS<sup>+</sup> radical. However, the percentage of inhibition is greater for the DCM fraction (41.59 %) than that of Faq (40.59 %). While that of Faq is more remarkable than that of the HOH extract (38.29 %). Of the two extracts, only the Aq extract has the lowest percentage of inhibition (25.51 %).



**Fig. 1. Percentage of DPPH inhibition as a function of different concentrations of ascorbic acid (reference molecule)**



**Fig. 2. Percentage of DPPH inhibition as a function of concentrations of various extracts and fractions of *Talinum triangulare* leaves**



**Fig. 3. Percentage of inhibition of the absorbance of the ABTS<sup>+</sup> radical various extracts and fractions of *Talinum triangulare* leaves**

### 3.3 Total Flavonoid Contents

Fig. 4 is the result of the total flavonoid contents of the extracts (aqueous and hydro-ethanolic) and fractions (DCM and Faq) of the leaves of *Talinum triangulare* (Talinaceae). It generally reveals that the aqueous extract (Aq) contains more total flavonoids ( $10.16667 \pm 0.289$  mg EQ/g of extract) than the DCM fraction and the HOH extract with approximately equal respective values of  $2.16667 \pm 0.577$  and  $1.8333 \pm 0.577$  mg EQ/g of extract. As for the Faq fraction, it is characterized by a low presence of total flavonoids ( $0.5 \pm 0.00$  mg EQ/g).

### 3.4 Total Polyphenol Contents

The results of the polyphenol contents are presented in Fig. 5. It should generally be noted that this content is greater in the HOH extract

( $4.8333 \pm 0.289$  mg EAG/g) than in the Faq fraction ( $4.5 \pm 0.289$  mg EAG/g), 0.500 mg EAG/g). After these, follow the Aq extract ( $4.333 \pm 0.289$  mg EAG/g) and the DCM fraction ( $3.8333 \pm 0.289$  mg EAG/g).

### 3.5 Total Tannin Contents

The results of the total tannin dosage test are given in Fig. 6. They show overall that the total tannin content is relatively low in all of the *Talinum triangulare* compounds studied. However, the tannin level is a little more abundant in the HOH extract ( $3.3228333 \pm 0.0254$   $\mu$ g EAT/g of dry extract) than in the Aq extract ( $2.970667 \pm 0.00318$   $\mu$ g EAT/g of dry extract). After these, the DCM fraction has more total tannins than the Faq fraction with respective contents of  $2.5096667 \pm 0.00029$  and  $1.835333 \pm 0.01384$   $\mu$ g EAT/g of dry extract.




Fig. 4. Total flavonoid contents of the Aq and HOH extracts, and of the DCM and Faq fractions of the leaves of *Talinum triangulare* (Talinaceae).




Fig. 5. Total polyphenol contents of the Aq and HOH extracts, and of the DCM and Faq fractions of the leaves of *Talinum triangulare* (Talinaceae).




Fig. 6. Total tannin contents of the Aq and HOH extracts, and of the DCM and Faq fractions of the leaves of *Talinum triangulare* (Talinaceae)

#### 4. DISCUSSION

The evaluation of the antioxidant activity of the 70 % aqueous and hydro-ethanolic extracts, as

well as the residual dichloromethane and aqueous fractions of *Talinum triangulare*, reveals that the aqueous and hydro-ethanolic extracts did not make it possible to determine the IC<sub>50</sub> via

the DPPH test. In comparison, vitamin C, used as a reference molecule, showed a low  $IC_{50}$ . These results suggest through the inhibition activity curves that the reduction of free radicals is a function of the concentration of the substances tested, confirming the dose-dependent nature of the antioxidant activity. Compared to the work of Kouakou [22] who reported  $IC_{50}$ s of 0.398 and 0.235 mg/mL for methanolic extracts of Shea hulls and cakes, *Talinum triangulare* fractions show lower anti-radical activity. Vitamin C is approximately 294 times (0.9 mg/3.056  $\mu$ g) more effective than the dichloromethane fraction and approximately 752 times (2.3 mg/3.056  $\mu$ g) more effective than the residual aqueous fraction, highlighting the superior effectiveness of the vitamin C as a pure antioxidant. In contrast, the dichloromethane fraction is approximately twice as effective as the residual aqueous fraction (2.3/0.9). This result is supported by the work of Athamena et al. [23] who showed that the percentage of inhibition of free radicals varies depending on the solvent used for the extraction and the concentrations of the anti-radical compounds. The results of Tidian et al. [24] report a lower vitamin C  $IC_{50}$  in *Vitellaria paradoxa* (shea) extract, but our  $IC_{50}$  for vitamin C is approximately three times higher than theirs, which may reflect differences in sample quality or the experimental conditions. According to Barkat and Laib [25] a lower  $IC_{50}$  is associated with better antioxidant activity. However, despite the higher  $IC_{50}$  of *Talinum triangulare* fractions compared to vitamin C, their antioxidant activity remains significant.

The results of the ABTS<sup>+</sup> radical absorbance inhibition test revealed a higher percentage for the dichloromethane fraction compared to the residual aqueous fraction, the hydro-ethanolic extract and the aqueous extract. They thus corroborate those obtained with the DPPH test on the antioxidant capacity of the different compounds contained in the plant products tested and show comparable performance, but highlight the hydro-ethanolic extract and the aqueous extract as having antioxidant capacities notable, although lower than the dichloromethane fraction.

This study also highlighted the presence of phenolic compounds such as flavonoids, polyphenols and total tannins at different proportions in the plant products used. The analysis of phenolic compounds showed a higher concentration of total flavonoids in the aqueous extract, followed by the dichloromethane fraction,

the hydro-ethanolic extract, and the residual aqueous fraction. Total polyphenols are most abundant in the hydro-ethanolic extract, followed by the residual aqueous fraction, the aqueous extract, and the dichloromethane fraction. Total tannins are most concentrated in the hydro-ethanolic extract, followed by the aqueous extract, the dichloromethane fraction, and the residual aqueous fraction.

These results confirm the observations of N'guessan et al. [26] and Bidié et al. *al.* [27] who found a correlation between the level of total phenols and antiradical activity. Furthermore, the work of Chen and Ho [28] demonstrated that the functional groups of phenolic compounds, such as flavonoids and polyphenols, are capable of donating electrons or protons to neutralize free radicals. Thus, the antioxidant activity of the requested plant products previously demonstrated by the DPPH test and the ABTS<sup>+</sup> radical-cation reduction test could be due to the presence of these phenolic compounds (polyphenols, flavonoids and total tannins) in the leaves of *Talinum triangulare*. The results of this study are different from those of Soro [29] who revealed the absence of phenolic compounds such as polyphenols and total flavonoids in *Rhynchospora corymbosa* extract.

## 5. CONCLUSION

The study demonstrated that *Talinum triangulare* leaf extracts and fractions have variable antioxidant activity, with notable effectiveness of the dichloromethane fraction. Although vitamin C remains the most potent antioxidant among the compounds tested, the plant's fractions offer significant antioxidant potential, justifying their traditional use in the treatment of malaria. The results suggest that phenolic compounds present in *Talinum triangulare* compounds contribute to this antioxidant activity. These results support the interest of this plant in traditional pharmacopoeia and encourage further research to isolate and characterize the active compounds responsible for its therapeutic properties.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.



## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Tagne MLL. Malaria: Knowledge, prevention practices and therapeutic routes in Koulouba, Sogonafing and point G. State Doctorate Thesis in Medicine, Faculty of Medicine and Odontostomatology (FMOS), University of Sciences, Techniques and Technologies of Bamako, Mali. 2021;71.
2. Gentilini M, Dufflo B. Malaria In: Medicine tropical ed, flammarion, 1993;81-108.
3. Bagayoko KN. Incidence and lethality of serious and complicated malaria in the pediatric intensive care unit of CHU Gabriel TOURE. Doctorate thesis in Medicine, Faculty of Medicine, Pharmacy and Odontostomatology of Bamako, Mali. 2008;89.
4. WHO ; 2023  
Available :<https://www.afro.who.int>. World Malaria Day 2024.
5. Salery MR. Antimalarial activity of girolin: *in vitro* and *in vivo* study. National Veterinary School of Toulouse, Paul-Sabatier University of Toulouse. Veterinary Medicine Thesis. 2007;158.
6. Tano KD. Study of the antiplasmodial activity of various extracts of *Terminalia glaucescens* (Combretaceae) and *Erigeron floribundus* (Asteraceae). Toxicological profile for the formulation of improved traditional antimalarial medicines (ATMs). Single thesis on Valorization of natural substances (Option: Parasitology), UFR of Pharmaceutical and Biological Sciences, Félix Houphouët-Boigny University, Ivory Coast. 2016;237.
7. Nkasa HL, Clément ICL, Muzomwe MJC, Masengo EI, Mumba CM, Taba KM. Phytochemical screening and antibacterial activity of phytomedicine mathesia, a drug use against buruli ulcer in Democratic Republic of Congo (DRC). European Journal of Pharmaceutical and Medical Research (EJPMR). 2020;7(2):52-56.
8. Mariam K, Obouayeba AP, Ackah BJAA, Dosso K. Phytochemical sorting and biotolerance of an improved traditional medicine used in the treatment of hepatitis B in Côte d'Ivoire. World Journal of Pharmaceutical and Life Sciences (WJPLS). 2024;10(3):25-36.
9. Swarna J, Ravindhran R. Pharmacognostical and phytochemical evaluation of *Talinum triangulare* (Jacq.) Willd. International Journal of Pharmacy and Pharmaceutical Sciences. 2013;5(2):249-256.
10. Okou OC, Tano KD, Koffi AJ, Koffi AF, Yavo W, Djaman AJ. Evaluation of the antiplasmodial activity of *talinum triangulare* (talinceae), a west African food plant. European Journal of Pharmaceutical and Medical Research; 2019.
11. Hemmer CJ, Lehr HA, Westphal K, Unverricht M, Kratzius M, Reisinger EC. *Plasmodium falciparum* Malaria: reduction of endothelial cell apoptosis *in vitro*." Infect Immun. 2005;73(3):1764-1770.
12. Pincemail J, Bonjean K, Cayeux K, Defraigne JO. Physiological action of antioxidant defense." Clinical Nutrition and Metabolism. 2002;16:233-239.
13. Zirih GN, Kra AKM, Guédé-Guina F. Evaluation of the antifungal activity *Microglossa pyrifolia* (Lamarck, O. Kuntze) (Asteraceae) "PYMI" on the *in vitro* growth of *Candida albicans*. Revue de Médecine et de Pharmacie. 2003;17:11-18.
14. Bolou GEK, Attioua B, N'guessan AC, Coulibaly A, N'guessan JD, Djaman AJ. *In vitro* evaluation of the antibacterial activity of *Terminalia glaucescens* plant extracts, on *Salmonella typhi* and *Salmonella Typhimurium*. Bulletin of the Royal Society of Sciences. 2011;80:772-790.
15. Brand-Williams W, Cuvelier ME, Berset CLWT. Use of radical method to evaluate antioxidant activity. LWT-Food Science and Technology, 1995;28(1) :25-30.
16. Benzie IFF, Szeto YT. Total antioxidant capacity of teas by the ferric reducing/antioxidant power assay. Journal Agricultural Food Chemical. 1999; 47(2):633 -636.
17. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. Songklanakarin Journal of Sciences and Technology. 2004;26(2) :211 -219.
18. Choong C, Van-Den TT, Roger F, Roger MCFL. Antioxidant activities, phenolic and beta-carotene contents of sweet potato

- genotypes with varying flesh colors. Food Chemistry. 2007;103(3):829-838.
19. Marinova D, Ribarova F, Atanassova M. Total phenolics in Bulgarian Fruits and vegetables. *Journal of the University of Chemical Technology and Metallurgy*, T. 2005 ;40(3):255-260.
  20. Wood JE, Senthilmohana ST, Peskinb AV. Antioxidant activity of procyanidin-containing plant extracts at different pHs. Food Chemistry. 2002;77(2):155–161.
  21. Julkunen-Titto R. Phenolic constituents in the leaves of Northern Willows: Methods for the Analysis of certain phenolics. *J. Agric. Food Chem.* 1985;33:213-217.
  22. Kouakou AB. Antioxidant, antimicrobial and healing effectiveness of cosmetic products formulated from shea nut shells and cakes (*Vitellaria paradoxa*). Agrifood Biotechnology Thesis. UFR Biosciences, Félix Houphouët-Boigny University, Ivory Coast. 2022;129.
  23. Athamena S, Chalghem I, Kassah-Laouar A, Laroui S, Khebri S. Antioxidant and antimicrobial activity of extracts of *Cuminum cyminum* L. *Lebanese Science Journal.* 2010;11(1):69-81.
  24. Tidiane K, Ouattara GA, Monon K, Abdoulaye T, Karamoko O. Phytochemical study and antioxidant activity of stem bark extracts of *Vitellaria Paradoxa* C.F.Gaertn, a medicinal plant used in the North of Ivory Coast. *European Scientific Journal.* 2021;17(34) :241. Available:<https://doi.org/10.19044/esj.2021.v17n34p241>
  25. Barkat M, Laib I. Chemical composition and antioxidant activity of the essential oil of dry flowers of *Lavandula officinalis*.2011;6:46-54.
  26. N'guessan JD, Zirih GN, Kra AKM, Kouakou K, Djaman AJ, Guede-Guina F. Free radical scavenging activity, flavonoid and phenolic contents of selected Ivorian plants. *IJONAS.* 2007;4:425-429.
  27. Bidie AP, N'guessan BB, Yapo AF, N'guessan JD, Djaman AJ. Antioxidant activities of ten medicinal plants from the Ivorian pharmacopoeia. Antioxidant activities of ten medicinal plants from the Ivorian pharmacopoeia. *Science & Nature.* 2011;8(1):1–11.
  28. Chen CW, Ho CT. Antioxidant properties of polyphenols extracted from green tea and black tea. *J Lipids.* 1995;2:35-46.
  29. Soro M. Identification and characterization of secondary metabolites that inhibit the polymerization of Hbs hemoglobins from *Rhynchospora corymbosa* and *Olax subscorpioidea*; plants used in the treatment of sickle cell disease in the Poro Region (Ivory Coast). Doctoral Thesis in Biology-Health, Specialty: Biochemistry and Pharmacology of Natural Substances, UFR Agroforestry, Jean Lorougnon Guédé University (Daloa), Ivory Coast. 2023;179.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124698>