



Establishment of *In vitro* Regeneration Protocol of *Aegle marmelos* (L.): A Sacred Medicinal Tree

Wagh YG^{a++}, Aher AR^{b#*}, Chimote VP^{b†}, Kale AA^{b‡}
and Jambhale VM^{c#}

^a Post Graduate Institute, MPKV, Rahuri (Maharashtra), India.

^b State Level Biotechnology Centre, MPKV, Rahuri (Maharashtra), India.

^c Medicinal and Aromatic Plant Project, MPKV, Rahuri (Maharashtra), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors contributed to the study's conception and design, study of the work, material preparation, data collection, analysis and manuscript preparation. All authors have read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i111685>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126784>

Original Research Article

Received: 09/09/2024

Accepted: 13/11/2024

Published: 19/11/2024

ABSTRACT

Bael (*Aegle marmelos* L.) is an important medicinal tree and sacred tree species of the Hindu religion. This study aimed to develop an efficient *in vitro* regeneration protocol for Bael, to overcome challenges in large-scale propagation due to low seed germination. For the regeneration study, the different explants such as the nodal segments and shoot tips were used for the proliferation of the

⁺⁺ Student (M. Sc. Agri);

[#] Assistant Professor (GPB);

[†] Associate Professor (Mol Bio & Biotech);

[‡] Associate Professor (Biochem);

*Corresponding author: E-mail: arahr76@gmail.com;

Cite as: YG, Wagh, Aher AR, Chimote VP, Kale AA, and Jambhale VM. 2024. "Establishment of *In Vitro* Regeneration Protocol of *Aegle Marmelos* (L.): A Sacred Medicinal Tree". *Journal of Advances in Biology & Biotechnology* 27 (11):1006-16. <https://doi.org/10.9734/jabb/2024/v27i111685>.

explant with MS medium, however, only the nodal segment showed positive results. To avoid the browning of the media the explants were kept in distilled water for 60 minutes and then sterilized. The maximum survival of explants (93.33%) and less contamination percentage (6.67%) was observed when explants were washed and treated with 0.1% Tween-20 concentration for 15 minutes + 0.1% bavistin and 0.05% streptocycline treatment for 25 minutes and finally 70% ethanol treatment for 30 seconds. Different hormonal combinations viz. BAP alone, BAP + IAA, and Kinetin + IAA were used to initiate shooting. Among all eighteen treatments, the BAP 2.5 mg/l showed a maximum number of shoots per explant (8.33 ± 2.88), the highest shoot initiation (93.33%), and the earliest initiation (30 ± 5.25 days). For shoot multiplication 2.5 mg/l BAP + 0.5 mg/l NAA showed maximum shoots per explant (28.00 ± 5.08) and fastest shoot proliferation. The regenerated multiple shoots were separated from each other and transferred to half MS media with IBA of different concentrations for rooting. Among the six different treatments only two treatments viz. $\frac{1}{2}$ MS media + 1.0 mg/l IBA and $\frac{1}{2}$ MS media + 1.5 mg/l IBA recorded best root length. For primary hardening the rooted plants were planted in vermicompost: sand: soil at 1:1:2 proportion and then after one and half months survived plants were again transferred to soil: organic manure (2:1) for secondary hardening.

Keywords: *Aegle marmelos*; regeneration; shooting; rooting; hardening.

1. INTRODUCTION

Aegle marmelos is a tropical deciduous and highly heterozygous fruit tree. The plant is very hardy and can adapt to a wide range of habitats, from arid to mesophytic (Arya & Shekawat, 1986; Singh et al. 1976; and Zaman, 1988). It belongs to the Rutaceae family and is an economically important multipurpose social tree of the Indo-Burman subcontinent (Hooker, 1875), that is now being introduced into tropical countries in Western Africa and South America. It has been around since prehistoric times in India and is also known as Bael, Bengal Quince, Golden Apple, Japanese Bitter Orange, Stone Apple or Wood Apple. It is considered auspicious, sacred and, a symbol of prosperity in ancient Sanskrit poems. Hindus worship Lord Shiva with its trifoliate compound leaves. The Bael appears frequently in the Sanskrit scriptures of Vedic times (C 2000BC - C 800BC). Bael is mentioned in the Yajurveda, early Buddhist and Jain literature (C 800BC - C 325BC), the Ramayana, and the '*Brihat Samhita*' (Aiyer, 1956; Prakash, 1961; Singh & Roy, 1984). As a sacred tree, the bael tree is usually planted around Hindu temples.

The root is a component of '*dasmula*' (10 roots), an ayurvedic treatment, widely used by practitioners. The roots have a variety of tastes and flavors, including sweet, bitter, and febrifuge. Bael fruit is significant because of its curative properties, which makes the tree one of India's most useful medicinal plants (Monika et al., 2023). The leaves are astringent, laxative, febrifuge, and expectorant, and can be used to

treat ophthalmia, deafness, inflammations, catarrh, diabetes, and asthmatic symptoms. The unripe fruits are bitter, acrid, sour, astringent, digestive, and stomachic, and are excellent in diarrhea and dysentery. The ripe fruits are sweet, aromatic, cooling, febrifuge, laxative, and tonic, and are beneficial to the brain and heart in disorders (Warrier et al., 1996). The pulp of the ripe fruit is used in the preparation of sherbet, which is an aromatic summer drink. Bael can also be used for forestation in dry areas due to its high drought tolerance capacity (Anonymous, 1985).

The Bael fruit is extremely nutritious and high in vitamins and minerals. Per 100g of edible portion, it contains 1.8 g protein, 0.39 g fat, 1.7 g minerals, 37.8 g carbohydrates, 55 mg carotene, 0.13 mg thiamine, 1.19 mg riboflavin, 1.1 mg niacin, and 8 mg vitamin C (Gopalan et al., 1971). Riboflavin is abundant in bael fruit (Mukherjee & Ahmad, 1957). It also contains marmelosin and has 88 calories of energy per 100g of fruit pulp, making it more nutritious than other fruits (Jauhari & Singh, 1971).

The Bael is typically propagated by seed. However, because it is a cross-pollinated species, seed propagation results in a highly heterozygous population (Ajithkumar & Seeni, 1998). Seeds also have short viability and are vulnerable to insect attack. Plantlets from seed grow slowly and are susceptible to pathogens and pests in the early stages. The rate of vegetative proliferation via root suckers is slow also difficult and time-consuming (Ray & Chattarjee, 1996).

Conventional propagation methods, such as seeds or root cuttings and layers are associated with problematic issues such as seasonal seed production (which takes a year), pest attacks and low seed germination. Non-conventional ways, such as *in vitro* propagation procedures, can help solve these issues. The *in vitro* multiplication of endangered species could be important for those taxa whose reproduction is problematic through conventional methods. Thus, the development of a specific sterilization technique for plant materials obtained from the field is an essential requirement before identifying the growth stage for the collection of plant materials and perfecting the medium composition for the multiplication of plants. Therefore, the present study was conducted to develop an *in vitro* regeneration protocol for the multiplication of bael.

2. MATERIALS AND METHODS

This study was conducted at the State Level Biotechnology Centre, Mahatma Phule Krishi Vidyapeeth, Rahuri. Plant materials *viz.* nodal segment and shoot tip were collected in the morning hours from healthy mature plants from the experimental orchard of Medicinal and Aromatic Plants Project (Dhanwantary Garden), Mahatma Phule Krishi Vidyapeeth, Rahuri.

The glassware, and instruments used for aseptic manipulations were sterilized as per standardized procedure. For the sterilization of explants, different sterilants were used and the best combination was identified. Sterilized explants were inoculated on MS (Murashige & Skoog, 1962) media with full-strength basal medium supplemented with 18 combinations of different plant growth hormones for shoot proliferation to identify the best treatment for shoot initiation. They were initially kept in the dark for a week and then incubated at 25°C with a 16 h light period and 8 h dark period. Initiated shoots were then transferred to 8 different media combinations to identify the best treatment for the shoot multiplication. For the root initiation study, different concentrations of IBA with half MS media were used. As per Das et al. (2008), six treatments were formulated to carry out rooting experiments, and the best treatment was identified. Plantlets thus produced were then transferred to pots containing different combinations of sterile soil, sand, vermicompost and coco peat. All the pots were irrigated at regular intervals. They were initially covered with plastic bags for a week and then kept in a polycarbonated polyhouse. All the experiments

were conducted in a Completely Randomized Block design with three replicates and 10 explants per replication were used. The data collected were analyzed for mean and standard error as given by Snedecor & Cochran (1967). Calculations of standard error for the number of days required for the multiplication, number of shoots formed per explant, shoot initiation percentage, number of days required for the rooting, length of roots formed and rooting percentage were done. Each replication's value was transformed by using the square root transformation method.

3. RESULTS AND DISCUSSION

3.1 Explant Preparation

The present study describes the development of a rapid and efficient plantlet regeneration protocol in bael using nodal segments. For the proliferation of the explants, different explants *viz.* nodal segments and shoot tips were used with a modified MS medium. For shoot initiation, only nodal segments showed positive results (73.33%), while in the case of shoot tip not a single explant developed the shoot (0%), and later on all the shoot tip explants were dried off. Hence, for further experimental study, only nodal segments were used as explants. Due to the presence of pre-existing meristem which can easily be developed into shoots, plant regeneration from nodal explant is one of the most promising ways of multiplying a selected species to its true type (Mandal et al., 2021).

The nodal segments were collected from newly emerging branches of healthy plants from the Dhanwantary Garden, MPKV, Rahuri. The explants of 0.5 cm to 1.0 cm in length were dissected from the mother plant and immediately the slant cut (near about 45° angle) was given at the bottom of the explants. In preliminary experiments, darkening or browning of the culture medium proved toxic to plant growth and hindered the growth; hence the explant was kept in double distilled water for 60 minutes to avoid the browning of media. Most of the phenolic compound was released in water and further, no browning was observed.

The various sterilants were used in different concentrations to standardize the surface sterilization protocol for obtaining the contamination-free culture. The part and partial similar procedures described by earlier workers were followed for surface sterilization of the

nodal segment. First, the explants were disinfected with Tween-20 solution (0.1%) for 15 minutes and then washed with double distilled water four times. Further sterilized by Bavistin and Streptocycline, maximum survival of explant 93.33% and less contamination (6.67%) was observed with 0.1% Bavistin and 0.05% Streptocycline concentration for 25 minutes (Table 1).

Then the explants were transferred to the inoculation room. In laminar airflow, the explants were treated with 0.1% HgCl₂ for 2 minutes and thoroughly washed with double distilled water 4 times to remove the traces of mercuric chloride and again treated with 70% ethanol for 30 seconds and washed with double distilled water 4 to 5 times and finally inoculated on culture media. The duration of treatment is important, long duration of treatment will damage the explants or can cause browning (Singhal et al., 2011).

3.2 Shoot Initiation

The type and concentration of a growth regulator were found to have a significant impact on morphogenetic responses in bael. The results of different experiments using different hormone concentrations and combinations are presented here. The MS media gave a good response to explant proliferation. The different types of growth regulator concentrations were used for the regeneration study (Fig. 1). The aim was to identify the best media combination for the shoot initiation. For the proliferation of explant BAP alone with different concentrations, the BAP + IAA combination and Kinetin + IAA combination were used. Among all the treatments, the T₅ treatment *i.e.* BAP 2.5 mg/l showed best shoot initiation (93.33%), fastest shoot proliferation

(30.00 ± 5.25 days) and higher shoot initiation (8.33± 2.88) followed by T₉ treatment (BAP 1.0 mg/l + IAA 0.3 mg/l) (86.67%, 31.00 ± 5.33, 6.33 ± 2.55) and T₁₇ Kinetin 2.0 mg/l + IAA 0.4 mg/l (86.67%, 31.33 ± 5.36, 5.00 ± 2.31). Because of the better response to shooting as compared to others, BAP (2.5 mg/l) proved to be the best media combination for the shoot proliferation (Table 2). Similar results were reported by Bindhu (2015) and Pathirana et al. (2020).

3.3 Shoot Multiplication

The initiated, well-developed, 45 to 60 days-old explants were transferred to the multiplication media. But, initially, all explants were dried off and could not survive. Then some of the bottom portion of initiated explants was removed and directly transferred to the multiplication media so the survival percentage was increased. All explants were cultured on multiplication media containing BAP + NAA growth regulator with different concentrations (Fig. 2). Among all the treatments M₈ *i.e.* BAP 2.5 mg/l + NAA 0.5 mg/l showed maximum shoots per explant (28.00 ± 5.08). The shoot establishment was 93.33% and the number of days required for shoot proliferation was 60.00 (± 7.82). Hence, it was accepted as the best treatment for shoot multiplication (Table 3). Das et al. (2008) reported that the highest percentage of multiple shoots (91.23%) within 40 days and the maximum number of multiple shoots per explant (22.7) was found in BAP 2.0 mg/l with NAA 0.2 mg/l. Hossain et al. (1994) reported the maximum frequency of adventurous bud proliferation was observed in MS medium with BAP and NAA. Agarwal & Bhardwaj (2024) reviewed advances in *in vitro* regeneration studies in bael.

Table 1. Effect of sterilization treatments on explants

Treatment	Time	Survival (%)	Contamination / Dead (%)
Bavistin (0.1%) + Streptocycline (0.05%)	15 min	46.66	53.33
	20 min	60.00	40.00
	25 min	93.33	6.67
Bavistin (0.2%) + Streptocycline (0.1%)	15 min	33.33	66.67
	20 min	40.00	60.00
	25 min	46.67	53.33
Bavistin (0.1%) + Streptocycline (0.2%)	15 min	53.33	46.67
	20 min	60.00	40.00
	25 min	66.67	33.33
Bavistin (0.2%) + Streptocycline (0.2%)	15 min	46.67	53.33
	20 min	66.67	33.33
	25 min	86.67	13.33

Table 2. Effect of different BAP concentrations on shoot proliferation

Treatment No.	Growth regulator (mg/l) added to MS media	Shoot establishment %	No. of days required for shoot proliferation (Mean ± SE)	No. of shoots per explant (Mean ± SE)
T ₁	0.5 BAP	60.00	33.00 ± 5.50	2.33 ± 1.72
T ₂	1.0 BAP	53.33	31.33 ± 5.36	3.33 ± 1.96
T ₃	1.5 BAP	86.67	31.00 ± 5.33	7.00 ± 2.67
T ₄	2.0 BAP	66.67	32.67 ± 5.47	6.00 ± 2.49
T₅	2.5 BAP	93.33	30.00 ± 5.25	8.33 ± 2.88
T ₆	3 BAP	66.67	34.67 ± 5.63	4.33 ± 2.18
T ₇	1.0 BAP + 0.1 IAA	73.33	31.83 ± 5.40	3.67 ± 2.04
T ₈	1.0 BAP + 0.2 IAA	80.00	32.27 ± 5.44	3.67 ± 2.04
T₉	1.0 BAP + 0.3 IAA	86.67	31.00 ± 5.33	6.33 ± 2.55
T ₁₀	1.0 BAP + 0.5 IAA	66.67	33.33 ± 5.52	5.00 ± 2.31
T ₁₁	2.0 BAP + 0.1 IAA	53.33	32.00 ± 5.42	3.33 ± 1.89
T ₁₂	2.0 BAP + 0.2 IAA	60.00	31.20 ± 5.35	3.00 ± 1.89
T ₁₃	1.0 Kinetin + 0.05 IAA	53.33	31.50 ± 5.37	3.00 ± 1.89
T ₁₄	2.0 Kinetin + 0.1 IAA	73.33	32.00 ± 5.42	5.33 ± 2.37
T ₁₅	2.0 Kinetin + 0.2 IAA	66.67	31.00 ± 5.33	3.33 ± 1.96
T ₁₆	2.0 Kinetin + 0.3 IAA	80.00	31.67 ± 5.39	4.00 ± 2.11
T₁₇	2.0 Kinetin + 0.4 IAA	86.67	31.33 ± 5.36	5.00 ± 2.31
T ₁₈	2.0 Kinetin + 0.5 IAA	53.00	33.33 ± 5.52	3.00 ± 1.89

Table 3. Effect of different concentrations of BAP + NAA on multiplication of shoots

Treatment No.	Growth Regulator BAP +NAA (mg/l) added to MS media	Shoot establishment %	No. of days required for shoot proliferation (Mean ± SE)	No. of shoots per explant (Mean ± SE)
M ₁	2.0 BAP + 0.1 NAA	46.67	60.43 ± 7.85	10.33 ± 3.17
M ₂	2.0 BAP + 0.2 NAA	80.00	60.70 ± 7.86	18.00 ± 4.11
M ₃	2.0 BAP + 0.3 NAA	73.33	60.55 ± 7.85	18.67 ± 4.18
M ₄	2.0 BAP + 0.5 NAA	66.67	60.45 ± 7.85	10.00 ± 3.13
M ₅	2.5 BAP + 0.1 NAA	86.67	60.30 ± 7.84	21.67 ± 4.49
M ₆	2.5 BAP + 0.2 NAA	60.00	61.67 ± 7.93	10.00 ± 3.13
M ₇	2.5 BAP + 0.3 NAA	80.00	61.33 ± 7.90	25.00 ± 4.81
M₈	2.5 BAP + 0.5 NAA	93.33	60.00 ± 7.82	28.00 ± 5.08

Table 4. Effect of different concentrations of IBA on root formation

Treatment No.	Growth Regulator IBA (mg/l) added to ½ MS media	% of root initiation	No. of days required for root proliferation (Mean ± SE)	Average length of roots (cm) (Mean ± SE)
R ₁	½ MS media + 0.1 IBA	-	-	-
R ₂	½ MS media + 0.5 IBA	-	-	-
R₃	½ MS media + 1.0 IBA	33.33	90.67 ± 9.03	2.90 ± 1.97
R ₄	½ MS media + 1.5 IBA	20.00	91.00 ± 9.04	2.67 ± 1.91
R ₅	½ MS media + 2.0 IBA	-	-	-
R ₆	½ MS media + 2.5 IBA	-	-	-



Fig. 1. Initiation of nodal segment of *Aegle marmelos* L. on MS medium with different concentration of BAP mg/l, BAP mg/l + IAA mg/l and Kinetin mg/l + IAA mg/l

1a. MS + BAP 0.5 mg/l

1c. MS + BAP 1.5 mg/l

1e. MS + BAP 2.5 mg/l

2a. MS + BAP 1.0 mg/l + IAA 0.1 mg/l

2c. MS + BAP 1.0 mg/l + IAA 0.3 mg/l

2e. MS + BAP 2.0 mg/l + IAA 0.1 mg/l

3a. MS + Kinetin 1.0 mg/l + IAA 0.05 mg/l

3c. MS + Kinetin 2.0 mg/l + IAA 0.2 mg/l

3e. MS + Kinetin 2.0 mg/l + IAA 0.4 mg/l

1b. MS + BAP 1.0 mg/l

1d. MS + BAP 2.0 mg/l

1f. MS + BAP 3 mg/l

2b. MS + BAP 1.0 mg/l + IAA 0.2 mg/l

2d. MS + BAP 1.0 mg/l + IAA 0.5 mg/l

2f. MS + BAP 2.0 mg/l + IAA 0.2 mg/l

3b. MS + Kinetin 2.0 mg/l + IAA 0.1 mg/l

3d. MS + Kinetin 2.0 mg/l + IAA 0.3 mg/l

3f. MS + Kinetin 2.0 mg/l + IAA 0.5 mg/l




Fig. 2. Multiplication of *Aegle marmelos* L. on MS medium and BAP mg/l + NAA mg/l combination with different concentrations

a. MS + BAP 2.0 mg/l + NAA 0.1 mg/l
c. MS + BAP 2.0 mg/l + NAA 0.3 mg/l
e. MS + BAP 2.5 mg/l + NAA 0.1 mg/l
g. MS + BAP 2.5 mg/l + NAA 0.3 mg/l

b. MS + BAP 2.0 mg/l + NAA 0.2 mg/l
d. MS + BAP 2.0 mg/l + NAA 0.5 mg/l
f. MS + BAP 2.5 mg/l + NAA 0.2 mg/l
h. MS + BAP 2.5 mg/l + NAA 0.5 mg/l




Fig. 3. Rooting of *Aegle marmelos* L. on $\frac{1}{2}$ MS medium and IBA mg/L with different concentrations
a & b. $\frac{1}{2}$ MS + IBA 1.0 mg/L
c & d. $\frac{1}{2}$ MS + IBA 1.5 mg/L mg/L



Fig. 4. Hardening of regenerated tissue culture plantlets of *Aegle marmelos* L.
a. Primary hardening (Vermicompost: Sand : Soil at 1:1:2)
b. Secondary hardening (Soil: Organic manure at 2:1)

Table 5. Hardening of the regenerated plants

Sr. No.	Type of hardening	Hardening composition	Plant survival percentage
1.	Primary hardening	Sterile soil: Coco peat: Sand (1:1:1) Vermicompost: Sand: Soil (1:1:2)	60% 80%
2.	Secondary hardening	Soil + Organic manure (2:1)	80%

3.4 *In vitro* Rooting

The regenerated multiple shoots were separated from each other and transferred to half MS media with IBA of different concentrations. The observations were recorded and depicted in. Among the six different treatments only two treatments R₃ (½ MS media + 1.0 mg/l IBA) and R₄ (½ MS media + 1.5 mg/l IBA) showed root proliferation (Table 4 and Fig. 3). The rooting treatment R₃ recorded 2.90 cm ± 1.97 and R₄ recorded root length of 2.67 cm ± 1.91. The percent root initiation was higher in R₃ (33.33%). The number of days required for root proliferation was approximately similar (R₃: 90.67 ± 9.03; R₄: 91.00 ± 9.04) in both treatments. As a hard woody plant species, the root formation was slow, and more duration was required for root induction. Similar results were reported by Das et al. (2008) and Akter et al. (2013).

3.5 Hardening for Plant Acclimatization

The regenerated *in vitro* plants were taken out after weeks from the inoculated bottle and washed with double distilled water to remove the media traces. The rooted portion was treated with 0.1% Bavistin solution to prevent further fungal infection. For primary hardening, the rooted plants were transferred to the small pots containing two different potting hardening mixtures of sterile Soil: Coco peat: Sand in the proportion of 1:1:1 and Vermicompost: Sand: Soil of 1:1:2 respective. To maintain the proper moisture level pots were drilled to make proper holes and kept under controlled environment conditions in the greenhouse for one month. The plant survival rate for sterile Soil: Coco peat: Sand (1:1:1) was 60% and Vermicompost: Sand: Soil (1:1:2) was 80%. After one month when new leaves emerged the plants were shifted to a new potting mixture containing Soil: Organic manure (2:1) for secondary hardening. After one month old plant survival rate was 80%. Results obtained in hardening are presented in Table 5 and Fig. 4. Similar results were reported by Das et al. (2008). After sufficient acclimatization, the plants were transferred to natural conditions and showed 80% plantlet survival.

4. CONCLUSION

The nodal segment 0.5cm to 1.0cm in length showed better response for shoot initiation. To avoid the browning of the media, the explants were kept in distilled water for 60 minutes immediately after removal from the mother plant. The explant was kept free from contamination by using surface sterilization such as Tween 20 (0.1%) for 15 minutes, Bavistin (0.1%) + Streptocycline (0.05%) for 25 minutes, Mercuric chloride 0.1% treated for 2 minutes and 70% ethanol for 30 seconds. The M.S. Medium supplemented with BAP 2.5 mg/l was the best media combination for shoot proliferation. For shoot multiplication, M.S. Medium supplemented with 2.5 mg/l BAP + 0.5 mg/l NAA proved to be the best media combination. Likewise, ½ MS medium with 1.0 mg/l IBA gave the best result for root development. For primary hardening Vermicompost: sand: soil at 1:1:2 proportion and for secondary hardening soil: organic manure (2:1) gave the better response for plant survival. The current investigation optimizes an effective high frequency *in vitro* regeneration from nodal segments of *A. marmelos*. This will help in micropropagation of desired bael plants.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

We acknowledge the Dean (Faculty of Agriculture) and Director of Instruction, Mahatma Phule Krishi Vidyapeeth, Rahuri and Associate Dean, Post Graduate Institute, Mahatma Phule Krishi Vidyapeeth, Rahuri for allowing us to undertake this research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Agarwal, K., & Bhardwaj, R. (2024). Advances in micropropagation techniques of *Aegle marmelos* (L.) Corr.: A review. *Micropropagation of Medicinal Plants*, 99–116.
<https://doi.org/10.2174/97898152383031240201>
- Aiyer, A. K. Y. N. (1956). *The antiquity of some field and forest flora of India*. Bangalore Printing and Publishing Co. Ltd., Bangalore.
- Ajithkumar, D., & Seeni, S. (1998). Rapid clonal multiplication through in vitro axillary shoot proliferation of *Aegle marmelos* (L.) Corr., a medicinal tree. *Plant Cell Reports*, 17(5), 422–429.
<https://doi.org/10.1007/s002990050418>
- Akter, S., Banu, T. A., Habib, M. A., Afrin, S., Khatun, A., Khan, S., & Islam, S. (2013). In vitro clonal multiplication of *Aegle marmelos* (L.) Corr. through cotyledonary node culture. *Bangladesh Journal of Scientific and Industrial Research*, 48(1), 13–18.
<https://doi.org/10.3329/BJSIR.V48I1.15408>
- Anonymous. (1985). *The wealth of India; Raw materials* (Vol. 1 (A), Revised Series, Ed. Chadha, Y. R., pp. 85–91). Council for Scientific and Industrial Research, New Delhi. <https://doi.org/10.2307/4109779>
- Arya, H. C., & Shekawat, N. S. (1986). Clonal multiplication of tree species in the Thar desert through tissue culture. *Forest Ecology and Management*, 16, 201–208.
[https://doi.org/10.1016/0378-1127\(86\)90020-4](https://doi.org/10.1016/0378-1127(86)90020-4)
- Bindhu, K. B. (2015). In vitro propagation of *Aegle marmelos* through nodal explants. *International Journal of Science and Research*, 4(4), 691–694.
- Das, R., Hasan, M. F., Rahman, M. S., Rashid, M. H., & Rahman, M. (2008). Study on in vitro propagation through multiple shoots proliferation in wood apple (*Aegle marmelos* L.). *International Journal of Sustainable Crop Production*, 3(6), 16–20.
- Gopalan, C. B. N., Rama Sastri, B., & Balasubramanian, R. B. (1971). *Nutritive value of Indian foods* (pp. 1–578). National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India.
- Hooker, J. (1875). *The flora of British India* (Vol. I, pp. 516–517). Reeve.
<https://doi.org/10.5962/bhl.title.678>
- Hossain, M., Islam, R., Karim, M. R., Joarder, O. J., & Biswas, B. K. (1994). Regeneration of plantlets from in vitro cultured cotyledons of *Aegle marmelos* Corr. (*Rutaceae*). *Scientia Horticulturae*, 57(4), 315–321.
[https://doi.org/10.1016/0304-4238\(94\)90114-7](https://doi.org/10.1016/0304-4238(94)90114-7)
- Jauhari, O. S., & Singh, R. D. (1971). Bael—valuable fruit. *Indian Horticulture*, 16(1), 9–10.
- Mandal, S., Parsai, A., Tiwari, P. K., & Nataraj, M. (2021). The effect of additional additives on the axillary shoot micropropagation of medicinal plant *Aegle marmelos* (L.) Correa. *An International Scientific Journal*, 34, 54–71.
- Monika, S., Thirumal, M., & Kumar, P. R. (2023). Phytochemical and biological review of *Aegle marmelos* Linn. *Future Science OA*, 9(3), FSO849. <https://doi.org/10.2144/fsoa-2022-0068>
- Mukherjee, B., & Ahmad, K. (1957). Riboflavin. *Pakistan Journal Biology & Agriculture Science*, 1, 47–51.
- Murashige, T., & Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15, 473–497.
<https://doi.org/10.1111/j.1399-3054.1962.tb08052.x>
- Pathirana, C., Attanayake, U., Dissanayake, U., Gamlath, L., Ketipearachchi, K., Madhujith, T., & Eeswara, J. (2020). Establishment of a micropropagation protocol for elite accessions of bael (*Aegle marmelos* (L.) Corr.), a tropical hardwood species. *Advances in Agriculture*, 2020, 8840386.
<https://doi.org/10.1155/2020/8840386>
- Prakash, O. (1961). *Food and drinks in ancient India: From earliest times to c. 1200 AD*. Munshi Ram Manohar Lal, Oriental Book Sellers and Publishers. NII Book ID BA01590741.
- Ray, D. P., & Chattarjee, B. K. (1996). Effect of different concentrations of growth regulators, etiolation and invigoration treatments on the rooting of stem cutting of bael (*Aegle marmelos* Corr.). *Orissa Journal of Horticulture*, 24(1–2), 36–41.
- Singh, R. N., & Roy, S. K. (1984). *The Bael, cultivation and processing* (pp. 1–25). Indian Council of Agricultural Research, New Delhi.
- Singh, U. R., Pandey, I. C., Upadhyaya, N. P., & Prasad, R. S. (1976). Propagation

- of Bael (*Aegle marmelos*) by budding. *Punjab Horticultural Journal*, 16(20), 57–59.
- Singhal, V. K., Salwan, A., Kumar, P., & Kaur, J. (2011). Phenology, pollination and breeding system of *Aegle marmelos* (Linn.) Correa (*Rutaceae*) from India. *New Forests*, 42(1), 85–100. <https://doi.org/10.1007/s11056-010-9239-3>
- Snedecor, G. W., & Cochran, W. G. (1967). *Statistical Methods 6th*. Edlowa State University Press. Ame, Iowa. USA, 2, 304-8.
- Warrier, P. K., Nambiar, V. P. K., & Ramankutty, C. (1996). *Indian medicinal plants* (Vol. 1, pp. 62–66). Orient Longman.
- Zaman, M. F. (1988). *Bangladesher phaller chash*. Bangla Academy. Dhaka, Bangladesh, 189–193.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/126784>