



Prevalence of Oral Screen Appliance Therapy in 6-12 Year Old Children Visiting a University Dental Hospital: A Retrospective Study

Arunn Jaikumarr Ram ^a and Mebin George Mathew ^{b*}

^a Dental Research Cell, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India.

^b Department of Pedodontics and Preventive Dentistry, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2022/v34i42B36300

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83925>

Original Research Article

Received 03 January 2022

Accepted 06 March 2022

Published 22 June 2022

ABSTRACT

The oral screen is a versatile and simple myofunctional instrument that is used to treat dental arc deformities early in the interception. A lip training device was described as an oral screen. Oral screens can be used for people who have bad habits such as mouth breathing, chewing nails, big overjet and lip muscle hypotonus. A sample of 4355 patients of age 6-12 years visiting the hospital were taken. We reviewed patients' records, analysed data of 7415 patients between June 2020 to March 2021 and clinical findings are recorded. Out of the sample size 4355,260 patients in the hospital database were diagnosed with orthodontic deformities and use of oral screen was recommended. The data is then tabulated in Microsoft Excel. Chi square test is used for comparison of groups. The data is analysed with the help of SPSS software. In this study we observed that 3.8% of the total patients (260) showed incidence of undergoing treatment using oral screen. Prevalence of oral screen appliance is observed to be more in age groups between 10-12 years followed by 6-9 years. Within the limitations of the study, 0.2% of the children visiting university hospital have undergone treatment using oral screen appliances.

^o Research Associate,

*Corresponding author: E-mail: mebingeorgem.sdc@saveetha.com, geomani.sdc@saveetha.com;

Keywords: Oral screen appliance; mouth breathing; children; orthodontic appliance and treatment; innovative technique.

1. INTRODUCTION

The oral screen is a versatile and simple myofunctional instrument that is used to treat dental arc deformities early in the interception. It is necessary to establish better muscle balance between the tongue and the buccinator mechanism, and to restore normal growth and development [1,2]. Functional devices refer to a variety of devices designed to modify the structure of different muscle groups which influence the function and the location of the mandible to transmit strength to the dentition and basal bone [3]. Newell introduced it for the first time in 1912. In Britain before the Second World War, oral screens were routinely used [4]. Kraus created a double oral screen and differs from the oral screen to the vestibular screen [5]. Later, Hotz, Nord and Fingeroth were in favour of oral screening [6-8].

In recent years, there has been questioning the importance of intermittent forces from the lips, cheeks and tongue for the position of the teeth [9]. This is particularly true of the forces exerted during swallowing and speech, while the constant strength of the rest of the muscles is assumed to be significant. Incompetent lips may contribute to the protuberance of the teeth by reducing the pressure on the teeth from those in the lips [10].

A lip training device was described as an oral screen. The screen extends the lip muscles and gives a reverse force to proclination incisors [11]. At the same time the arc of the lips are strengthened. If a loop is provided on the labial surface, the training effect can be increased [12]. The tendency of the screen to discharge from the mouth causes the lips to contract with the external force when the loop is tightened by finger strength [13]. It is believed to strengthen the lips in the regular application of external force with accompanying lip contraction; that is to say, the lips will be able to exercise more force as a result of their training [14]. Our team has extensive knowledge and research experience that has translate into high quality publications [15-30].

Oral screens can be used for people who have bad habits such as mouth breathing , chewing nails, big overjet and lip muscle hypotonus [31,32]. The oral screen is one of the choices

[33,34]. The objective of the study was to assess the prevalence of oral screen appliances in 6–12-year-old children visiting a dental hospital.

2. MATERIALS AND METHODS

This retrospective study examined the records of patients from 01 June 2020 to 31st March 2021 who visited university dental hospital. The study population included patients with age ranging between 6-12 years. The study sample included both male and female gender, predominantly south Indians.

The study population was 7415 pediatric patients who visited university hospital. Sample size was 4355 pediatric patients in which 260 patients in the hospital database were diagnosed with orthodontic deformities and use of oral was recommended. The necessary data such as age, gender and number of people undergone treatment was recorded. Incomplete patient records were excluded. Data was recorded in Microsoft Excel and exported to the statistical package of social science for windows (SPSS) and subjected to statistical analysis. Chi square tests are used for comparison of groups.

3. RESULTS AND DISCUSSION:


The study sample consisted of 4355 patients, in which about 260 patients were recommended the use of oral screen appliances. Out of 260 about 139 children belonged to the age group between 6-9 years and 121 children belonged to the age group between 10-12 years (Graph -1). Among 260, 137 of the children were boys and 123 of the children were girls (Graph -2).

Among 260 patients only 10 underwent treatment with oral screen appliance and remaining 250 did not opt for the treatment with oral screen appliance (Graph -3).


When age and treatment were compared, Children in groups 6-9 years, out of 139 children only 4 underwent treatment with oral screen and 135 underwent treatment without oral screen. Children in the group 10-12 years, out of 121 children only 6 underwent treatment with oral screen and 115 underwent treatment without oral screen (Graph -4).

When gender and treatment were compared, out of 137 boys only 6 boys underwent treatment with oral screen and 131 underwent treatment without oral screen. Out of 123 girls only 4 girls underwent treatment with oral screen and 119 underwent treatment without oral screen (Graph - 5).


This study found that of 4355 patients attending the university hospital ,260 were advised of the use of oral screen appliance, out of which only 10 underwent treatment using oral screen appliance.




Graph 1. Pie chart showing age distribution of children in sample population requiring oral screen appliance therapy. Dark blue colour denotes children of age group between 6 to 9 years and green colour denotes children of age group between 10 to 12 years. 121 of the children belonged to the age group of 10 to 12 years and 139 of the children belonged to the age group between 6 to 9 years




Graph 2. Pie chart showing gender distribution of children in sample population requiring oral screen appliance therapy. green colour denotes girls and blue colour denotes boys. 137 of the children were boys and 123 of the children were girls



Graph 3. Pie Chart showing prevalence of treatment in the study population where blue colour denotes treatment with oral screen appliance. Green colour denotes treatment without oral screen. Prevalence of oral screen appliance was only 0.2%, out of 4355 patients 260 patients was suggested with oral screen appliance therapy but only 10 patients underwent oral screen appliance therapy.



Graph 4. Bar chart shows the association between various age groups and treatment .X axis represents age group and Y axis represents frequency distribution of treatment. Blue colour denotes treatment with oral screen appliance, green colour denotes treatment without oral screen appliance. Age group of 10-12 has more patients(6) using oral screen for treatment when compared to age group 6-9(4).Chi square test was done and the association was found to be not significant.(p value was =0.757 p >0.05 statistically not significant). The maximum number of people using oral screen was seen in the age group of 10-12 years old however the difference was not significant



Graph 5. Bar chart showing association of treatment with gender. X axis represents gender and Y axis represents frequency of treatment. Blue colour denotes treatment with oral screen appliance, green colour denotes treatment without oral screen appliance. Between boys and girls, boys have undergone treatment with oral screen more than the girls. Chi square test was done, and the association was found to be not significant. (p value was =0.223 p>0.05 statistically not significant). The maximum number of people using oral screen was seen in males rather than females

The prevalence was 0.2 % which is almost equal compared to the study of Casey [35]. The results of oral screen appliance are similar to the results of other studies conducted in Asian region [36]. The prevalence of male who underwent the treatment was more than the female which is similar to the study of Vasui [37].

Early diagnosis and treatment should be followed in these patients to improve the quality of their life by preventing the crowding of teeth. The limitation of this study was, since our study was carried out in a uncenter study in the South Indian population, it cannot be generalised into any other population. Other cases which could underestimate the common occlusal properties were not included in addition to orthodontic cases. Subjects for direct examination are not available. Further research is needed as it is related with aesthetics and has a functional concern. For additional diagnosis and treatment planning, the study may be extended. Proper advice on early correction awareness is

necessary. The research thus serves as proof and adds to the consensus that may be used with the broader population and clinical studies for further studies.

4. CONCLUSION

Within the limits of the study, about 0.2 % of children visiting university dental hospitals among 4355 had the prevalence of oral screen appliances. Children of 10 to 12years had significantly higher prevalence of the oral screen appliance therapy. Higher prevalence of treatment with oral screen appliance was seen in boys (131) than in girls although the difference was not statistically significant (p=0.223).

ETHICAL APPROVAL

Ethical Approval was taken from the institutional review board/ SDC/SIHEC/DIASDATA/0618-0320.

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

ACKNOWLEDGEMENT

The authors would like to acknowledge the help and support rendered by the Department of Pedodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gadda D, Nosetti LM, Caprioglio A, Levrini L, Nespoli L. Oral Modifications Due To Mouth Breathing In Children With Sleep-Disordered Breathing (SDB) [Internet]. B73. PEDIATRIC SLEEP; 2011. Available: http://dx.doi.org/10.1164/ajrccm-conference.2011.183.1_meetingabstracts.a3715
2. Sano M, Sano S, Kato H, Arakawa K, Arai M. Proposal for a screening questionnaire for detecting habitual mouth breathing, based on a mouth-breathing habit score [Internet]. BMC Oral Health. 2018;18. Available:<http://dx.doi.org/10.1186/s12903-018-0672-6>
3. Jallad WA, Al Jallad W. Restoration of Missing Upper Anterior Teeth using Dental Implant Simultaneous with Bone Grafting- A Case Report [Internet]. Journal of Dentistry & Oral Health. 2014;1-8. Available:<http://dx.doi.org/10.17303/jdoh.2013.303>
4. Phulari B. History of Removable Orthodontic Appliances [Internet]. History of Orthodontics. 2013;160-160. Available:http://dx.doi.org/10.5005/jp/books/12065_17
5. Babu S, Schutt CA, Bojrab DI. Diagnosis and Treatment of Vestibular Disorders [Internet]. Springer. 2019;391. Available:<https://play.google.com/store/books/details?id=XCWFDwAAQBAJ>
6. Hotz R. Orthodontie in der täglichen Praxis: Möglichkeiten und Grenzen im

- Rahmen der Kinderzahnheilkunde [Internet]. 1980;502. Available:https://books.google.com/books/about/Orthodontie_in_der_t%C3%A4glichen_Praxis.html?hl=&id=nk9qAAAAMAAJ
7. Graber TM, Neumann B. Removable Orthodontic Appliances [Internet]. W B Saunders Company; 1984;631. Available:https://books.google.com/books/about/Removable_Orthodontic_Appliances.html?hl=&id=8htqAAAAMAAJ
 8. Levine JM, Fingerth JM. Historical and Current Nomenclature Associated with Intervertebral Disc Pathology [Internet]. Advances in Intervertebral Disc Disease in Dogs and Cats. 2014;25-31. Available:<http://dx.doi.org/10.1002/9781118940372.ch4>
 9. Owman-Moll P, Ingervall B. Effect of oral screen treatment on dentition, lip morphology, and function in children with incompetent lips [Internet]. American Journal of Orthodontics. 1984;85:37-46. Available: [http://dx.doi.org/10.1016/0002-9416\(84\)90121-0](http://dx.doi.org/10.1016/0002-9416(84)90121-0)
 10. Tosello DO, Vitti M, Berzin F. EMG activity of the orbicularis oris and mentalis muscles in children with malocclusion, incompetent lips and atypical swallowing-Part II [Internet]. Journal of Oral Rehabilitation. 1999;26:644-9. Available:<http://dx.doi.org/10.1046/j.1365-2842.1999.00409.x>
 11. Nesbitt RA, Marley JJ, James J. Cheilitis glandularis: report of an unusual case affecting the upper lip [Internet]. Vol.3 Oral Surgery. 2010. Oral Surg.; 32-35. – no. Available: <http://dx.doi.org/10.1111/j.1752-248x.2010.01075.x>
 12. Kumar M, Goyal M, Bhavna J, Tomar S, Kushwah A. An Innovative Procedure for Lip Lengthening in a Patient with a Short Upper Lip and High-Angle Skeletal Class II Pattern: A Case Report [Internet]. Journal of Indian Orthodontic Society. 2021;030157422110076. Available:<http://dx.doi.org/10.1177/03015742211007618>
 13. Serdev NP. Excision of Prominent Posterior Septal Angle and Nasal Spine for Downward Tip Rotation, in Short Upper Lip, or Over-Rotated Tip [Internet]. Miniinvasive Techniques in Rhinoplasty; 2016. Available: <http://dx.doi.org/10.5772/62062>
 14. Graber TM. The use of muscle forces by simple orthodontic appliances [Internet].

- V American Journal of Orthodontics. 1979;76:1–20.
Available: [http://dx.doi.org/10.1016/0002-9416\(79\)90295-1](http://dx.doi.org/10.1016/0002-9416(79)90295-1)
15. Subramanyam D, Gurunathan D, Gaayathri R, Vishnu Priya V. Comparative evaluation of salivary malondialdehyde levels as a marker of lipid peroxidation in early childhood caries. *Eur J Dent* [Internet]. 2018;12(1):67–70. Available: http://dx.doi.org/10.4103/ejd.ejd_266_17
 16. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig* [Internet]. 2019;23(9):3543–50. Available: <http://dx.doi.org/10.1007/s00784-018-2775-5>
 17. Ramakrishnan M, Dhanalakshmi R, Subramanian EMG. Survival rate of different fixed posterior space maintainers used in Paediatric Dentistry – A systematic review [Internet]. *The Saudi Dental Journal*. 2019;31:165–72. Available: <http://dx.doi.org/10.1016/j.sdentj.2019.02.037>
 18. Jeevanandan G, Thomas E. Volumetric analysis of hand, reciprocating and rotary instrumentation techniques in primary molars using spiral computed tomography: An in vitro comparative study. *Eur J Dent* [Internet]. 2018;12(1):21–6. Available: http://dx.doi.org/10.4103/ejd.ejd_247_17
 19. Princeton B, Santhakumar P, Prathap L. Awareness on Preventive Measures taken by Health Care Professionals Attending COVID-19 Patients among Dental Students. *Eur J Dent* [Internet]. 2020;14(S 01):S105–9. Available: <http://dx.doi.org/10.1055/s-0040-1721296>
 20. Saravanakumar K, Park S, Mariadoss AVA, Sathiyaseelan A, Veeraraghavan VP, Kim S, et al. Chemical composition, antioxidant, and anti-diabetic activities of ethyl acetate fraction of *Stachys riederi* var. *japonica* (Miq.) in streptozotocin-induced type 2 diabetic mice. *Food Chem Toxicol* [Internet]. 2021 Jun 26;155:112374. Available: <http://dx.doi.org/10.1016/j.fct.2021.112374>
 21. Wei W, Li R, Liu Q, Devanathadesikan Seshadri V, Veeraraghavan VP, Surapaneni KM, et al. Amelioration of oxidative stress, inflammation and tumor promotion by Tin oxide-Sodium alginate-Polyethylene glycol-Allyl isothiocyanate nanocomposites on the 1,2-Dimethylhydrazine induced colon carcinogenesis in rats. *Arabian Journal of Chemistry* [Internet]. 2021;14(8):103238. Available: <https://www.sciencedirect.com/science/article/pii/S1878535221002537>
 22. Gothandam K, Ganesan VS, Ayyasamy T, Ramalingam S. Antioxidant potential of theaflavin ameliorates the activities of key enzymes of glucose metabolism in high fat diet and streptozotocin - induced diabetic rats. *Redox Rep* [Internet]. 2019;24(1):41–50. Available: <http://dx.doi.org/10.1080/13510002.2019.1624085>
 23. Su P, Veeraraghavan VP, Krishna Mohan S, Lu W. A ginger derivative, zingerone-a phenolic compound-induces ROS-mediated apoptosis in colon cancer cells (HCT-116). *J Biochem Mol Toxicol* [Internet]. 2019;33(12):e22403. Available: <http://dx.doi.org/10.1002/jbt.22403>
 24. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial [Internet]. *Clinical Oral Investigations*. 2020;24:3275–80. Available: <http://dx.doi.org/10.1007/s00784-020-03204-9>
 25. Sekar D, Johnson J, Biruntha M, Lakhmanan G, Gurunathan D, Ross K. Biological and Clinical Relevance of microRNAs in Mitochondrial Diseases/Dysfunctions. *DNA Cell Biol* [Internet]. 2020;39(8):1379–84. Available: <http://dx.doi.org/10.1089/dna.2019.5013>
 26. Velusamy R, Sakthinathan G, Vignesh R, Kumarasamy A, Sathishkumar D, Nithya Priya K, et al. Tribological and thermal characterization of electron beam physical vapor deposited single layer thin film for TBC application. *Surf Topogr: Metrol Prop* [Internet]. 2021 Jun 24 [cited 2021 Aug 10];9(2):025043. Available: <https://iopscience.iop.org/article/10.1088/2051-672X/ac0c61/meta>
 27. Aldhuwayhi S, Mallineni SK, Sakhamuri S,

- Thakare AA, Mallineni S, Sajja R, et al. Covid-19 Knowledge and Perceptions Among Dental Specialists: A Cross-Sectional Online Questionnaire Survey. Risk Manag Healthc Policy [Internet]. 2021;14:2851–61. Available:<http://dx.doi.org/10.2147/RMHP.S306880>
28. Abirami S, Navaneethan R, Varghese R. Comparison between antegonial notch depth, symphysis morphology and ramus morphology among different growth patterns in skeletal class I and class II subjects [Internet]. International Journal of Research in Pharmaceutical Sciences. 2020;11:1975–83. Available:<http://dx.doi.org/10.26452/ijrps.v11ispl3.3703>
29. G NK, Nithya KG, Bhagya LT, Prabhu D. Radiographic evaluation of permanent second molar development based on nollas stage of tooth development in 9-11-year-old male children [Internet]. International Journal of Research in Pharmaceutical Sciences. 2020;11:469–74. Available:<http://dx.doi.org/10.26452/ijrps.v11ispl3.2968>
30. Radiographic Evaluation of Permanent Canine Development Based on Nollas Stage of Tooth Development In 6–8-Year-Old Male Children [Internet]. Journal of Contemporary Issues in Business and Government. 2021;27. Available:<http://dx.doi.org/10.47750/cibg.2021.27.02.035>
31. Singh H. To Assess the Influence of Habitual Mouth Breathing and its Associated Taste Alterations [Internet]. Open Access Journal of Dental Sciences. 2020;5. Available: <http://dx.doi.org/10.23880/oajds-16000274>
32. Lee SY, Guilleminault C, Chiu HY, Sullivan SS. Mouth breathing, “nasal disuse,” and pediatric sleep-disordered breathing [Internet]. Sleep and Breathing. 2015;19:1257–64. Available:<http://dx.doi.org/10.1007/s11325-015-1154-6>
33. Sjögreen L. Review for “Reliability and comparability of methods for assessing oral function: chewing, tongue pressure, and lip force” [Internet]; 2020. Available:<http://dx.doi.org/10.1111/joor.12976/v1/review2>
34. Hansen H, Leber AB, Saygin ZM. What sound sources trigger misophonia? Not just chewing and breathing [Internet]. Available:<http://dx.doi.org/10.31234/osf.io/8y9ek>
35. Caseley R. Principles and construction of the oral screen as a functional appliance. Dent Rec [Internet]. 1947;67(10):230–5. Available:<https://www.ncbi.nlm.nih.gov/pubmed/18915843>
36. Prevalence of various removable functional appliances - an institutional set up [Internet]. Journal of Contemporary Issues in Business and Government. 2021;27. Available:<http://dx.doi.org/10.47750/cibg.2021.27.02.247>
37. Yasui EM, Kimura RK, Kawamura A, Akiyama S, Morisaki I. A modified oral screen appliance to prevent self-inflicted oral trauma in an infant with cerebral palsy: a case report [Internet]. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology. 2004;97:471–5. Available:<http://dx.doi.org/10.1016/j.tripleo.2003.10.010>

© 2022 Ram and Mathew; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83925>