



A Cross Sectional Study on Some Aspects of Reproduction Scenario and Prevalence of Reproductive Disorders in Dairy Cattle in Bangladesh

M. R. Islam ^{a*}, M. N. Zahan ^a, H. Hossain ^a, A. A. Maruf ^{b^o}, M. R. Molla ^{c^o},
M. S. I. Sajib ^{d[#]} and M. M. Hossain ^e

^a Department of Surgery and Theriogenology, Faculty of Animal Science and Veterinary Medicine, Sher-e-Bangla Agricultural University, Dhaka - 1207, Bangladesh.

^b Department of Livestock Services, Austagram, Kishoreganj - 2380, Bangladesh.

^c Department of Livestock Services, Rupganj, Narayanganj - 1464, Bangladesh.

^d Livestock and Dairy Development Project, Department of Livestock Services, Dhaka - 1215, Bangladesh.

^e Department of Anatomy, Histology and Physiology, Faculty of Animal Science and Veterinary Medicine, Sher-e-Bangla Agricultural University, Dhaka - 1207, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91021>

Original Research Article

Received 17 June 2022
Accepted 22 August 2022
Published 27 August 2022

ABSTRACT

A cross sectional survey was carried out in dairy herds in four districts of Bangladesh to determine some aspects of reproduction and the prevalence of reproductive disorders that affect the dairy cattle industry. A total of 2850 dairy cows (local 1020; 35.8% and crossbred 1830; 64.2%) were surveyed in 256 local (204; 79.7%) and commercial (52; 20.3%) dairy herds located in Dhaka, Narayanganj, Gazipur, and Manikganj were investigated using interviewer-administered questionnaire survey and face-to face interviews with the owners. Sizes of the dairy herds visited

^o Upazila Livestock Officer;

[#] Livestock Extension Officer;

*Corresponding author: E-mail: rashed.suth@sau.edu.bd, rashed036@gmail.com;

were small (1-5 cows; 47.7%), medium (6-10 cows; 30.5%), and large (≥ 10 cows; 21.8%). The estrus of most of the dairy herd cows (197/256; 76.9%) were determined by observing signs or taking history from owners. Furthermore, only 26.1% (67/256) of the dairy herds kept reproductive records using a log book or a computer, whereas 32.8% (84/256) of the dairy herds used a reproductive calendar for breeding purposes. The results indicated that 71.5% (146/204) locals were using artificial insemination (AI) for the dairy cows reproduction when classified according to service method while, 73.1% (38/52) commercials were using AI other than natural service (NS), and AI and NS. The use of the Department of Livestock Service (DLS) semen rate was higher 67.2% (137/204) in locals while, Bangladesh Rural Advancement Committee (BRAC) semen rate was higher 61.5% (32/52) in commercials. According to the study's findings, anestrus rates of 8.6% (88/1020) in locals and repeat breeding syndrome (RBS) rates of 9.8% (180/1830) in commercials were the major reproductive issues in the investigated dairy herds, followed by poor heat detection at 5.2% (54/1020) in locals and anestrus at 4.7% (86/1830) in commercials. Finally, it is recommended that nationwide studies of reproductive characteristics and diseases be conducted, as well as owner education, to establish effective control measures for reproductive management in Bangladesh's dairy cattle industry.

Keywords: Dairy cows; reproductive disorders; prevalence; artificial insemination; semen.

1. INTRODUCTION

Livestock's contribution to Bangladesh's gross domestic product (GDP) was 1.9% in 2021-22 [1]. Livestock plays a major role in agriculture and livelihoods. Cattle, buffalo, sheep, and goat are mainly considered as livestock animals in Bangladesh. However, dairy farming is an important and potential agricultural sector in Bangladesh. Nearly 85% of the population of the country is engaged in agriculture and livestock sector [2]. In Bangladesh, there are about 24.7 million cattle, 1.5 million buffaloes, 3.7 million sheep, and 26.7 million goats [1] are available as per latest livestock census. Among them the total of 6 million milking cows, 85–90% of them are indigenous and 10–15% are crossbred [3].

Reproduction is a vital factor in determining the efficiency of animal production. In cattle production, good reproductive performance is essential for efficient management and production as a whole. However, successful economy of a dairy farm either large or small scale lies in proper and optimal reproductive rhythm of each individual cow within the normal physiological range. Reproductive and production disorders of dairy cattle significantly reduce their productivity which is of great concern to dairy producers worldwide because most reproductive disorders adversely affect the future fertility. Furthermore, mastitis is a major threat of dairy industry in Bangladesh and its effect on conception rate [4]. Poor reproduction performance is a critical limitation factor. It has been reported that reproductive

disorders are accountable for huge economic losses to dairy farmers in Bangladesh [5]. In European and American dairy herds, about a third of all cows are culled because of reproductive disturbances [6]. The occurrence of various reproductive disorders is increased in Bangladesh due to introduction of intensive cross breeding programme through artificial insemination [6,7]. The occurrence of different reproductive disorders in cows has been reported in Bangladesh by several authors [8,9]. Despite the large number of cattle and their economic importance in Bangladesh, the productivity is low due to the constraints of reproductive disorders, nutrition, poor management, poor performance of local breeds, lack of use of reproductive tools and technologies to manage the disorders. Therefore, the present study was undertaken and designed with the objectives of investigating the prevalence of reproductive disorders and collecting the base line data needed for formulating possible strategies for prevention and control of reproductive disorders.

2. MATERIALS AND METHODS

2.1 Study Area

The study covered four districts (out of 64 districts) in Bangladesh; these were Dhaka, Narayanganj, Gazipur, and Manikganj (Fig. 1). The dairy herds were randomly selected from different Upazila's among these districts from September 2021 to June 2022.



Fig. 1. The map of the study area (Modified from <https://www.google.com.bd/search?> Black circle: 1Dhaka, 2Narayanganj, 3Gazipur, and 4Manikganj)

2.2 Study Design and Animals

Cross-sectional study using questionnaire survey and field visit, was carried out in dairy farms in four districts in Bangladesh. A total of 2850 dairy cows (local 1020; 35.8% and crossbred 1830; 64.2%) were surveyed in 256 local (204; 79.7%) and commercial (52; 20.3%) (Table 1) dairy herds located in Dhaka (local 31; 15.3% and commercial 10; 19.3%), Narayanganj

(local 54; 26.4% and commercial 12; 23%), Gajipur (local 57; 27.9% and commercial 19; 36.7%), and Manikganj (local 62; 30.4% and commercial 13; 25%) (Table 2) were investigated using questionnaire survey and face-to face interviews with the owners. The dairy herds were categorized into small (1-5 cows; 47.7%), medium (6-10 cows; 30.5%), and large (≥ 10 cows; 21.8%) (Table 1).

Table 1. Study areas and categories of dairy herds investigated

Locality	Number of dairy herds investigated			
	Local	Percentage (%)	Commercial	Percentage (%)
Dhaka	31	15.3	10	19.3
Narayanganj	54	26.4	12	23
Gazipur	57	27.9	19	36.7
Manikganj	62	30.4	13	25

Table 2. Frequency table showing the characteristics of the dairy herds visited

Types of dairy Herd	Local	204	79.7%
	Commercial	52	20.3%
Herd Size	Small (1-5)	122	47.7%
	Medium (6-10)	78	30.5%
	Large (≥ 10)	56	21.8%
Dairy Cows	Local cows	1020	35.8%
	Cross-bred	1830	64.2%

2.3 Questionnaire Survey

A total of 256 owners of dairy herds were interviewed face-to-face using close ended questions to ease data processing; minimize variation and improve precision of responses. In the survey, information on reproductive parameters (estrus detection method, reproductive calendar, reproductive records, service methods, source of semen, and breed of semen) as well as reproductive disorders were collected.

2.4 Data Processing and Analysis

Questionnaires were numbered (coded) before processing and the collected data were sorted manually. Data collected on reproductive parameters, prevalence and relative frequency of reproductive disorders were calculated in percentages.

3. RESULTS AND DISCUSSION

A total of 256 questionnaire was obtained to all the dairy herds investigated. Out of the 256 dairy herds visited, 57 were local and 19 were commercial in Gazipur district while, 147 were local and 33 were commercial dairy herds in the other areas (Table 1).

A total of 256 dairy herds (Local 204 and Commercial 52) were examined with a maximum of 47.7% ($n = 122/256$) of small dairy herds categorized against medium and large dairy

herds. Of these, 35.8% ($n=1020/2850$) were locals and 64.2% ($n=1830/2850$) were crossbred dairy cows (Table 1). In Bangladesh, there are about 24.7 million cattle, 1.5 million buffaloes, 3.7 million sheep, and 26.7 million goats [1]. Out of the 6 million milking cows, 85–90% are indigenous and 10–15% are crossbred [3]. Smallholder producers dominate the dairy sector in Bangladesh. More than 70% of the dairy farmers are smallholders and produce around 70–80% of the country's total milk [10]. It is estimated that there are about 1.4 million dairy farms with an average herd size of 1–3 cows [11].

The details on estrus detection methods, reproductive calendar and records of these dairy herds are presented in Table 3. Most of the dairy herds ($n=197$; 76.9%) were observing estrus signs or taking history from the herd owner to detect the estrus of dairy cows rather than calling of a veterinarian and using digital technologies. Moreover, 32.8% ($n=84/256$) of the dairy herds were using reproductive calendar for breeding purposes while only 26.1% ($n=67/256$) were keeping the reproductive records either by log book or a computer. Detection of estrus in cows may often be difficult due to short periods of standing estrus, silent estrus, changing nutritional regimens, environmental temperatures and estrus onset during the late night to early morning hours [12]. There is a report that about 40% of cows remained undetected when they were in estrus in small holding farms in Bangladesh [13].

Table 3. Reproductive aspects of dairy herds reported by the participants

Estrous detection	History/Signs	197	76.9%
	Veterinarian observation	57	22.3%
	Digital technologies (Ultrasound, CCTV camera etc) observation	2	0.8%
Reproductive calendar	Yes	84	32.8%
	No	172	67.2%
Reproductive records	Log book	48	18.7%
	Computerization	19	7.4%
	None	189	73.9%

CCTV: Closed circuit camera

Table 4. Dairy cows service methods reported by the participants

Number dairy herds	Type of dairy herds	AI only	Percentage (%)	NS only	Percentage (%)	AI and NS	Percentage (%)
204	Local	146	71.5	23	11.2	35	17.1
52	Commercial	38	73.0	0	0	14	26.9

AI: Artificial insemination; NS: Natural Service

Table 5. Sources of semen used for insemination of the dairy cows

Type of dairy herds	Number dairy herds	DLS	Percentage (%)	BRAC	Percentage (%)	Lal Teer	Percentage (%)
Local	204	137	67.1	44	21.5	23	11.2
Commercial	52	18	34.6	32	61.5	2	3.8

DLS: Department of Livestock Service; BRAC: Bangladesh Rural Advancement Committee

Table 6. Breeds from which semen used for insemination were sourced

Type of dairy herds	Number dairy herds	Holstein	Percentage (%)	Sahiwal	Percentage (%)	Red Chittagong	Percentage (%)
Local	181	109	60.2	77	37.1	5	2.7
Commercial	52	39	75	10	19.2	3	5.7

Table 7. Types and prevalence of reproductive disorders in the dairy cows

Dairy cows	Breed of cows		Reproductive disorders	Affected cows		Prevalence (%)	
	Local	Crossbred		Local	Crossbred	Local	Crossbred
2850	1020	1830	Anoestrus	88	86	8.6	4.7
			Repeat breeding	28	180	2.7	9.8
			Metritis	9	57	0.8	3.1
			Retained placenta	11	32	1.1	1.7
			Early embryonic death	7	13	0.7	0.7
			Poor heat detection	54	65	5.2	3.5
			Ovarian cyst	0	7	0	0.3
			Uterine prolapsed	0	12	0	0.6
			Vaginal prolapsed	1	7	0.09	0.3
			Still birth	0	12	0	0.6
			Abortion	1	8	0.09	0.4
			Fetal mummification	1	5	0.09	0.2
			Dystocia	19	27	1.9	1.4
			Pyometra	4	9	0.3	0.4
			Ovarian tumor	0	1	0	0.05
			Cervicitis	0	4	0	0.2
			Sulphingitis	0	3	0	0.1
			Ovobarsal adhesion	1	2	0.09	0.1
			Ovarian atrophy	0	3	0	0.1
			Total			309	533

Of the 256 dairy herds in the study, 71.5% (n=146/204) locals used artificial insemination (AI) for the dairy cows reproduction when classified according to service method while, 73% (n=38/52) commercials were using AI other than natural service (NS), and AI and NS (Table 4). Artificial Insemination (AI) is the first-generation reproductive biotechnology that has made a profound contribution to the genetic improvement of the cattle. Failure of cows to become pregnant and the need for repeated AI result to frustration and economic losses of the cattle farmers [14]. Although AI is a widely recognized breeding tool but farmers in Bangladesh are not yet able to adopt it perfectly. This is due to facing some problems while practicing AI at farmers' level. Some of the problems are poor ability to adapt services, low pregnancy rate, poor communication, inability to select the desired breed and to practice AI inappropriate time etc [15,16].

The source of semen for dairy cows reproduction are presented in Table 5. The use of Department of Livestock Service (DLS) semen rate was higher (67.1 %) in locals compared with BRAC (21.5%) and Lal Teer (11.2%). On the other hand, BRAC semen rate was higher 61.5% (n=32/52) in commercials compared with DLS 34.6% (n=18/52) and Lal Teer 3.8% (n=2/52). The present finding is consistent with other investigators where cows inseminated with frozen semen derived from BRAC showed higher (61.6%) pregnancy rate than that of semen derived from DLS (55.0%) [17].

Moreover, the breed of semen of dairy cows reproduction are presented in Table 6. Most of the locals were using Holstein breed 60.2% (n=109/204) semen compared with Sahiwal 37.1% (n=77/204) and Red Chittagong 2.7% (n=5/204) while, 75% (n=39/52), 19.2% (n=10/52) and 5.7% (n=3/52) using in commercials, respectively. The crossbreds and purebreds are mostly in Bangladesh Sindhi, Sahiwal, and Holstein Friesian breeds [18].

Prevalence of reproductive disorders was 21.6% in locals and 31.4% in commercials in total investigated population. Among the disorders, anestrus was 8.6% (n=88/1020) in locals and repeat breeding syndrome (RBS) was 9.8% (n=180/1830) followed by poor heat detection was 5.2% (n=54/1020) in locals and anestrus was 4.7% (n=86/1830) in commercials (Table 7). The other disorders reported in locals included

RBS was 2.7% (n=28/1020), dystocia was (n=19/1020), and retained placenta (RP) was 1.1% (n=11/1020) while, poor heat detection was 3.5% (n=65/1830), metritis was 3.1% (n=57/1830), RP was 1.7% (n=32/1830), and dystocia was 1.4% (n=27/1830) in commercials. In addition, less than 1% of local and crossbred reproductive disorders are detailed in Table 7. Similar prevalence of reproductive disorders in cross-bred dairy cows was reported as post calving anoestrus was 8.6%, delayed puberty was 6.8%, repeat breeding was 5.7%, retained placenta was 4.7%, metritis was 3.9%, dystocia was 3.3%, vaginal prolapse was 2.7%, abortion was 2.1% and uterine prolapse was 1.6% [19]. Contrasting to the present finding, 23.0% prevalence of reproductive diseases was reported in crossbred cows in Chittagong district [20]. Reproductive disorders reported in goats were dystocia (41.2%), abortion (21.8%), mastitis (21.8%), retained placenta (11.8%), and pyometra (3.4%) [21]. Similarly, major reproductive disorders in sheep were dystocia (53.7%), abortion (25.0%), pyometra (7.1%), mastitis (7.1%), and retained placenta (7.1%) (Sultan et al. 2015). The occurrence of anoestrus was 6.79%, which was lower than that (26.52%) observed in South West Ethiopia [22], and 27.73% cases of retained placenta reported in Karan Fries cows [23]. Other researchers reported lower incidence of retained fetal membrane (0.31%, 1.47%, 8.28%, and 7.84% respectively) [24]. The occurrence of metritis was recorded 25.57% in South West Ethiopia [25]. Retained placenta is an important post-parturient problem in cattle farming. The prevalence of RP was 12% even in normal delivery, which was about 63% of the RP on 750 calvings on the Savar dairy farm [8]. The incidence of repeat breeding in the present studies was 3.20% which was lower than the findings in Ethiopia [26]. Similarly, communal use of bulls for natural services is also considered as contributing factor reported 5% repeat breeding cases [21]. The prevalence rate of pyometra 0.2% recorded in this study is lower than the prevalence rate of 8.2% observed in Savar dairy farm [8]. A 47.7% of dairy cows were having at least one of the reproductive problems observed by Benti and Zewdie, (2014) in indigenous Borena breed cows in Borena zone in Southern Ethiopia. Contrasting to the present finding, 39.4% prevalence of reproductive disorders was reported in cross-bred dairy cows at Ullapara upazila under Sirajganj district [19]. Haile et al. [27] recorded 43.07 % of cows were found to be affected either with one or more of reproductive problems.

Variation in the prevalence of reproductive disorders in the dairy industry between studies may be due to variation in dairy herd management and agroclimatic conditions in the regions.

4. CONCLUSION

This study revealed a wide variety of reproductive parameters and reproductive disorders affecting local and commercial dairy industry in Bangladesh. The situation calls for further investigations in both crossbred and local breed of dairy cows to determine causes, economic impact, and appropriate control strategies under different management systems.

ACKNOWLEDGEMENT

The authors would like to thank the Level-4 Semester-1 (2022) students of Animal Science and Veterinary Medicine Faculty, Sher-e-Bangla Agricultural University, Dhaka-1207 for investigating and collecting data by using prescribed questionnaire.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. DLS. Department of Livestock Services: livestock economy at a glance. Ministry of Fisheries and Livestock. Dhaka: Government of the People Republic of Bangladesh. 2021-22.
2. Raha SK. Development of livestock sector: issues and evidences. Eskaton Garden Road, Dhaka: Changing Rural Economy of Bangladesh. Bangladesh Economic Association; 2000.
3. DLS. Department of livestock services: an overview. Ministry of Fisheries and Livestock. Dhaka: Government of the People Republic of Bangladesh. 2012-13.
4. Khokon MS, Azizunnesa A, Islam MM, Chowdhury KB, Rahman ML, Ali MZ. Effect of mastitis on post-partum conception of cross bred dairy cows in Chittagong District of Bangladesh. *J Adv Vet Anim Res.* 2017;4(2):155-60..
5. MAST, MAMYK, MGMR, MRI, MAAK. Reproductive problems of cow at Bangladesh Agricultural University Dairy Farm and possible remedies. *Pak J Biol Sci.* 2005;8(11):1561-7.
6. Faruq MBH. Clinical and abattoir studies on reproductive diseases of cows in Bangladesh [MS thesis]. Mymensingh, Bangladesh: Department of Surgery and Obstetrics, Faculty of Veterinary Science, Bangladesh Agricultural University; 2001.
7. Khair A, Alam MM, Rahman AK, Islam MT, Azim A, Chowdhury EH. Incidence of reproductive and production diseases of cross-bred dairy cattle in Bangladesh. *Bangl J Vet Med.* 2013;11(1):31-6.
8. Shamsuddin M, Alam MGS, Ahmed JU. Reproductive disorder of cross- bred cows. *Bangladesh Vet J.* 1988;22:21-8.
9. Das SC, Rahman MA, Ahmed JU, Alam MG. Reproductive disorders in zebu cows of Tangail milk shed area. *Bd J Anim Sci.* 1995;24:19-25.
10. Uddin MM, Sultana MN, Bruermer B, Peters KJ. Assessing the impact of dairy policies on farm-level profits in dairy farms in Bangladesh: benchmarking for rural livelihoods improvement policy. *J Revi Glob Eco.* 2012;1:124-38.
11. Hemme T, Deeken E, Ramanovich M. IFCN dairy report. Kiel: International Farm Comparison Network (IFCN), Dairy Research Center; 2008.
12. Hasib FM, Reza MM, Alam MM, Azizunnesa TH. Occurrence and risk factors of repeat breeding on household dairy cows of Hathazari in Chattogram. *Bangladesh J Vet Anim Sci.* 2020;8:1.
13. Shamsuddin M, Bhuiyan MM, Sikder TK, Sugulle AH, Chanda PK, Alam MG et al. Constraints limiting the efficiency of artificial insemination of cattle in Bangladesh. *Int Atom Ergol Agcy.* 2001;1220:9-27.
14. Stevenson JS, Call EP, Scoby RK, Phatak AP. Double Insemination and Gonadotrophin releasing hormone treatment of repeat breeding dairy cattle. *J Dairy Sci.* 1990;73(7):1766-72.
15. Alam MGS. National livestock policy and action plan, veterinary services and animal health, FAO office, Dhanmandi, Dhaka, Bangladesh. 2005.
16. Paul AK, Alam MG, Shamsuddin M. Factors that limit first service pregnancy rate in cows at char management of Bangladesh. *Parity.* 2013;215:45-5.
17. Shikder A. Post AI Pregnancy rate of crossbred and zebu cattle using frozen semen at Debhata and Kaligonj Upazilla in

- Satkhira district. MS in Theriogenology Thesis. Department of Surgery and Obstetrics, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh. 2011.
18. Miaz OF, Hossain ME, Hassan MM. Productive and reproductive performance of crossbred and indigenous dairy cows under rural conditions in Comilla, Bangladesh. Univ j zool Rajshahi Univ. 2007;26:67-70.
 19. Sultan A, Islam MR, Yadav RK, Akhter R, Ahmed JU. Prevalence of different reproductive disorders of small ruminants in five upazillas of Mymensingh district. Asian J Med Biol Res. 2015;1: 74-9.
 20. Alam MA, Bhuiyan MMU, Parvin MS, Rahman MM, Bari FY. Prevalence of reproductive diseases and its associated risk factors in crossbred dairy cows. Res Agric Livest Fish. 2014;1(1): 71-9.
 21. Maruf AA, Islam MR, Rahman MM, Bhuiyan MMU, Shamsuddin M. Prevalence of Reproductive Disorders of Dairy Cows in the Chittagong District of Bangladesh. Bangladesh Vet J. 2012;46:11-18.
 22. Bitew M, Shiv P. Study on major reproductive health problems in indigenous and cross breed cows in and around Bedelle, south West Ethiopia. J Anim Vet Adv. 2010;10:723-7.
 23. Satya P. Investigation on health disorder in dairy cattle and buffaloes during pre and post-partum period Phd [thesis] submitted to NDRI. Karnal; 2003.
 24. Shamsuddin M, Bhattacharjee J, Goodger WJ, Momont H, Frank G, Talukdar AK et al. Community-based productivity veterinary service for smallholder dairy farmers in Bangladesh. In: Sustainable improvement of animal production and health. Rome: Food and Agriculture Organization of the United Nations; 2010; 247-53.
 25. Bitew M, Prasad S. Study on major reproductive health problems in indigenous and cross-breed cows in and around Bedelle, south West Ethiopia. J Anim Vet Adv. 2011;10(6):723-7.
 26. Getachew E, Nibret M. Major Reproductive Health Disorders in cross breed Dairy Cows in Ada'a District, East Shoa, Ethiopia. Department of Veterinary Clinical Studies. Ethio Glo Veterinaria. 2014;13(4):444-9.
 27. Haile A, Tsegaye Y, Tesfaye N. Assessment of major reproductive disorders of dairy cattle in urban and per urban area of Hosanna, Southern Ethiopia. Anim Vet Sci. 2014;2:135-41.

© 2022 Islam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91021>