



Assessing of Water Quality of Lake Dogodogo in Burundi Based on Zooplankton Diversity Indices

S. Buhungu ^{a*}, C. Sibomana ^a, L. Niyoyitungiye ^a,
E. Niyikiza ^a, P. Bararunyeretse ^a and S. Ntiharirizwa ^b

^a Department of Biology, Faculty of Sciences, Natural Sciences and Environmental Research Center, University of Burundi, Bujumbura, Burundi.

^b Department of Earth Sciences, Faculty of Sciences, Natural Sciences and Environmental Research Center, University of Burundi, Bujumbura, Burundi.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v21i4547

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95957>

Original Research Article

Received: 18/11/2022

Accepted: 27/01/2023

Published: 22/02/2023

ABSTRACT

A study of zooplankton diversity and water quality of Lake Dogodogo located in West Burundi was carried out from April to July 2021 in six sampling stations, with four selected in the littoral zone and two remaining in pelagic one. The main objective was to assess the water quality using zooplankton diversity indices for a better management of this fishery resource essential for the surrounding populations. The zooplankton sampling was collected twice a month between 8 A.M and 11 A.M each time. The samples were taken vertically using plankton net with a 50 µm of mesh size, 26 cm in diameter and 0.5m depth. We recorded 30 zooplankton species, 19 of rotifers, 9 of copepods and 2 of cladocerans. The number of species varied from one station to another and from the littoral zone to pelagic zone. The abundance of zooplankton was higher in the pelagic zone than the littoral one. We then highlighted species specific for pelagic zone, such as

*Corresponding author: Email: simon.buhungu@ub.edu.bi;

Anuraeopsis fissa, *Brachionus angularis*, *Brachionus calyciflorus*, *Brachionus falcatus* and *Keratella tropica*. These latter are indicators of a high trophic level. According to the diversity indices calculated, the Dogodogo's water quality is moderately polluted by a low level contamination of agricultural origin. Therefore, it is necessary to protect this important resource for fish production purposes, which is essential in protein supply for human life.

Keywords: Aquatic ecosystem; Cibitoke province; zooplankton composition; biodiversity.

1. INTRODUCTION

Agricultural activities carried out by humans, organic matter loads and dissolved substances from domestic effluents contribute to changes of water composition in aquatic environments [1]. Many substances are washed away by drainage water or runoff leaching fertilizers and phytosanitary products and wastewater from urban areas enriched with organic matter.

In rural areas, these waters are subject to agricultural pollution due to pesticides, fertilizers and livestock residues. The long-term result is deterioration in water quality, reduction in diversity and even the disappearance of water bodies [2]. These modifications affect significantly the food webs [3].

For thousands years and especially during the last decade, water engineering has been conducted to ensure a reliable supply of water, limit flood damage, and to avoid or clean up pollution [4]. In the absence of scientific criteria, water quality was assessed using simple criteria. Knowing its origin, a simple visual estimate of its color was enough to declare that the water was drinkable. The physico-chemical criteria of water need to be supplemented by the characterization of biological community indicators to determine the quality of the environment [5]. The use of diversity indices can then be an approach for estimating the biological and ecological quality of an ecosystem through the structure of the community. It can also provide information on pollution levels in an environment [6,7].

Both zooplankton and macroinvertebrates constitute a tool for biological monitoring of aquatic environments. It is one of the first links in the food chain in aquatic environments [8]. Depending on their lifestyle, their reactions to various disturbances as well as the modification of the composition of the water, zooplankton have become an excellent indicator of climate change [9]. It can be used to study the disturbances that can occur in an aquatic ecosystem [10]. Zooplankton holds a key place in

food webs, link primary producers and secondary consumers [11] and serve as an important food source for aquatic organisms, especially fish [12]. Planktivorous fish are in turn hunted by predatory fish. The chain continues until human nutritional needs are met. These organisms are also used for the study of ecological interactions [13].

Regarding the management of aquatic ecosystems, without measuring physico-chemical parameters each time, zooplankton species can be used as a biological indicator of pollution and eutrophication [8,14].

The study of zooplankton helps in the development of an effective strategy for managing fishery resources [15]. The alteration of food states in the water column leads to profound changes in the structure of zooplankton communities. It is made up of organisms that are very sensitive to the variations in their environment conditions, with the relatively short period of time [14,16]. Some species are largely tolerant to certain environmental conditions and are therefore used as indicators of water pollution [17].

The study of taxonomy, distribution and biology of species is an essential tool for the further development of research, but the inventory of aquatic species remains an unfinished task. In tropical regions, the fauna and flora are poorly known. An estimated of 90% of microorganisms present in the environment have not yet been well described [18]. Burundi has adopted in its commitments through the water code, the establishment of legislation in favor of the environment and biodiversity in general and the limitation of water pollution, in particular those of the Lake Tanganyika [19]. Burundi has many lakes that are subject to anthropogenic pressures. Lake Dogodogo is one of the aquatic systems of Burundi located in the northwest of the country in the province of Cibitoke. As per our knowledge, no study has been conducted on its biotic and abiotic components. Lake Dogodogo is a shallow Lake surrounded mostly by fields of rice, maize, and cassava. Thus, its

waters receive different types of agricultural fertilizers from the amendment of crops and pesticides. These substances discharged into the lake change the chemical composition of the water and the animal community's composition living there—resulting in degradation of biodiversity in general and zooplankton in particular. The overall objective of the study was to assess zooplankton diversity and water quality of Lake Dogodogo using zooplankton diversity indices for the sustainable management of this fishery resource, which is very important for the surrounding populations, and finally to have a database for decisions makers and researchers.

2. MATERIALS AND METHODS

2.1 Study area

Lake Dogodogo is located in northwest of Burundi, Cibitoke Province, in the Imbo plain. It is situated between 2° 50' 23.00" South latitude and 29° 05' 52.00" East longitude and 910m of altitude. It is a young reservoir located at 67 km far away from Bujumbura City and in the southern part, at 4 km far away from the chief town of Rugombo commune, along the National Road 5, in right side. This region is the westernmost and has the lowest altitude in Burundi [20].

The Lake occupies an area of 80 hectares with the largest pond having marshy valleys covering nearly 450 hectares. It is characterized by aquatic vegetation formed by water lilies and has a remarkable algal flora and some fish species. The edges of the lake are invaded by *Phragmites mauritianus* and *Typha domingensis* and tall grasses sailing on its water in an unexpected and mysterious way. Lake Dogodogo biodiversity also includes aquatic birds such as ducks [21] and its soil is sandy clay.

2.2 Sampling Stations

A total of six stations were selected depending on their characteristics (Fig. 1) and four stations among them (St 1, St 2, St 3 and St 4) were selected in the littoral zone while the two remaining others (St 5 and St 6) were taken from the pelagic zone. The first station St 1 with

coordinates (S 2.84191°, E 29.09506°, altitude: 912 m) is located in front of the water outlet of the lake through the discharge canal towards Rusizi River. It is chosen to reflect the water quality of the lake at the outlet. The second station St2 with coordinates (S 2.84312°, E 29.09919°, altitude: 913 m) is located to the right side of the station St 1 near the rice fields and was selected to show the impact of irrigation water on zooplankton diversity.

The Stations St 3 and St 4 respectively with coordinates (S 2.83797°, E 29.09956°, altitude: 914 m) and (S 2.83543°, E 29.09529°, altitude: 910 m) are located at a few meters far away from the mouth of the irrigation canal from the rice fields. Station St 4 is located at the left side of St 3 and was chosen near the landing area for small fishing boats. Both Stations St 3 and St 4 were chosen to assess the influence of the nutrient supply in water on zooplankton composition.

The St 5 with coordinates (S 2.83949°, E 29.09763°, altitude: 913 m) and station St 6 with coordinates (2.83689°, E 29.09626°, altitude: 912m) are located in the middle of the lake, especially in the pelagic zone. These two stations were chosen to characterize pelagic zooplankton diversity.

2.3 Sampling

The zooplankton sampling was carried out twice a month over a four-month period, from 4th April to 25th July 2021. Sampling was done between 8 A.M and 11 A.M each time. The samples were taken vertically using plankton net with a 50 µm of mesh size, 26 cm in diameter and 0.5m depth. At each sampling, the plankton net was immersed and then raised up to the surface. The water retained in the tap was allowed to flow and then collected. The zooplankton concentrate was then collected and preserved in a bottle by adding 5ml of 5% formalin solution [17,22,23].

At each station, sampling was repeated three successive times to constitute a composite sample. Six bottles labeled according to the order of the stations were used to preserve the samples and then sent to the laboratory for microscopic analysis.




Fig. 1. Location of Lake Dogodogo and sampling stations

2.4 Observation, Identification and Enumeration of Zooplankton

In the laboratory of Limnology, each zooplankton sample was concentrated to a volume of 50 ml. Zooplankton species were identified by microscopic observation using WF 10 X-18 MM optical microscope. This species identification operation was based on the specific morphological characters observable using different determination keys [24-28]. The individuals of each identified species were also counted using a Burker Turk enumeration cell [17] and the counting effort was fixed at 400 individuals for each inventoried species. Thus, the count rate varied according to species abundance up to 100% of sample for rare species. An extrapolation was then made on total volume of sample, on the one hand, and the volume of filtered water, on the other hand, to assess the densities per liter of the Lake water. The density was calculated using the following relation:

$$D = \frac{1000 * (ni * \frac{50}{AR})}{V}$$

Where,

D: The density (expressed in individuals per liter);

ni: The number of individuals recorded for species i;

AR: sample analysis rate corresponding to ni;

V: volume of filtered Lake Water (ml) [17,29].

2.5 Determination of Diversity Indices

- i. **Simpson's abundance index** was used to assess the imbalance in terms of individual within the population. It is given by the formula:

$$D = \sum (ni / N)^2,$$

Where:

ni: The number of species i,

N: The total number of individuals considering all the species.

If the calculated value is closer to 100%, there is dominance of individuals of a few taxa [23].

- ii. **Shannon-Weaver diversity index** establishes the link between the number of species and the number of individuals in the same ecosystem or the same community [15]. It was calculated according to the following formula:

$$H' = -\sum \left(\frac{ni}{N} \right) \log_2 \left(\frac{ni}{N} \right)$$

Where:

- H'**: The value of the Shannon diversity index,
Σ: The sum of the results obtained for each of the species present,
ni: The number of species i,
N: The total number of individuals of all species,
log₂: The logarithm of base 2.

This index helps knowing the diversity of the species that make up the stands of a given environment [23].

- iii. **Evenness index** was calculated using the formula of Piélou [30]:

$$J = H' / \log_2 S$$

Where:

- H'**: The Shannon-Weaver species diversity index;
log₂ S: H'max,
S: The total number of species in the sample,
J: The Piélou evenness index.

On one hand, the evenness index is derived from the specific diversity index and consists of comparing the diversity H' to its maximum value (log₂N). At the other hand, the evenness index helps in comparing the measured diversity to the theoretical maximum diversity [15].

- iv. **Margalef species richness index** was used to assess the biological diversity of the different stations. This Index is used to estimate the absolute species richness. Its value is obtained by the following formula:

$$RMg = S - 1 / \log (N)$$

Where,

- N**: The number of individuals and
S: The total number of species [31].

The Margalef species richness index is the simplest way to measure biodiversity. It counts

the number of different species in a given area [32].

2.6 Statistical analysis

The Statistical analyzes were performed using IBM SPSS Statistics version 22 and Canoco 5 software and those analyzes includes: One way Analysis of Variance (ANOVA-1) used to test the significance (at p=.05) of the differences between the mean data found for the species identified at the different stations while a Principal Component Analysis (PCA) was applied to all the biotic parameters for establishing the relationship between the biotic parameters and study stations, and to better understand their spatial distribution.

3. RESULTS

3.1 Species Composition of Zooplankton

A total of 30 zooplankton species belonging to three groups of zooplankton have been identified in Lake Dogodogo. Indeed, the rotifers group comprises of 19 species (63.33%), copepods group contains 9 species (30%) whereas cladocerans group contains only 2 species (6.66%). These different species are distributed into 8 families and 16 genera. The most diversified families are Brachionidae with 8 species divided into 4 genera followed by the Cyclopidae with 8 species and the Filinidae family with 4 species. The Trichocercidae, Asplanchnidae and Moinidae families are each represented by two species, while the rest of the families are monospecific like the family of Lecanidae and Synchaetidae. The number of species varies from station to another and the highest species richness was observed at station 4 with 30 species while 25 species were found at station 5. Among the 30 species listed above, only seven species were not identified in all the stations: four species of rotifers, one species of copepods and two species of cladocerans. These are *Brachionus calyciflorus*, *Brachionus plicatilis*, *Brachionus quadridentatus*, *Keratella sp*, *Macrocyclus albidus*, *Moina macrocopa* and *Moina micrura* (Table 1).

3.2 Temporal Variation of Identified Species

The study showed that April had the highest density for all the identified species and in all stations except the species *Brachionus angularis*

whose highest density was recorded in May and July. The months of June and July were characterized by a lower density for almost all species. *Brachionus angularis*, *Brachionus falcatus*, *Polyarthra vulgaris* and *Nauplii* species show high densities in all months and were dominant in the zooplankton population found in Lake Dogodogo (Fig. 2).

3.3 Spatio-temporal Variation in the Abundance of Different Zooplankton Groups

The composition of zooplankton has fluctuated considerably over time. The highest density was recorded at station St6 while the lowest density is recorded at station St2 for both rotifers and copepods. The rainy season (April-May) recorded the highest abundances than the dry season (June-July). Rotifers remained the most abundant over time except for the samples collected on 20th June where copepods were higher than rotifers. As for water fleas, they

remained weakly abundant in space than in time (Fig. 3).

3.4 Spatio-temporal Variation of Zooplankton Diversity

The spatio-temporal variation of the diversity indices of Shannon, Piélou equitability, Simpson and Margalef were given. The highest value of the Shannon diversity index was recorded on 4th April at station St 1 while the lowest value was recorded on 25th July at the same station. Regarding the Piélou equitability index, the highest and lowest values were recorded respectively on 11th July and 25th July at the first station.

The highest and lowest values of the Simpson index were noted respectively on 25th July and 16th May at station St 1. For Margalef diversity index, the highest value was recorded on 20th June while the smallest was recorded on 11th July at station 1 (Table 2).




Fig. 2. Temporal variation in total zooplankton identified




Fig. 3. Spatial (A) and Temporal (B) variation in the abundance of zooplanktonic groups

Table 1. Densities (mean, standard deviation) of species identified at different stations, comparison with one way ANOVA at (p= .05) Tukey's pairwise comparisons

Groups	Families	Species	St1	St2	St3	St4	St5	St6	mean± SD	Sig		
Rotifers	Brachionidae	<i>Brachionus falcatus</i>	69.05	92.73	227.90	206.21	368.01	206.58	195.07±107.32	S		
		<i>Brachionus calyciflorus</i>	2.24	2.24	1.55	0.36	0.00	0.00	1.06±1.07	S		
		<i>Brachionus plicatilis</i>	0.00	0.00	0.00	0.18	0.00	0.23	0.06±0.10	S		
		<i>Brachionus angularis</i>	237.65	128.40	357.82	249.79	495.35	595.11	344.02±174.82	NS		
		<i>Brachionus quadridentatus</i>	0.00	0.18	0.00	0.18	0.00	1.37	0.28±0.53	S		
		<i>Brachionus manjavacas</i>	3.92	6.57	9.22	9.67	9.90	6.43	7.61±2.37	NS		
		<i>Keratella tropica</i>	7.62	1.37	1.87	2.24	7.07	4.01	4.03±2.72	S		
		<i>Keratella sp</i>	7.66	1.37	0.64	0.46	2.24	0.00	2.06±2.85	S		
	Trichocercidae	<i>Anuraeopsis fissa</i>	1.78	2.51	1.46	2.65	0.73	6.89	2.66±2.18	S		
		<i>Trichocera similis</i>	2.69	1.96	9.99	37.38	5.84	39.05	16.15±17.33	S		
	Synchaetidae	<i>Trichocera elongata</i>	14.01	9.72	16.15	9.17	11.91	17.11	13.00±3.29	NS		
		<i>Poyartha vulgaris</i>	76.79	74.23	134.70	98.58	156.49	229.98	128.46±59.38	NS		
	Lecanidae	<i>Lecane mira</i>	0.46	0.87	0.23	2.55	1.09	1.28	1.08±0.82	NS		
		Asplanchnidae	<i>Asplanchna girodi</i>	1.00	3.42	1.55	2.92	3.79	4.70	2.89±1.39	S	
	<i>Asplanchna sp</i>		6.66	6.57	10.25	21.76	7.12	7.30	9.94±5.94	NS		
	Filinidae	<i>Filinia terminalis</i>	55.98	29.61	73.99	18.57	59.85	51.37	48.22±20.47	S		
		<i>Filinia opoliensis</i>	26.46	1.96	3.79	2.01	29.20	34.58	16.33±15.29	NS		
<i>Filinia longiseta</i>		19.16	5.52	29.66	4.97	27.83	15.19	17.05±10.60	NS			
Copepods	Philodinidae	<i>Rotaria sp</i>	4.20	3.65	3.65	3.24	3.01	3.97	3.61±0.44	NS		
	Cyclopoidae	<i>Macrocyclus albidus</i>	0.14	0.32	0.96	0.14	0.00	0.59	0.35±0.35	S		
		<i>Cyclops sp</i>	0.23	0.00	1.19	3.10	1.41	2.92	1.47±1.30	S		
		<i>Diacyclops thomasi</i>	1.78	1.05	5.70	1.55	12.86	7.85	5.13±4.65	S		
		<i>Termocyclops leukarti</i>	22.35	12.82	23.36	15.08	103.27	97.57	45.74±42.58	S		
		<i>Thermocyclops crassus</i>	0.55	0.55	1.87	0.41	16.33	12.68	5.39±7.16	NS		
		<i>Acanthocyclops sp</i>	0.41	2.83	1.78	0.78	0.96	2.65	1.56±1.01	NS		
		<i>Eucyclops macrurus</i>	2.83	1.14	3.24	0.91	15.05	8.67	5.30±5.53	S		
		<i>Eucyclops serrulatus</i>	4.79	1.55	9.40	1.41	20.76	10.99	8.15±7.33	S		
		Nauplii sp.	140.98	126.03	180.50	126.88	269.40	348.68	198.74±91.20	NS		
		Cladocerans	Moinidae	<i>Moina micrura</i>	0.09	0.00	0.00	0.14	0.68	0.09	0.16±0.25	NS
				<i>Moina macrocopa</i>	0.00	0.09	0.00	0.05	0.00	0.05	0.03±0.03	S

NS : non significance with (p> .05) ; S : significance with (p< .05) ; Sig : Significativity ; SD : Standard Deviation

Species such as: *Brachionus falcatus*, *Brachionus calyciflorus*, *Brachionus plicatilis*, *Brachionus quadridentatus*, *Keratella tropica*, *Keratella sp*, *Anuraeopsis fissa*, *Trichocera similis*, *Asplanchna girodi*, *Filinia terminalis*, *Macrocyclus albidus*, *Cyclops sp*, *Diacyclops thomasi*, *Termocyclops leukarti*, *Eucyclops macrurus*, *Eucyclops serratus*, *Moina macrocopa* varied significantly (p<.05). While *Brachionus angularis*, *Brachionus manjavacas*, *Trichocera elongata*, *Polyartha vulgaris*, *Lecane mira*, *Asplanchna sp*, *Filinia opoliensis*, *Filinia longiseta*, *Rotaria sp*, *Thermocyclops crassus*, *Acanthocyclops sp*, *Nauplii sp*, *Moina micrura* did not vary significantly (p >.05)

Table 2. Spatio-temporal variation of Shannon, Piélou, Simpson and Margalef indices

Stations	Dates	04-April	18-April	02-May	16-May	06-June	20-June	11-July	25-July	mean
St1	H'	3.23	2.93	2.86	2.96	1.9	2.62	2.97	1.74	2.65
	J	0.73	0.64	0.66	0.71	0.48	0.31	0.8	0.42	0.59
	D	0.14	0.17	0.17	0.01	0.42	0.27	0.15	0.49	0.23
	RMG	6.7	6.48	7.89	6.55	5.74	8.74	4.14	5.45	6.46
St2	H'	2.8	2.44	2.53	3.08	2.24	2.61	2.05	1.8	2.44
	J	0.61	0.61	0.58	0.73	0.57	0.59	0.51	0.5	0.59
	D	0.19	0.29	0.25	0.14	0.28	0.26	0.34	0.44	0.27
	RMG	7.31	6.18	6.63	6.88	6.51	8.79	5.51	5.25	6.63
St3	H'	2.89	1.73	1.96	2.58	2.56	2.43	2.45	2.6	2.40
	J	0.64	0.4	0.5	0.62	0.58	0.61	0.56	0.67	0.57
	D	0.2	0.48	0.35	0.21	0.26	0.23	0.3	0.22	0.28
	RMG	6.69	5.91	4.44	5.56	7.05	5.13	6.82	5.44	5.88
St4	H'	2.59	2.75	2.29	2.4	2.52	2.67	2.09	2.11	2.43
	J	0.54	0.66	0.59	0.55	0.63	0.63	0.49	0.54	0.58
	D	0.01	0.2	0.26	0.27	0.25	0.25	0.38	0.34	0.25
	RMG	7.8	6.22	5.63	6.39	5.69	7.58	6.16	4.9	6.30
St5	H'	2.66	2.37	2.44	2.83	2.78	2.8	2.57	2.86	2.66
	J	0.58	0.52	0.56	0.63	0.66	0.67	0.68	0.66	0.62
	D	0.2	0.27	0.29	0.19	0.2	0.17	0.25	0.2	0.22
	RMG	6.56	6.65	5.91	6.41	6.23	5.83	4.57	6.16	6.04
St6	H'	2.62	3.04	2.31	2.55	2.6	2.46	2.66	2.42	2.58
	J	0.57	0.66	0.56	0.58	0.62	0.59	0.68	0.61	0.61
	D	0.22	0.16	0.29	0.25	0.22	0.27	0.23	0.3	0.24
	RMG	6.56	7.4	5	5.88	5.5	5.51	4.31	4.88	5.63

H': Shannon's index, *J*: Piélou's equitability index, *D*: Simpson's index, *RMG*: Margalef's index

3.5 Distribution of Zooplankton Species in the Sampled Stations and Their Similarities

A Principal Component Analysis (PCA) was performed to all the biotic parameters to highlight the relationships between the species and the sampled stations. The two axes explained respectively 40.38% and 27.41% of the global distribution of zooplankton species, a total variation of 63.23%. They provide sufficient information on their distribution in sampling stations. The PCA serves in distinguishing of the stations of the pelagic environment (St 5 and St 6) and those of the coastal area (St1, St 2, St 3 and St 4). The two axes selected the species of copepods like *Eucyclops serratus*, *Eucyclops macrurus*, *Diacyclops thomasi* and *Termocyclops leukarti* which are positively correlated to them because of their abundance on the littoral zone. The species *Brachionus angularis* and *Polyarthra vulgaris* are negatively correlated to them. The species *Keratella tropica*, *Brachionus quadridentatus*, *Macrocyclus albidis* and

Acanthocyclops sp are weakly represented on the both axes due to their low abundances (Fig. 4).

An ascending hierarchical classification grouped the stations similar to more than 75% St 1 and St 2, St 3 and St 4 and finally St 5 and St 6 compared to the total zooplankton. This classification also highlighted the coastal stations (St 1, St 2, St 3, St 4) and those of the pelagic zone (St 5, St 6) (Fig. 5).

4. DISCUSSION

4.1 Abundance and Variation of Zooplankton in Lake Dogodogo

During this study, three groups of zooplankton were identified. These groups are Rotifers, Copepods and Cladocera. The rotifers group was the most dominant followed by copepods and finally cladocerans. Similar results were found by Buhungu [17] in Kinyankonge River in Burundi and Fofana [2] in Kaby Lake in Ivory Coast.




Fig. 4. Principal Component Analysis (PCA) of zooplankton species in Lake Dogodogo




Fig. 5. Similarity between sampled stations

In Lake Dogodogo, the rotifers group appeared to be more important than copepods and cladocerans. According to several authors, rotifers tend to dominate in terms of abundance in aquatic environments. Two main factors are mentioned to explain this numerical superiority of rotifers: (i) their opportunistic nature, which allows them to better resist to variations in environmental conditions, (ii) their greater competitiveness in these environments not only because of their food plasticity towards available resources, but also because of their small size, which makes them less vulnerable to predator pressure [13,33]. In Lake Dogodogo, the zooplankton composition fluctuates over time, the evolution seems to be similar between rotifers and copepods. In general, higher zooplankton densities recorded during the rainy season decrease dramatically during the dry season. The decrease in the population and specific richness of zooplankton in a lake system could be explained by the increase in temperature and acidity as well as the decrease in nutrients [34]. The dominance of rotifers in Lake Dogodogo during two seasons is a characteristic of tropical lakes [2,23]. Fluctuations in the Shannon, Pielou and Margalef index revealed certain instability in the structure of the zooplankton community over time.

Indeed, in May, which marks the transition between the wet and dry seasons, the abundance of rotifers has decreased. Species of the Brachionidae and Filinidae families dominate

from April to July. These families are generally represented by the largest number of species belonging to the genera *Brachionus*, *Keratella* and *Anuraeopsis*. Thus, whatever the season, the species *Brachionus angularis*, *Brachionus falcatus*, *Polyarthra vulgaris*, were dominant in the group of rotifers. These results coincide with those found by Tchagnouo [15] of the Ossa lake complex (Ossa and Mwembè) in Cameroon.

During the dry season when temperatures are highest, the density of zooplankton, especially rotifers is low. This decrease in rotifers would be probably due to the variation in environmental conditions, especially pollution. The copepods population tended to approach that of rotifers. The development of copepods observed in dry periods would be linked in part to their ability to accept and withstand widely varying environmental conditions [26,35]. This could also be explained by their possibility of surviving in the state of resting stages where certain species of this group can be transported from one environment to another and thus have a wider distribution area [36].

According to Maier [37], the development time of eggs, larvae and copepodid stages is inversely proportional to temperature. The abundance of zooplankton varied from station to station. This variation is attributed to hydrobiological conditions and trophic status [38]. It can be also due to the speed of the low current and the less turbid waters [39] as well as a high transparency

of the waters, due to the sedimentation of heavy particles in suspension [22]. Station S3 is characterized by a high density of total zooplankton compared to stations S1 and S2 in the littoral zone. This proliferation of zooplankton in this coastal zone is probably due to the enrichment of the water which could contain nitrates and phosphates from rice crops favorable to the development of phytoplankton grazed by zooplankton.

The study of zooplankton was carried out in the littoral zone on the one hand and in the pelagic zone on the other hand. A non-homogeneous distribution of rotifers and copepods between the littoral zone and the pelagic zone was noted. The pelagic zone represented by stations St 5 and St 6 has a high abundance compared to the littoral zone. This abundance could be explained by the existence of stable waters. Similar results were reported by Onana [33] with essentially pelagic species such as *Anuraeopsis fissa*, *Brachionus angularis*, *Brachionus calyciflorus*, *Brachionus falcatus*, *Brachionus leydigii* and *Keratella tropica*. Water fleas did not vary much at the sampled stations. This would be caused by the strong predation pressure towards these Water fleas.

4.2 Assessment of the Water Quality of Lake Dogodogo

Zooplankton organisms are sensitive and very reactive to variations in environmental conditions [40]. As a result, they are used as bioindicators of water quality [38]. The study revealed a high representativeness of rotifers compared to other zooplankton groups. Thus, the presence or absence of species could be an indicator of water pollution [41]. According to Margalef [42], a high representativeness of rotifers in freshwater aquatic environments can be considered as a biological indicator of a high trophic level [2]. According to Pourriot and Hillbricht-Ilkowsk [25] the presence of species of the genus *Brachionus*, indicates hypertrophic water. The zooplankton composition recorded in Lake Dogodogo is similar to the results recorded in other organic pollution environments [33,43,44] mainly the waters of tropical environments [45].

A few diversity indices were used to assess the water quality of Lake Dogodogo. According to Simboura [46], the Pielou equitability index can be used to assess the water quality. These authors mentioned the following water classification categories:

The range $0.77 < E < 0.96$ indicates "very good quality water", $0.46 < E < 0.77$ indicates "good

quality water", $0.30 < E < 0.46$ indicates "average quality water", $0.21 < E < 0.30$ indicates "low quality water" and $0.09 < E < 0.21$ indicates "very poor quality water" [47]. In the present study, the average values of the Pielou index are in the range of 0.57 to 0.62. These values show that the waters of Lake Dogodogo are classified as "good quality waters".

Shannon and Simpson indices were also used to assess the water quality [48] according to the following classification categories: $3 < H'$ (or $6 < D$) indicates clean water; $2 < H' < 3$ (or $3 \leq D \leq 6$) indicates low contamination; $1 < H' < 2$ (or $2 \leq D \leq 3$) indicates moderate contamination and $0 < H' < 1$ (or $2 < D$) indicates water with a high level of pollution. In the present study, the mean values of Shannon and Simpson indices fall within the respective ranges of 2.40-2.66 and 0.22-0.28. Only Shannon index indicates that the waters are of "low contamination level" while the Simpson index reveals high pollution. In general, among the three evaluated indices, two of them indicate that the waters of Lake Dogodogo are of low contamination level.

5. CONCLUSION

The study revealed that the zooplankton of Lake Dogodogo is essentially composed of Rotifers, Copepods and Cladocera. The Rotifera group was the most dominant followed by Copepoda and Cladocera. This zooplankton population is dominated by species belonging to the families Brachionidae and Filinidae characteristic of eutrophic conditions. The study also revealed that some identified species are essentially pelagic and indicative of a high trophic level. Some calculated diversity indices showed that the waters of Lake Dogodogo have a low level of contamination and this contamination is of agricultural origin.

To ensure a good future management of Lake Dogodogo, further research could focus on other taxa of animal or plant organisms living in Lake Dogodogo by including physico-chemical parameters over a long period. As a certain pollution threshold is discovered in the lake, some measures must be taken to protect this ecosystem which has a high potential for producing protein resources for human life.

ACKNOWLEDGEMENTS

The authors are sincerely grateful to the Limnology Laboratory of the Natural Sciences

and Environmental Research Center of the University of Burundi for facilitating the analyzes.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Côté R, Bussi eres D, Desgagn es P. Spatio-temporal distribution of phytoplankton and zooplankton in Lac Saint-Jean (Quebec), a hydroelectric reservoir. *Rev. Science. Water.* 2002; 15(3):597–614. DOI: <https://doi.org/10.7202/705471ar>.
2. Fofana NM, Etile RN, Bi GG. Seasonal distribution of zooplankton in relation to environmental characteristics in Lake Kaby (Bongouanou, Ivory Coast). *J Appl Biosci.* 2019;140:14256–14267. DOI:<https://dx.doi.org/10.4314/jab.v140i1.5>
3. Lacroix G, Danger M. From food webs to the functioning of lake ecosystems: Towards an integration of heterogeneity and complexity. *Rev. Science. Water.* 2008;21(2):155-172. DOI: <https://doi.org/10.7202/018464ar>.
4. Karr JR, Chu EW. Introduction: Sustaining living rivers. In *Assessing the Ecological Integrity of Running Waters.* 2000;1-14. Springer, Dordrecht.
5. Beauger A. Bio-assessment of water quality: establishment of a simplified sampling protocol, based on the collection of benthic macroinvertebrates on the sills of rivers with gravelly bedload. *Biodiversity and Ecology.* Doctoral thesis, University Blaise Pascal - Clermont-Ferrand II; University of Auvergne: - Clermont-Ferrand I. French. 2008.
6. Ravera O. Monitoring of the aquatic environment by species accumulator of pollutants: a review. *J Limnol.* 2001;60(Supp.1):63-78. DOI:<https://doi.org/10.4081/jlimnol.2001.s1.63>.
7. Bandeira B. Ecology of zooplankton communities within two Mediterranean coastal ecosystems: treatment of time series. Doctoral dissertation, University of Toulon ; 2013.
8. Koudenoukpo CZ, Chikou A, Z ebaz e SHT, Mvondo N, Hazoume RU, Houndonougbo PK et al. Zooplanktons and aquatic macroinvertebrates: towards an assembly of bioindicators for better monitoring of aquatic ecosystems in tropical regions. *Int. J. Innov. Appl. Stud.* 2017;20(1):276-287.
9. Richardson AJ. In hot water: zooplankton and climate change. *ICES Journal of Marine Science,* 2008;65(3):279–295. DOI:<https://doi.org/10.1093/icesjms/fsn028>
10. Kisekelwa T, Isumbisho M. Qualitative composition of zooplankton upstream and downstream of the Mururu dam. *Cahiers du CERUKI, New Series.* 2010;40:28-32.
11. Espinasse MB. Distribution and structure of zooplankton communities in two coastal ecosystems. Analysis of the impact of physical and trophic factors on the spatial distributions and size spectra of zooplankton. Doctoral thesis. Aix-Marseille University ; 2012.
12. Ismail AH, Adnan AAM. Zooplankton composition and abundance as indicators of eutrophication in two small man-made lakes. Too much. *Life Sci. Res.* 2016; 27(supp1):31-38. DOI: 10.21315/tlsr2016.27.3.5.
13. Sonya L. Spatial-temporal structure of lake zooplankton: Contribution of currents, predation and resources. Dissertation, University of Quebec in Montreal; 2008.
14. Gazonato NAJ, Silva LC, Saggio AA, Rocha O. Zooplankton communities as eutrophication bioindicators in tropical reservoirs. *Biota Neotrop.* 2014;14(4): 1-12. DOI: <http://dx.doi.org/10.1590/1676-06032014001814>.
15. Tchagnouo JGN, Njin e T, Togouet SHZ, Segnou SCD, Tahir TSM, Tchakont e S et al. Specific diversity and abundance of communities of copepods, cladocerans and rotifers in the lakes of the Ossa complex (Dizangu e, Cameroon). *Physio-Geo. Physical geography and environment.* 2012; 6: 71-93. DOI: <https://doi.org/10.4000/physio-geo.2430>.
16. Mimouni EI-A. Diversity and structure of zooplankton: importance for the conservation of aquatic biodiversity in urban areas. Doctoral thesis, University of Montreal; 2015.
17. Buhungu S, Montchowui E, Sibomana C, Barankanira E, Ntakimazi G, Bonou CA. Zooplankton diversity and abundance of the Kinyankonge River (Bujumbura, Burundi) in relation to environmental variables. *Annals of Agricultural Sciences.* 2019;23 (1):1-20.

18. Leveque C. Biodiversity and management of continental aquatic systems. *Rev. Science. Water*: 1998;11(special issue): 211-221.
DOI: <https://doi.org/10.7202/705339ar>.
19. Burundi Nature Action. Strategy for the limitation of pollution of Lake Tanganyika. Critical ecosystem partnership fund, Bujumbura, Burundi; 2014.
20. Allison E, Paley R, Ntakimazi G, Cowan V, West K. Pollution Control and Other Measures to Protect Biodiversity in Lake Tanganyika (RAF/92/G32); 2000.
21. INECN. Regional plan for implementing the National Strategy and Action Plan on Biodiversity in the Imbo plain. Bujumbura, Burundi ; 2013.
22. Ouattara IN, Ouattara A, Kone T, N'Douba V, Gourene G. Distribution of zooplankton along two small West African coastal basins (Bia and Agnebi; Ivory Coast). *African Agronomy*. 2007;19 (2):197–210.
23. Adandedjan D, Makponse E, Hinvu LC, Laleye P. Preliminary data on the diversity of zooplankton in Lake Nokoué (southern Benin). *J Appl Biosci*. 2017;115:11476-11489.
Available: <https://dx.doi.org/10.4314/jab/v115i1.7>.
24. Dussart B. The copepods of continental waters of Western Europe II. Cyclopoids and Biology. Boubee and Cie, Paris ; 1967.
25. Pourriot R, Hillbricht-Ilkowska A. Research on the biology of some planktonic rotifers. *Bull. Soc. zool*. 1969 ;94 (1):111-118.
26. Amoros C. Practical introduction to the systematics of organisms in French inland waters-5. Crustaceans Cladocera. *Bull. Soc. Linn. Lyons*. 1984;53 (3):72-107; 53(4):120-144.
27. Pourriot R, Francez AJ. Rotifers. Practical introduction to the systematics of organisms in French continental waters. *Bull. Mens. Soc. Linn. Lyons*. 1986;8:1-37.
28. Sharma BK. Rotifers (Rotifera: Eurotatoria) from wetlands of Majuli the largest river island, the Brahmaputra river basin of upper Assam, northeast India. *Check List*. 2014;10(2):292-298.
DOI: 10.15560/10.2.292.
29. Bouzidi MA, Amar Y, Attaoui I, Latrèche A, Benyahia M, Bouguenaya N et al. Copepods, Cladoceres and Rotifers from Lake Sidi M'hamed Benali (north-western Algeria). *Physio-Geo. Physical geography and environment*. 2010;4:69-85.
30. Pielou EC. *Ecological diversity*. Wiley Interscience, New York, USA; 1975.
31. Khalifa N, El-Damhogy KA, Fishar MR, Nasef AM, Hegab MH. Using zooplankton in some environmental biotic indices to assess water quality of Lake Nasser, Egypt. *Int. J. Fisheries Aquat. Stud*. 2015;2(4):281-289.
32. Zhushi Etemi F, Çadraku H, Bytyçi A, Kuçi T, Desku A, Ymeri P et al. Correlation between physical and chemical parameters of water and biotic indices: The case study the White Drin River basin, Kosovo. *Journal of Water and Land Development*. 2020; 46(VII–IX): 229–241.
DOI: 10.24425/jwld.2020.134585.
33. Mamert OF, Hubert ZTS, Ernest K, Tchatcho N, Lié N, Siméon T. Influence of municipal and industrial pollution on the diversity and the structure of benthic macro-invertebrate community of an urban river in Douala, Cameroon. *J.Bio. Approx. Science*. 2016;8(6):120-133.
34. Okogwu OI. Seasonal variations of species composition and abundance of zooplankton in Ehoma Lake, a floodplain lake in Nigeria. *Rev. Biol. Too much*. 2009;58(1):171-182.
DOI: 10.15517/rbt.v58i1.5202
35. Dussart B. Calanoid copepod crustaceans from African inland waters. *Crustaceana*. 1989; Supplement (15):III-205.
36. El Khalki A, Gaudy R, Moncef M. Study of seasonal variations in the population of copepods in the Oum Er Rbia estuary (Atlantic coast of Morocco): impact of urban pollution in the city of Azemmour. *Marine Life (Marseilles)*. 2004;14 (1-2):19-29.
37. Maier G. The effect of temperature on the development times of eggs, naupliar and copepodite stages of five species of cyclopoid copepods. *Hydrobiology*. 1989;184(1):79-88.
38. Dolbeth M, Cusson M, Sousa R, Pardal MA. Secondary production as a tool for better understanding of aquatic ecosystems. *Can. J. Fish. Aquat. Science*. 2012;69 (7):1230-1253.
39. Richardson JS, Mackay RJ. Lake outlets and the distribution of filter feeders: an assessment of hypotheses. *Oikos*. 1991; 370-380.
40. Harris R, Wiebe P, Lenz J, Skjoldal HR, Huntley M. *ICES Zooplankton Methodology Manual*. Academic Press, San Diego, USA; 2000.

41. Zirirane DD, Bagalwa JJ, Isumbisho M, Mulengezi M, Mukumba I, Bora M et al. Comparative evaluation of the A pollution of the Kahuwa and Mpungwe rivers by the use of benthic macroinvertebrates. *Vertigo. The Electronic Journal of Environmental Sciences*. 2014; 14(3). DOI: 10.4000/vertigo.15365.
42. Margalef R. *Limnologia*. Omega, Barcelona; 1983.
43. Véronique L. Relations between planktonic populations and the physico-chemistry of the waters of the Scarpe and the Petite Sensée in the region of Douai (North). *Memory. Pierre and Marie Curie University*; 1997.
44. Zébazé TSH, Njiné T, Kemka N, Nola M, Foto Menbohan S, Monkiedje A et al. Spatial and temporal variations in the richness and abundance of rotifers (Brachionidae and Trichocercidae) and water fleas in a small eutrophic artificial lake located in a tropical zone. *Rev. Science. Water*. 2005;18 (4):485–505.
45. Branco CW, Rocha MIA, Pinto GF, Gômara GA, Filippo RD. Limnological features of Funil Reservoir (RJ, Brazil) and indicator properties of rotifers and cladocerans of the zooplankton community. *Lakes and Reservoirs: Research and Management*. 2002;7 (2):87-92. Available:<https://doi.org/10.1046/j.1440-169X.2002.00177.x>
46. Simboura N, Panayotidis P, Papathanassiou E. A synthesis of the biological quality elements for the implementation of the European Water Framework Directive in the Mediterranean ecoregion: the case of Saronikos Gulf. *Ecological Indicators*. 2005;5 (3):253–266. DOI:10.1016/j.ecolind.2005.03.006.
47. Goffart A. Development of the composition index as part of the phytoplankton indicator composition indicesn *Phytoplankton in coastal waters – bibliographic summary*, University of Liège Belgium and Stareso Calvi Corse; 2010.
48. Javaid AS, Ashok KP. Application of diversity indices to crustacean community of Wular Lake, Kashmir Himalaya. *Int. J Biodivers. Conserv*. 2013;5(6):311-316. DOI: 10.5897/IJBC2013.0567.

© 2023 Buhungu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95957>